

α -Alkylation of Carbonyl Compounds by Direct Addition of Alcohols to Enol Acetates**

Yoshihiro Nishimoto, Yoshiharu Onishi, Makoto Yasuda, and Akio Baba*


The catalytic α -alkylation of carbonyl compounds has contributed remarkably to the development of organic synthesis, and the reaction of enolate derivatives with alkyl electrophiles is a powerful alkylation method.^[1] Alcohols are appealing electrophiles, as they are plentiful and readily synthesized; however, the direct use of alcohols is effectively prevented by the fact that the hydroxy group is a particularly poor leaving group, and it often tends to decomposition of catalysts and active intermediates.^[2,3] The α -alkylation of monocarbonyl compounds using an alcohol remains problematic whereas many research groups have reported the direct alkylation of 1,3-dicarbonyl compounds.^[4] The transfer hydrogenation method has achieved some direct α -alkylations of ketones, although this is only applicable to primary alcohols and requires a strong base to generate the enolate species.^[5] Hidai, Uemura et al. published a ruthenium-catalyzed system, which was limited to the reactions of 1-arylpropargylic alcohols, in the presence of an excess of ketone.^[6] Methods using metal enolates suffer from decomposition of the enolates by the hydroxy group, critically narrowing the scope of available alcohol substrates;^[7] furthermore, the α -alkylation of aldehydes has not been investigated to the degree that ketones have.^[8] Herein, we report a synthesis of α -alkylated carbonyl compounds from enol acetates and alcohols, catalyzed by InI_3 , GaBr_3 , or FeBr_3 , in which the α -alkylation of not only ketones but also of aldehydes has been successfully achieved. Furthermore, the exploitation of enol acetates as readily available, stable, and easily-handled enolate reagents enhances the practicality of this α -alkylation method.^[9] We employed Lewis acids to selectively activate alcohols and thereby suppress side reactions, such as transesterification.

We recently found that the moderate Lewis acidity of indium trihalide effectively promoted the direct coupling of alcohols with nucleophiles, such as allyl and alkenyl sila-

nes;^[3,4e] these results suggested that indium trihalide selectively interacts with hydroxy groups in the presence of other oxygen-containing moieties such as carbonyl groups.

As a model reaction, 1-phenylethanol (**1a**) and 2-propenyl acetate (**2a**) were heated to reflux in 1,2-dichloroethane in the presence of 5 mol % indium trihalide to afford the desired product **3aa** in satisfactory yields, of which InI_3 gave the highest (83 %; Table 1, entries 1–3). By contrast, the more

Table 1: Effects of catalyst and solvent in the reaction of 1-phenylethanol (**1a**) with isopropenyl acetate (**2a**).^[a]



Entry	Catalyst	Yields [%]		
		3 aa	4	5
1	InCl_3	60	0	0
2	InBr_3	70	0	0
3	InI_3	83	0	0
4	$\text{In}(\text{OTf})_3$	25	0	0
5	GaCl_3	62	0	0
6	GaBr_3	73	0	0
7	FeBr_3	77	0	0
8	FeCl_3	48	0	0
9	FeBr_2	38	31	31
10	FeCl_2	0	17	66
11	$\text{BF}_3 \cdot \text{OEt}_2$	0	46	43
12	AlCl_3	0	10	27
13	TiCl_4	0	34	0
14	$\text{Sc}(\text{OTf})_3$	25	0	0
15	RuCl_3	0	6	35
16	$\text{Cu}(\text{OTf})_2$	7	0	0
17 ^[b]	CH_3COOH	0	0	0
18 ^[c]	InI_3	55	0	0
19 ^[d]	InI_3	50	0	0
20 ^[e]	InI_3	64	30	6

[a] Reaction conditions: **1a** (1 mmol), **2a** (2 mmol), catalyst (5 mol %), $\text{ClCH}_2\text{CH}_2\text{Cl}$ (2 mL), 83°C , 2 h. [b] 99% recovery of **1a**. [c] $\text{CF}_3\text{C}_6\text{H}_5$ used instead of $\text{ClCH}_2\text{CH}_2\text{Cl}$. [d] Toluene used instead of $\text{ClCH}_2\text{CH}_2\text{Cl}$. [e] CH_2Cl_2 used instead of $\text{ClCH}_2\text{CH}_2\text{Cl}$; 40°C .

Lewis acidic $\text{In}(\text{OTf})_3$ was less effective, giving only a 25 % yield of the α -alkylated product (Table 1, entry 4). Examination of various Lewis acids revealed that gallium and iron halides also had high catalytic abilities (Table 1, entries 5–10); however, representative strong Lewis acids such as $\text{BF}_3 \cdot \text{OEt}_2$, AlCl_3 , and TiCl_4 all promoted transesterification (**4**) and/or dimerization (**5**) of the parent alcohol, with no formation of

[*] Dr. Y. Nishimoto, Y. Onishi, Dr. M. Yasuda, Prof. Dr. A. Baba
Department of Applied Chemistry
Center for Atomic and Molecular Technologies (CAMT)
Graduate School of Engineering, Osaka University
2-1, Yamada-oka, Suita, Osaka University (Japan)
Fax: (+81) 6-6879-7387
E-mail: baba@chem.eng.osaka-u.ac.jp

[**] This work was supported by Grant-in-Aid for Scientific Research on Priority Areas (No. 18065015, “Chemistry of Concerto Catalysis” and No. 20036036, “Synergistic Effects for Creation of Functional Molecules”) and for Scientific Research (No. 19550038) from Ministry of Education, Culture, Sports, Science and Technology (Japan).

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.200904069>.

the desired product **3aa** (Table 1, entries 11–13). $\text{Sc}(\text{OTf})_3$, RuCl_3 , and $\text{Cu}(\text{OTf})_2$, which are known to catalyze the coupling reaction of alcohols with 1,3-dicarbonyl compounds, was a poor catalyst (Table 1, entries 14–16);^[4n] acetic acid, generated in situ, resulted in no reaction (Table 1, entry 17). Trifluorotoluene, toluene, and dichloromethane were also tested as the solvent and gave satisfactory results (Table 1, entries 18–20).^[10] After analyzing these results, InI_3 , GaBr_3 , and FeBr_3 were taken forward as promising catalysts for further investigations.

We then assessed the scope and limitations of the alcohols **1** and enol acetates **2** (Table 2). In almost all cases, InI_3 , GaBr_3 , and FeBr_3 showed similar catalytic activity. Enol acetates **2b**, **2c**, and **2d**, bearing one or no substituents at the 2-position, gave high yields (Table 2, entries 1–3); disubstituted acetate **2e** afforded ketone **3ae**, including formation of a new quaternary carbon center, although the yields were poorer because of steric hindrance (Table 2, entry 4). Enol acetate **2f**, which was derived from an unsymmetrical ketone, afforded the corresponding product **3af** with no isomerization of the enol acetate (Table 2, entry 5). These results showed that the reaction system had a wide scope of applicable enol acetates. Various types of alcohols were also examined and found to be exploitable. Secondary benzylic alcohol **1b**, bearing an electron-donating group, gave the product **3ba** almost exclusively, whereas the electron-withdrawing chloro substituent decreased the yield (ca. 40%; Table 2, entries 6 and 7). Tertiary benzylic alcohol **1d** was found to react at room temperature, furnishing **3dd** in 74–79% yield (Table 2, entry 8). Allylic, propargylic, and ferrocenyl alcohols **1e–1g** (Table 2, entries 9–11) gave satisfying results, except in the cases of propargylic alcohol **1f** and ferrocenyl alcohol **1g** catalyzed by FeBr_3 (Table 2, entries 10, 11). By contrast, primary benzylic alcohol **1h** was catalysed solely in the presence of

Table 2: Testing the scope of the reaction using various types of alcohols and enol acetates.^[a]

$\text{ROH } \mathbf{1} + \text{Enol Acetate } \mathbf{2} \xrightarrow[\text{ClCH}_2\text{CH}_2\text{Cl, 83 } ^\circ\text{C, 2 h}]{\text{catalyst (5 mol\%)}} \text{Product } \mathbf{3}$					
Entry	1	2	Product 3	Catalyst	Yield [%] ^[b]
1		1a 2b (69:31)		InI_3	80 (60:40)
				GaBr_3	86 (60:40)
				FeBr_3	92 (62:38)
2		1a 2c		InI_3	86 (55:45)
				GaBr_3	75 (59:41)
				FeBr_3	80 (53:47)
3		1a 2d		InI_3	92
				GaBr_3	98
				FeBr_3	92
4		1a 2e		InI_3	28
				GaBr_3	47
				FeBr_3	45
5		1a 2f (67:33)		InI_3	86 (58:42)
				GaBr_3	94 (57:43)
				FeBr_3	81 (56:44)
6		1b 2a		InI_3	97
				GaBr_3	97
				FeBr_3	82
7		1c 2a		InI_3	43
				GaBr_3	38
				FeBr_3	37
8 ^[c]		1d 2d		InI_3	75
				GaBr_3	79
				FeBr_3	74
9		1e 2d		InI_3	82
				GaBr_3	89
				FeBr_3	77
10 ^[d]		1f 2a		InI_3	96
				GaBr_3	74
				FeBr_3	0
11 ^[e]		1g 2a		InI_3	82
				GaBr_3	77
				FeBr_3	26
12		1h 2a		InI_3	40
				GaBr_3	0
				FeBr_3	0
13 ^[f]		1i 2d		InI_3	64
				GaBr_3	69
				FeBr_3	67
14 ^[g]		1j 2a		InI_3	99
				GaBr_3	99
				FeBr_3	99
15 ^[h]		1k 2d		InI_3	62
				GaBr_3	63
				FeBr_3	68


[a] Reaction conditions: **1** (1 mmol), **2** (2 mmol), catalyst (5 mol%), $\text{ClCH}_2\text{CH}_2\text{Cl}$ (2 mL), 83 °C.

[b] Diastereomeric ratio is shown in parentheses. [c] CH_2Cl_2 (2 mL), room temperature, **2d** (3 mmol) was slowly added over 30 min. [d] MeCN (2 mL) used instead of $\text{ClCH}_2\text{CH}_2\text{Cl}$; room temperature. [e] **2a** (5 mmol), MeCN (2 mL) used instead of $\text{ClCH}_2\text{CH}_2\text{Cl}$; 81 °C. [f] **2d** (5 mmol), MeCN (2 mL) used instead of $\text{ClCH}_2\text{CH}_2\text{Cl}$; room temperature. **1i** was added slowly over 30 min. [g] 10 mol % catalyst used. [h] MeCN (2 mL) used instead of $\text{ClCH}_2\text{CH}_2\text{Cl}$; 30 min, 81 °C.

InI₃ to afford the adduct **3ha** in 40 % yield (Table 2, entry 12). These results suggested that the mechanism went via a carbocation intermediate, and that InI₃ could be a more general catalyst than either GaBr₃ or FeBr₃. The reaction of alcohol **1i**, which bears a thiophene moiety, proceeded successfully without deactivation of the metal halide catalyst by the sulfur atom (Table 2, entry 13). 1-Adamantanol **1j** gave the desired product **3ja** in quantitative yield (Table 2, entry 14), although the simple aliphatic alcohol, 2-methyl-4-phenyl-2-butanol, gave no product. Hemiacetal **1k** was effectively transformed into the desired ketone **3kd** without ring cleavage (Table 2, entry 15).

We then focused on the reaction of an aldehyde-derived enol acetate because the reported precedent was crucially limited to a combination of highly reactive 1,1-diarylmethanols.^[8] The reaction of heptanal-derived enol acetate **6a** with 1-phenylethanol **1a** provided the α -alkylated aldehyde **7aa** in high yields (Table 3, entry 1). Other benzylic, allylic, and propargylic alcohols smoothly furnished their corresponding aldehydes, **7ba**, **7ea**, and **7fa** (Table 3, entries 2–4). As for the scope of enol acetates, the reactions using isovaleraldehyde and acetaldehyde-derived enol acetates **6b** and **6c** proceeded successfully (Table 3, entries 5, 6). Furthermore, direct isolation of the aldehyde products is indicative of the mildness of these Lewis acid catalysts.

A plausible reaction mechanism is given in Scheme 1. In the cases of InI₃, GaBr₃, and FeBr₃, the metal halides (MtX₃) are weakly coordinated by oxygen atoms on either the alcohol



Scheme 1. Proposed reaction mechanism.

(**1**) or carbonyl moieties (enol acetate **2**, product **3**, or acetic acid). When MtX₃ interacts with alcohol **1**, a carbocation (R⁺) is generated.^[11] The electrophilic R⁺ is subsequently attacked by enol acetate **2** to afford product **3** with the expulsion of acetic acid and regeneration of MtX₃.^[12] By contrast, Lewis acids which have a high oxophilicity, such as BF₃·OEt₂, AlCl₃, TiCl₄, and In(OTf)₃, are too strongly coordinated by some oxygen moieties to act as a catalyst. Therefore, tolerance of protic conditions and a moderate Lewis acidity of MtX₃ are the particularly important factors in this system.

In summary, we have developed a new procedure for the α -alkylation of aldehydes and ketones that involves the direct use of alcohols and enol acetates in the presence of either InI₃, GaBr₃, or FeBr₃ catalysts. This reaction is applicable to various types of alcohols and enol acetates, and the synthesis of α -alkylated aldehydes is particularly noteworthy. Further investigation to elucidate the reaction mechanism is underway.

Table 3: Reactions using aldehyde-derived enol acetates **6**.^[a]

Entry	1	6	Product	Catalyst	Yield [%]
1				InI ₃ GaBr ₃ FeBr ₃	91 (55:45) 75 (60:40) 84 (61:39)
2				InI ₃ GaBr ₃ FeBr ₃	78 (60:40) 79 (63:37) 78 (66:34)
3				InI ₃ GaBr ₃ FeBr ₃	76 (55:45) 59 (54:46) 57 (58:42)
4 ^[b]				InI ₃ GaBr ₃	74 (54:46) 88 (53:47)
5				InI ₃ GaBr ₃ FeBr ₃	28 (58:42) 54 (56:44) 49 (62:38)
6				InI ₃ GaBr ₃ FeBr ₃	47 16 40

[a] Reaction conditions: **1** (1 mmol), **6** (5 mmol), catalyst (5 mol%), ClCH₂CH₂Cl (2 mL), 83 °C, 1 h. [b] MeCN was used instead of ClCH₂CH₂Cl.

Experimental Section

Typical procedure for the reaction of **1a** with **2a** (Table 1): **1a** (1 mmol) was added to a mixture of catalyst (0.05 mmol) and **2a** (2 mmol) in 1,2-dichloroethane (2 mL) under an inert nitrogen atmosphere. The reaction mixture was stirred at 83 °C for 2 hours, then allowed to cool before diethyl ether (10 mL) and saturated NaHCO₃aq (10 mL) were added. The solution was extracted with diethyl ether, the organic layer dried over MgSO₄, and the solvent removed under reduced pressure to afford the crude product, which was analyzed by NMR spectroscopy.

Received: July 23, 2009

Revised: August 29, 2009

Published online: October 28, 2009

Keywords: alcohols · alkylations · enol acetates · homogeneous catalysis · Lewis acids

- [1] For a review of the α -alkylation of carbonyl compounds, see: a) D. Caine in *Comprehensive Organic Synthesis*, Vol. 9 (Eds.: B. M. Trost, I. Fleming), Oxford, New York, **1991**, pp. 1–63; b) M. T. Reetz, *Angew. Chem.* **1982**, 94, 97–109; *Angew. Chem. Int. Ed. Engl.* **1982**, 21, 96–108.
- [2] a) K.-G. Chung, Y. Miyake, S. Uemura, *J. Chem. Soc. Perkin Trans. 1* **2000**, 15–18; b) M. Rubin, V. Gevorgyan, *Org. Lett.* **2001**, 3, 2705–2707; c) F. Ozawa, H. Okamoto, S. Kawagishi, S.

- Yamamoto, T. Minami, M. Yoshifuji, *J. Am. Chem. Soc.* **2002**, *124*, 10968–10969; d) M. R. Luzung, F. D. Toste, *J. Am. Chem. Soc.* **2003**, *125*, 15760–15761; e) G. W. Kabalka, G. Dong, B. Venkataiah, *Org. Lett.* **2003**, *5*, 893–895; f) S. H. Kim, C. Shin, A. N. Pae, H. Y. Koh, M. H. Chang, B. Y. Chung, Y. S. Cho, *Synthesis* **2004**, 1581–1584; g) Y. Kayaki, T. Koda, T. Ikariya, *Eur. J. Org. Chem.* **2004**, 4989–4993; h) H. Tsukamoto, M. Sato, Y. Kondo, *Chem. Commun.* **2004**, 1200–1201; i) S. K. De, R. A. Gibbs, *Tetrahedron Lett.* **2005**, *46*, 8345–8350; j) M. Georgy, V. Boucard, J.-M. Campagne, *J. Am. Chem. Soc.* **2005**, *127*, 14180–14181; k) G. W. Kabalka, M.-L. Yao, S. Borella, *Org. Lett.* **2006**, *8*, 879–881; l) Y. Kuninobu, E. Ishii, K. Takai, *Angew. Chem.* **2007**, *119*, 3360–3363; *Angew. Chem. Int. Ed.* **2007**, *46*, 3296–3299; m) G.-W. Wang, Y.-B. Shen, X.-L. Wu, *Eur. J. Org. Chem.* **2008**, 4999–5004; n) W. Huang, Q. Shen, J. Wang, X. Zhou, *J. Org. Chem.* **2008**, *73*, 1586–1589; o) G. Chen, Z. Wang, J. Wu, K. Ding, *Org. Lett.* **2008**, *10*, 4573–4576.
- [3] For previous reports from our group, see: a) M. Yasuda, T. Saito, M. Ueba, A. Baba, *Angew. Chem.* **2004**, *116*, 1438–1440; *Angew. Chem. Int. Ed.* **2004**, *43*, 1414–1416; b) T. Saito, M. Yasuda, A. Baba, *Synlett* **2005**, 1737–1739; c) T. Saito, Y. Nishimoto, M. Yasuda, A. Baba, *J. Org. Chem.* **2006**, *71*, 8516–8522; d) A. Baba, M. Yasuda, Y. Nishimoto, T. Saito, Y. Onishi, *Pure Appl. Chem.* **2008**, *80*, 845–854; e) Y. Nishimoto, M. Kajioka, T. Saito, M. Yasuda, A. Baba, *Chem. Commun.* **2008**, 6396–6398.
- [4] For selected recent reports, see: a) K. Manabe, S. Kobayashi, *Org. Lett.* **2003**, *5*, 3241–3244; b) M. Kimura, R. Mukai, N. Tanigawa, S. Tanaka, Y. Tamaru, *Tetrahedron* **2003**, *59*, 7767–7777; c) M. Noji, T. Ohno, K. Fuji, N. Futaba, H. Tajima, K. Ishii, *J. Org. Chem.* **2003**, *68*, 9340–9347; d) H. Kinoshita, H. Shinokubo, K. Oshima, *Org. Lett.* **2004**, *6*, 4085–4088; e) M. Yasuda, T. Somyo, A. Baba, *Angew. Chem.* **2006**, *118*, 807–810; *Angew. Chem. Int. Ed.* **2006**, *45*, 793–796; f) M. Rueping, B. J. Nachtsheim, A. Kuenkel, *Org. Lett.* **2007**, *9*, 825–828; g) J. Kischel, K. Mertins, D. Michalik, A. Zapf, M. Beller, *Adv. Synth. Catal.* **2007**, *349*, 865–870; h) M. Noji, Y. Konno, K. Ishii, *J. Org. Chem.* **2007**, *72*, 5161–5167; i) W. Huang, J. Wang, Q. Shen, X. Zhou, *Tetrahedron Lett.* **2007**, *48*, 3969–3973; j) K. Motokura, N. Nakagiri, T. Mizugaki, K. Ebitani, K. Kaneda, *J. Org. Chem.* **2007**, *72*, 6006–6015; k) R. Sanz, D. Miguel, A. Martinez, J. M. Alvarez-Gutierrez, F. Rodriguez, *Org. Lett.* **2007**, *9*, 2027–2030; l) S. Shirakawa, S. Kobayashi, *Org. Lett.* **2007**, *9*, 311–314; m) P. N. Liu, Z. Y. Zhou, C. P. Lau, *Chem. Eur. J.* **2007**, *13*, 8610–8619; n) S. A. Babu, M. Yasuda, Y. Tsukahara, T. Yamauchi, Y. Wada, A. Baba, *Synthesis* **2008**, 1717–1724.
- [5] G. Guillena, D. J. Ramón, M. Yus, *Angew. Chem.* **2007**, *119*, 2410–2416; *Angew. Chem. Int. Ed.* **2007**, *46*, 2358–2364.
- [6] Y. Nishibayashi, I. Wakiji, Y. Ishii, S. Uemura, M. Hidai, *J. Am. Chem. Soc.* **2001**, *123*, 3393–3394.
- [7] a) P. Vicennati, P. G. Cozzi, *Eur. J. Org. Chem.* **2007**, 2248–2253; b) P. Rubenbauer, T. Bach, *Tetrahedron Lett.* **2008**, *49*, 1305–1309; c) M. Yoshimatsu, T. Otani, S. Matsuda, T. Yamamoto, A. Sawa, *Org. Lett.* **2008**, *10*, 4251–4254.
- [8] P. G. Cozzi, F. Benfatti, L. Zoli, *Angew. Chem.* **2009**, *121*, 1339–1342; *Angew. Chem. Int. Ed.* **2009**, *48*, 1313–1316.
- [9] Only one reaction of 1-naphthylethanol with 1-tetralone-derived enol acetate has been reported: M. Noji, T. Ohno, K. Fuji, N. Futaba, H. Tajima, K. Ishii, *J. Org. Chem.* **2003**, *68*, 9340–9347.
- [10] The screening of solvents is summarized in the Supporting Information.
- [11] The reaction of (*R*)-(+)-phenylethanol (*R*)-**1a** with isopropenyl acetate **2a** in the presence of either InI_3 , GaBr_3 , or FeBr_3 gave the racemic product **3aa**. Experimental details are provided in the Supporting Information.
- [12] T. Mukaiyama, T. Izawa, K. Saigo, *Chem. Lett.* **1974**, 323–326.