

Bridging Binding Modes of Phosphine-Stabilized Nitrous Oxide to $\text{Zn}(\text{C}_6\text{F}_5)_2$ **

Rebecca C. Neu, Edwin Otten, and Douglas W. Stephan*

In 1969, Armor and Taube formulated $[\text{Ru}(\text{NH}_3)_5(\text{N}_2\text{O})]^{2+}$ as the first example of a metal complex of nitrous oxide.^[1] Subsequent studies have supported this formulation with spectroscopic and computational data.^[2] Since then, the interactions of nitrous oxide with transition metals have been shown to play important roles across the discipline. For example, in inorganic synthetic chemistry, reactions of N_2O with transition metal species have been shown to result in oxidation of low-valent metal centers,^[3] insertion of O into metal–carbon or metal–hydride bonds,^[4] and very recently, O-atom transfer to a Ni–carbene complex.^[5] In addition, reactions of N_2O have led to metal mediated N–N bond cleavage^[6] and hydrogenation yielding N_2 and H_2O .^[7] Applications to organic synthesis have recently exploited (transition metal catalyzed) N_2O oxidations of organic substrates.^[8] As a component of the global nitrogen cycle, N_2O is produced and consumed by anaerobic bacteria in denitrification processes that convert NO_3^- or NO_2^- to gaseous products.^[9] The four enzymes that are sequentially involved contain Mo, Fe, and Cu centers in their active sites, of which the latter is required for the last step of N_2O reduction.^[9a] In these nitrous oxide reductases, an unusual Cu_4S cluster is responsible for the conversion of N_2O to N_2 and H_2O ,^[10] and functional synthetic analogues have recently been prepared.^[11] In the field of heterogeneous catalysis, various systems containing transition metals have been developed that decompose N_2O , but these invariably require high temperatures.^[12]

Investigations into the conversion of N_2O to less harmful chemicals have been fueled recently by the realization that N_2O contributes to global warming and stratospheric ozone destruction.^[13] In all the cases mentioned above, the inference of metal– N_2O interactions is clear. Nevertheless, the nature of that interaction remains unknown.

We have recently reported the synthesis of the N_2O complexes $[\text{tBu}_3\text{PN}_2\text{OB}(\text{C}_6\text{F}_5)_2(\text{Ar})]$ ($\text{Ar} = \text{C}_6\text{F}_5$, Ph),^[14] derived from the reaction of the corresponding “frustrated Lewis pairs” and N_2O . Herein, we describe the exploitation of

the reactivity of related main group species to prepare Zn complexes incorporating the $\text{tBu}_3\text{PN}_2\text{O}$ fragment.^[15] These species exhibit two unique bridging modes of the phosphine-stabilized N_2O fragment with the transition metal atoms.

The reactions of $[\text{tBu}_3\text{PN}_2\text{OB}(\text{C}_6\text{F}_5)_3]$ ^[14] with the toluene adduct of $\text{Zn}(\text{C}_6\text{F}_5)_2$ ($\text{tol}\cdot\text{Zn}(\text{C}_6\text{F}_5)_2$) were explored. NMR data for reaction mixtures containing up to 5 equivalents of $\text{tol}\cdot\text{Zn}(\text{C}_6\text{F}_5)_2$ showed no discernible reaction, although resonances for the components were slightly broadened. Diffusion of pentane into a CH_2Cl_2 solution resulted in the precipitation of a mixture of two different types of crystals. Manual separation and subsequent NMR analysis showed these to be the starting material $[\text{tBu}_3\text{PN}_2\text{OB}(\text{C}_6\text{F}_5)_3]$ and a new $\{\text{Zn}(\text{C}_6\text{F}_5)_2\}$ -containing compound, **4**, that is silent in the ^{11}B NMR spectrum, suggesting the possibility of a Zn/B exchange process. Seeking a clean synthesis of this new product, we engineered a scheme to facilitate such an exchange. The species $[\text{tBu}_3\text{PN}_2\text{OB}(\text{C}_6\text{H}_4\text{F})_3]$ (**1**), containing a relatively weakly Lewis acidic borane, was prepared in a fashion similar to that described for $[\text{tBu}_3\text{PN}_2\text{OB}(\text{C}_6\text{F}_5)_2(\text{Ar})]$ ($\text{Ar} = \text{C}_6\text{F}_5$, Ph).^[14] NMR spectral parameters for **1** were similar to those reported for the perfluoroarylborane derivatives. However, in contrast to the known compounds, **1** undergoes a clean and facile reaction with an equivalent of $\text{tol}\cdot\text{Zn}(\text{C}_6\text{F}_5)_2$ resulting in the precipitation of a white solid **2**, which was isolated in essentially quantitative yield. NMR spectroscopic analysis in CD_2Cl_2 showed a new single ^{31}P resonance at 66.5 ppm. The fully ^{15}N labeled isotologue **2- ^{15}N** was synthesized from $[\text{tBu}_3\text{P}^{15}\text{N}_2\text{OB}(\text{C}_6\text{H}_4\text{F})_3]$ (**1- ^{15}N**). ^{15}N NMR signals at $\delta = 318.0$ and 599.1 ppm which exhibit N–P coupling of 9.3 and 54 Hz, respectively, and a coupling constant of $^1J_{\text{NN}} = 18$ Hz establish that the PN_2O fragment remains intact upon formation of **2**. ^{11}B and ^{19}F NMR spectra of the reaction mixture support the quantitative liberation of $\text{B}(\text{C}_6\text{H}_4\text{F})_3$. In addition, the ^{19}F NMR spectrum shows resonances at $\delta = -117.4$, -157.7 , and -162.6 ppm attributable to a $\{\text{Zn}(\text{C}_6\text{F}_5)_2\}$ -containing product. These data suggest the empirical formula of **2** is $[\text{tBu}_3\text{PN}_2\text{OZn}(\text{C}_6\text{F}_5)_2]$. A crystal structure determination established the centrosymmetric and dimeric nature of **2** (Figure 1)^[16] in which two $\text{tBu}_3\text{PN}_2\text{O}$ fragments bridge two Zn centers forming a $\{\text{Zn}_2\text{O}_2\}$ core. The Zn–O distances were found to be 2.088(2) and 2.144(2) Å, while the corresponding Zn–O–Zn' and O–Zn–O' angles are 107.15(10) and 72.85(8)°, respectively. The N–N and N–O distances in **2** are 1.266(4) and 1.308(3) Å, and are significantly elongated in comparison to free N_2O (1.127 and 1.186 Å).^[17]

The dimeric nature of the complex positions Zn(1) proximal to N(1) at a non-bonded distance of 3.035(2) Å. The substituents around the N=N double bond are disposed in

[*] R. C. Neu, Dr. E. Otten, Prof. Dr. D. W. Stephan
Department of Chemistry, University of Toronto
80 St. George St., Toronto, Ontario, M5S3H6 (Canada)
E-mail: dsteph@chem.utoronto.ca
Homepage: <http://www.chem.utoronto.ca/staff/dstephan>

[**] D.W.S. gratefully acknowledges the financial support of NSERC of Canada and the award of a Canada Research Chair and a Killam Research Fellowship. E.O. is grateful for the support of a Rubicon postdoctoral fellowship from the Netherlands Organisation for Scientific Research (NWO). R.C.N. is grateful for the award of an NSERC of Canada scholarship.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.200905650>.





Figure 1. POV-ray depiction of the molecular structure of 2.

a *trans* position, as is observed in the main group species $[t\text{Bu}_3\text{PN}_2\text{OBAr}_3]$.^[14]

Reaction of 1 with 1.5 equivalents of $[\text{tol}\cdot\text{Zn}(\text{C}_6\text{F}_5)_2]$ resulted in the clean formation of a new species 3, which was isolated in 81 % yield after crystallization (Scheme 1). A



Scheme 1. Synthesis of 2–4 starting from 1 (a, b, c=2, 3, 4 equivalents $[\text{tol}\cdot\text{Zn}(\text{C}_6\text{F}_5)_2]$, respectively, per 2 equivalents of 1) and conversion of 2→3→4.

crystallographic study established the structure of 3 as the C_2 symmetric compound $[(t\text{Bu}_3\text{N}_2\text{OZn}(\text{C}_6\text{F}_5)_2)_2\text{Zn}(\text{C}_6\text{F}_5)_2]$ (Figure 2)^[16] in which a single pseudo-tetrahedral Zn center bridges two $\{t\text{Bu}_3\text{PN}_2\text{OZn}(\text{C}_6\text{F}_5)_2\}$ units with $\text{Zn}(1)\text{--O}(1)$ distances of 2.118(2) Å. The $\text{Zn}(2)$ atoms in the latter units are coordinated to $\text{O}(1)$ and $\text{N}(1)$ of the N_2O fragment at distances of 2.184(2) and 2.242(2) Å, respectively. This yields a chelating four-membered $\{\text{ZnN}_2\text{O}\}$ ring and results in a $\text{N}(1)\text{--Zn}(2)\text{--O}(1)$ angle of 56.91(9)°. The ^{31}P NMR resonance of 3 is shifted slightly downfield ($\delta = 68.5$ ppm) compared to 2. The room temperature ^{19}F NMR spectrum shows only one




Figure 2. POV-ray depiction of the molecular structure of 3.

set of resonances for the C_6F_5 rings, suggesting that exchange between the two different $\{\text{Zn}(\text{C}_6\text{F}_5)_2\}$ environments is facile. Measuring the spectrum at -75°C reveals two distinct $\{\text{Zn}(\text{C}_6\text{F}_5)_2\}$ fragments in a 2:1 ratio, which is consistent with the solid state structure of 3. ^{15}N NMR signals for the isotopologue 3- ^{15}N are observed at $\delta = 323.8$ and 595.2 ppm with N–P and N–N couplings of 9.4, 54, and 18 Hz, respectively.

In an analogous reaction, 1 was treated with two equivalents of $[\text{tol}\cdot\text{Zn}(\text{C}_6\text{F}_5)_2]$ affording a new species 4 in 80 % isolated yield. Compound 4 gave rise to a ^{31}P resonance at $\delta = 71.7$ ppm, and ^{15}N NMR signals for the isotopologue 4- ^{15}N are observed at $\delta = 349.3$ and 582.5 ppm with N–P and N–N couplings of 11, 54, and 17 Hz, respectively.

The precise structural details of 4 were confirmed crystallographically (Figure 3), unambiguously establishing the formula as $[t\text{Bu}_3\text{PN}_2\text{O}(\text{Zn}(\text{C}_6\text{F}_5)_2)_2]$.^[16] This molecule contains two Zn atoms, one of which has a rare^[18] three-coordinate geometry being bound to the O atom of the N_2O fragment and two perfluoroaryl rings. The $\text{Zn}(1)\text{--O}(1)$ distance in this case is 2.0912(9) Å while the $\text{C}\text{--Zn}(1)\text{--C}$ angle is 153.23(6)°. A second Zn atom, $\text{Zn}(2)$, has a pseudo-tetrahedral coordination sphere comprised of two perfluoroaryl





Figure 3. POV-ray depiction of the molecular structure of 4.

rings, an O, and the P-bound N of N₂O, creating a ZnN₂O four-membered chelate ring similar to that seen in **3**. The resulting Zn(2)–O(1) and Zn(2)–N(1) distances in this case are 2.1435(10) and 2.3086(12) Å, respectively, while the chelate bite-angle at Zn(2) is 56.38(4)°.

As in **3**, the room temperature ¹⁹F NMR spectrum of **4** shows rapid exchange between the two {Zn(C₆F₅)₂} moieties. Decoalescence of the *o*-F resonances is observed at –34.6°C, corresponding to Δ*G*[‡] = 10.9 kcal mol^{–1} for the process exchanging the {Zn(C₆F₅)₂} environments. This low barrier suggests a mechanism involving the dissociation of the weak Zn–N interaction, followed by rotation about the N–O bond (Scheme 2).



Scheme 2. Proposed mechanism of {Zn(C₆F₅)₂} exchange in **4**.

A comparison of the metrical parameters of **2–4** (Table 1) shows that there is little variation in the bond lengths of the PN₂O fragment. A marginal elongation of the N=N bond is observed upon coordination of a {Zn(C₆F₅)₂} group to the

Table 1: Comparison of pertinent metrical parameters in **2–4**.^[a]

	2	3	4
P(1)–N(1)	1.703(2)	1.702(3)	1.7103(11)
N(1)–N(2)	1.266(4)	1.287(4)	1.2793(15)
N(2)–O(1)	1.308(3)	1.301(3)	1.3057(15)
Zn(1)–O(1)	2.088(2)	2.118(2)	2.0912(9)
Zn(2)–N(1)		2.242(2)	2.3086(12)
Zn(2)–O(1)		2.184(2)	2.1435(10)
N(1)–N(2)–O(1)	111.7(2)	109.2(2)	109.29(10)
Zn(1)–O(1)–Zn(2) ^[b]	107.15(10)	135.57(9)	139.95(5)
N(1)–Zn(2)–O(1)		56.91(9)	56.38(4)

[a] Distances in Å, angles in °. [b] Zn(1)' in case of **2**.

N₂O moiety (cf. **2** vs. **3** or **4**). At the same time, the N–N–O bond angle becomes slightly more acute in order to accommodate binding of Zn(2). It thus appears that coordination of a {Zn(C₆F₅)₂} group does not lead to a substantial perturbation of the PN₂O fragment. The terminal, three-coordinate Zn(1) center in **4** is more tightly bound to O(1) than the bridging Zn(1) in **3**, as expected based on its coordinative unsaturation. In addition, the greater steric congestion around the central Zn(1) in **3** forces the {Zn(C₆F₅)₂} fragment to be almost perpendicular to the PN₂O plane (C–Zn(1)–C/N–N–O interplanar angle **3**: 67.1(3)°; **4**: 23.84(16)°). This results in a close approach of two C₆F₅ rings in **3**, with concomitant displacement of Zn(2) away from O(1) towards N(1).

The formation of **2–4** from **1**^[19] is presumably driven by several factors, including the greater Lewis acidity of Zn–

(C₆F₅)₂ compared to B(C₆H₄F)₃ and the basicity of the N and O atoms within the PN₂O fragment that facilitates binding to additional Lewis acidic centers. In addition, the diminished steric congestion about {Zn(C₆F₅)₂} in comparison to triarylboranes allows the interaction of the PN₂O fragment with multiple Zn centers.

The chemistry described herein demonstrates that frustrated Lewis pairs can be employed to generate unusual species such as phosphine-stabilized N₂O fragments that can undergo exchange with other Lewis acids offering a unique route to Zn complexes containing the PNNO moiety. Moreover, the characterization of **2**, **3**, and **4** illustrates multiple binding modes for the interaction of an N₂O fragment with a metal. We continue to actively examine the further chemistry of frustrated Lewis pairs and in particular the potential for small-molecule complexation and activation.

Experimental Section

Synthesis of 2: A 20 mL scintillation vial was charged with **1** (0.100 g, 0.184 mmol) and [tol-Zn(C₆F₅)₂] (0.091 g, 0.185 mmol) in CH₂Cl₂ (5 mL). The solution was initially opaque but cleared after a few seconds of stirring. The reaction was left stirring for 1 h at room temperature. At this time, the solution was cloudy. Hexanes (10 mL) was added precipitating a fine white solid. The solid was isolated by filtration, washed with hexanes (3 × 5 mL), and dried in vacuo for 2 h. Crystals suitable for X-ray diffraction were grown from a layered CH₂Cl₂/pentane solution at –35°C. Yield: 0.118 g (99%). ¹⁹F NMR (376 MHz, CD₂Cl₂, 25°C): δ = –117.44 (m, *o*-C₆F₅), –157.71 (t, ³*J*_{F–F} = 19 Hz, *p*-C₆F₅), –162.64 ppm (m, *m*-C₆F₅); ³¹P{¹H} NMR (162 MHz, CD₂Cl₂, 25°C): δ = 66.50 ppm (s); ¹⁵N NMR (40.6 MHz, CD₂Cl₂, 25°C): δ = 599.07 (dd, ²*J*_{N–P} = 9.3, ¹*J*_{N–N} = 18 Hz, PNNO), 317.97 ppm (dd, ¹*J*_{N–P} = 54, ¹*J*_{N–N} = 18 Hz, PNNO).

Synthesis of 3: A 20 mL scintillation vial was charged with **1** (0.060 g, 0.111 mmol) and [tol-Zn(C₆F₅)₂] (0.082 g, 0.167 mmol) in CH₂Cl₂ (10 mL). The clear solution was left stirring for 1 h at room temperature. At this time, pentane (10 mL) was added precipitating a fine white solid. The product was allowed to settle and the solvent was decanted followed by washing of the solid with pentane (3 × 5 mL). The product was dried in vacuo for 1 h. Yield: 0.076 g (81%). Crystals suitable for X-ray diffraction were grown from a layered CH₂Cl₂/pentane solution at –35°C. ¹⁹F NMR (376 MHz, CD₂Cl₂, 25°C): δ = –117.56 (m, *o*-C₆F₅), –156.73 (t, ³*J*_{F–F} = 19 Hz, *p*-C₆F₅), –162.42 ppm (m, *m*-C₆F₅); ³¹P{¹H} NMR (162 MHz, CD₂Cl₂, 25°C): δ = 68.54 ppm (s); ¹⁵N NMR (40.6 MHz, CD₂Cl₂, 25°C): δ = 595.17 (dd, ²*J*_{N–P} = 9.4, ¹*J*_{N–N} = 18 Hz, PNNO), 323.78 ppm (dd, ¹*J*_{N–P} = 54, ¹*J*_{N–N} = 18 Hz, PNNO).

Synthesis of 4: A 20 mL scintillation vial was charged with **1** (0.064 g, 0.118 mmol) and [tol-Zn(C₆F₅)₂] (0.116 g, 0.236 mmol) in CH₂Cl₂ (10 mL). The clear solution was left stirring for 1 h at room temperature. At this time, pentane (10 mL) was added precipitating a fine white solid. The product was allowed to settle and the solvent was decanted followed by washing with pentane (3 × 5 mL). The solid was dried in vacuo for 1 h. Yield: 0.099 g, (80%). Crystals suitable for X-ray diffraction were grown from a layered CH₂Cl₂/cyclohexane solution at 25°C. ¹⁹F NMR (376 MHz, CD₂Cl₂, 25°C): δ = –117.62 (m, *o*-C₆F₅), –156.26 (t, ³*J*_{F–F} = 20 Hz, *p*-C₆F₅), –162.18 ppm (m, *m*-C₆F₅); ³¹P{¹H} NMR (162 MHz, CD₂Cl₂, 25°C): δ = 71.65 (s); ¹⁵N NMR (40.6 MHz, CD₂Cl₂, 25°C): δ = 582.52 (dd, ²*J*_{N–P} = 11, ¹*J*_{N–N} = 17 Hz, PNNO), 349.33 ppm (dd, ¹*J*_{N–P} = 54, ¹*J*_{N–N} = 17 Hz, PNNO).

Received: October 8, 2009

Published online: November 17, 2009

Keywords: frustrated Lewis pairs · nitrous oxide · zinc

- [1] J. N. Armor, H. Taube, *J. Am. Chem. Soc.* **1969**, *91*, 6874.
- [2] a) F. Bottomley, W. V. F. Brooks, *Inorg. Chem.* **1977**, *16*, 501; b) F. Bottomley, J. R. Crawford, *J. Am. Chem. Soc.* **1972**, *94*, 9092; c) F. Paulat, T. Kuschel, C. Näther, V. K. K. Praneeth, O. Sander, N. Lehnert, *Inorg. Chem.* **2004**, *43*, 6979; d) D. F. T. Tuan, R. Hoffmann, *Inorg. Chem.* **1985**, *24*, 871.
- [3] a) F. Bottomley, *Polyhedron* **1992**, *11*, 1707; b) J. T. Groves, J. S. Roman, *J. Am. Chem. Soc.* **1995**, *117*, 5594; c) W. H. Harman, C. J. Chang, *J. Am. Chem. Soc.* **2007**, *129*, 15128; d) W. A. Howard, G. Parkin, *J. Am. Chem. Soc.* **1994**, *116*, 606; e) H. Z. Yu, G. C. Jia, Z. Lin, *Organometallics* **2009**, *28*, 1158.
- [4] a) A. W. Kaplan, R. G. Bergman, *Organometallics* **1997**, *16*, 1106; b) G. A. Vaughan, G. L. Hillhouse, A. L. Rheingold, *J. Am. Chem. Soc.* **1990**, *112*, 7994; c) G. A. Vaughan, P. B. Rupert, G. L. Hillhouse, *J. Am. Chem. Soc.* **1987**, *109*, 5538; d) G. A. Vaughan, C. D. Sofield, G. L. Hillhouse, A. L. Rheingold, *J. Am. Chem. Soc.* **1989**, *111*, 5491; e) H. Z. Yu, G. C. Jia, Z. Y. Lin, *Organometallics* **2007**, *26*, 6769; f) H. Z. Yu, G. C. Jia, Z. Y. Lin, *Organometallics* **2008**, *27*, 3825.
- [5] N. D. Harrold, R. Waterman, G. L. Hillhouse, T. R. Cundari, *J. Am. Chem. Soc.* **2009**, *131*, 12872.
- [6] a) J.-P. F. Cherry, A. R. Johnson, L. M. Baraldo, Y.-C. Tsai, C. C. Cummins, S. V. Kryatov, E. V. Rybak-Akimova, K. B. Capps, C. D. Hoff, C. M. Haar, S. P. Nolan, *J. Am. Chem. Soc.* **2001**, *123*, 7271; b) A. R. Johnson, W. M. Davis, C. C. Cummins, S. Serron, S. P. Nolan, D. G. Musaev, K. Morokuma, *J. Am. Chem. Soc.* **1998**, *120*, 2071; c) C. E. Laplaza, A. L. Odom, W. M. Davis, C. C. Cummins, J. D. Protasiewicz, *J. Am. Chem. Soc.* **1995**, *117*, 4999.
- [7] J.-H. Lee, M. Pink, J. Tomaszewski, H. Fan, K. G. Caulton, *J. Am. Chem. Soc.* **2007**, *129*, 8706.
- [8] a) S. Poh, R. Hernandez, M. Inagaki, P. G. Jessop, *Org. Lett.* **1999**, *1*, 583; b) V. N. Parmon, G. I. Panov, A. Uriarte, A. S. Noskov, *Catal. Today* **2005**, *100*, 115.
- [9] a) D. Richardson, H. Felgate, N. Watmough, A. Thomson, E. Baggs, *Trends Biotechnol.* **2009**, *27*, 388; b) I. M. Wasser, S. de Vries, P. Moënne-Loccoz, I. Schröder, K. D. Karlin, *Chem. Rev.* **2002**, *102*, 1201.
- [10] a) P. Chen, S. I. Gorelsky, S. Ghosh, E. I. Solomon, *Angew. Chem.* **2004**, *116*, 4224; *Angew. Chem. Int. Ed.* **2004**, *43*, 4132; b) S. I. Gorelsky, S. Ghosh, E. I. Solomon, *J. Am. Chem. Soc.* **2006**, *128*, 278.
- [11] I. Bar-Nahum, A. K. Gupta, S. M. Huber, M. Z. Ertem, C. J. Cramer, W. B. Tolman, *J. Am. Chem. Soc.* **2009**, *131*, 2812.
- [12] a) G. Centi, A. Galli, B. Montanari, S. Perathoner, A. Vaccaria, *Catal. Today* **1997**, *35*, 113; b) P. J. Smeets, M. H. Groothaert, R. M. van Teeffelen, H. Leeman, E. J. M. Hensen, R. A. Schoonheydt, *J. Catal.* **2007**, *245*, 358; c) N. Russo, D. Fino, G. Saracco, V. Specchia, *Catal. Today* **2007**, *119*, 228; d) H. Jiang, F. Liang, S. Werth, T. Schiestel, J. Caro, *Angew. Chem.* **2009**, *121*, 3027; *Angew. Chem. Int. Ed.* **2009**, *48*, 2983.
- [13] a) D. J. Wuebbles, *Science* **2009**, *326*, 56; b) A. R. Ravishankara, J. S. Daniel, R. W. Portmann, *Science* **2009**, *326*, 123.
- [14] E. Otten, R. C. Neu, D. W. Stephan, *J. Am. Chem. Soc.* **2009**, *131*, 9918.
- [15] a) Transition metal complexes with related phosphazide (R_3PN_3R') ligands have been reported, see: b) W. Beck, W. Rieber, H. Kirmaier, *Z. Naturforsch. B* **1977**, *32*, 528; c) G. L. Hillhouse, G. V. Goeden, B. L. Haymore, *Inorg. Chem.* **1982**, *21*, 2064; d) A. A. Danopoulos, R. S. Hay-Motherwell, G. Wilkinson, S. M. Cafferkey, T. K. N. Sweet, M. B. Hursthouse, *J. Chem. Soc. Dalton Trans.* **1997**, 3177; e) G. C. Fortman, B. Captain, C. D. Hoff, *Organometallics* **2009**, *28*, 3587.
- [16] See Supporting Information.
- [17] J. J. L. Griggs, K. Narahari Rao, L. H. Jones, R. M. Potter, *J. Mol. Spectrosc.* **1968**, *25*, 34.
- [18] a) H. J. Gais, G. Bulow, G. Raabe, *J. Am. Chem. Soc.* **1993**, *115*, 7215; b) E. Martin, D. L. Hughes, S. J. Lancaster, *Eur. J. Inorg. Chem.* **2006**, 4037.
- [19] Compounds **3** and **4** are also cleanly obtained by consecutive addition of $[tol\cdot Zn(C_6F_5)]$ to **2** (Scheme 1).