


Linear Trimer of Diruthenium Linked by Butadiyn-Diyl Units: A Unique Electronic Wire**

Jie-Wen Ying, Isiah Po-Chun Liu, Bin Xi, You Song, Charles Campana, Jing-Lin Zuo, and Tong Ren*

The proposal of molecular wires based on metal oligoynyl compounds has been around for some time.^[1–3] Two types of structural motifs are featured prominently in these endeavors: dimers ($[M]-(C\equiv C)_n-[M]$) and oligomers ($\{[M]-(C\equiv C)_n\}_m-[M]$) bridged by a μ -C,C'-oligoyn-diyl unit. The earliest examples of dimers and oligomers were based on Pd^{2+} and Pt^{2+} species,^[4] which are insulators with band gaps generally larger than 3 eV.^[5] The significant electronic delocalization that is desired for molecular wires has been demonstrated in many dimeric compounds with metal centers, such as Fe, Re, and Ru.^[6] Recent efforts in measuring the current-voltage characteristics of metal alkynyl compounds in nanojunctions yielded encouraging results as well.^[7] Nevertheless, realizing similar delocalization in the $\{[M]-(C\equiv C)_n\}_m-[M]$ type oligomers remains synthetically challenging.^[8] The electronic delocalization in the $[M]-(C\equiv C)_n-[M]$ type dimers is mediated by the polyynediyl chains,^[6] while the electronic delocalization across the $[M]$ units has rarely been addressed. Hence, synthetic access to oligomers will provide the opportunity to assess electronic delocalization across both the carbon-rich backbone and metal centers. Studies of *trans*- $[Fc(C\equiv C)_n-Ru_2(DMBA)_4-(C\equiv C)_n-Fc]$ (DMBA is *N,N*-dimethylbenzamidinate, Fc is ferrocenyl) type compounds revealed that the central $Ru_2(DMBA)_4$ unit mediates strong electronic couplings between two Fc termini over distances up to 2.7 nm.^[9] Encouraged by these results, we began to explore the utility of $Ru_2(DMBA)_4$ species in forming oligomeric compounds. Reported herein are the synthesis and characterization of linear trimers **1a/b** (Scheme 1) that exhibit a multitude of unusual physical properties as a result of the strong inter-metallic delocalization, and a theoretical rationale on the basis of density functional theory (DFT) calculations.



Scheme 1. Preparation of compounds **1**.

The preparation of trimeric compounds **1a/b** was accomplished from the reaction between $[Ru_2(DiMeOap)_4(C_4H)]$ (**2**)^[10] and $[Ru_2(X-DMBA)_4(NO_3)_2]$ ^[11] in a 2:1 ratio in the presence of Et_2NH .^[12] Compounds **1a** and **1b** were purified by recrystallization from THF/hexanes in yields of 77 % and 58 %, respectively. Although the high-spin nature of **1a/b** prevents clean characterization by NMR spectroscopy, they were analyzed satisfactorily with mass spectrometry and elemental analysis. The trimeric nature of **1** was further confirmed through an X-ray diffraction study of **1a**^[13] (Figure 1). The structural analysis of **1a** reveals that the coordination spheres of both the $\{Ru_2(DiMeOap)_4\}$ termini and central $\{Ru_2(m-MeO-DMBA)_4\}$ unit bear close resemblance to those of $[Ru_2(DiMeOap)_4(C_4SiMe_3)]$ ^[10] and $[Ru_2(DMBA)_4(C_{2n}R')_2]$,^[11] respectively. The C–C bond lengths in




Figure 1. ORTEP plot of **1a** (thermal ellipsoids set 30 % probability). The asymmetric unit of **1a** consists of one half of the trimer, which is related to the other half through an inversion center. Selected bond lengths [Å]: Ru1–Ru2 2.326(1), Ru3–Ru3' 2.440(2), Ru2–C1 2.09(1), Ru3–C4 1.99(1), C1–C2 1.19(2), C2–C3 1.39(2), C3–C4 1.20(2).

[*] Dr. J.-W. Ying, Dr. I. P.-C. Liu, Dr. B. Xi, Prof. Dr. T. Ren

Department of Chemistry, Purdue University
West Lafayette, IN 47907 (USA)

Fax: (+1) 765-494-0239

E-mail: tren@purdue.edu

Prof. Dr. Y. Song, Prof. Dr. J.-L. Zuo

School of Chemistry and Chemical Engineering, Nanjing University
Nanjing 210093 (P. R. China)

Dr. C. Campana

Bruker AXS Inc

Madison, WI 53711 (USA)

[**] This work was supported in part by the US National Science Foundation (CHE 0715404), and the National Natural Science Foundation of China (20725104, 20771057, 20531040).

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/ange.200904674>.

the butadiyn-diyl fragment follow the expected pattern of alternating single and triple bonds. The distance between two $\{\text{Ru}_2(\text{DiMeOap})_4\}$ termini ($\text{Ru}_2\cdots\text{Ru}_2'$) is 18.04 Å.

The cyclic and differential voltammograms (CVs and DPVs) of compounds **1a/b** were examined to gauge the electronic delocalization therein, and the DPV of **1a** shown in Figure 2 reveals a very complex pattern in the window from




Figure 2. DPVs of compounds **1a**, and its precursors **2'** and **3** recorded in THF. The numbers denote the Ru oxidation states; “*” denotes peaks attributed to the degraded species. A plausible assignment is provided in the Supporting Information.

1.0 to -2.0 V. To facilitate the analysis of the DPV of **1a**, the DPVs of $[\text{Ru}_2(\text{DiMeOap})_4(\text{C}_4\text{SiMe}_3)]$ (**2'**)^[10] and $[\text{Ru}_2(m\text{-MeO-DMBA})_4(\text{C}_4\text{SiMe}_3)_2]$ (**3**)^[11] both precursors to **1a**, were also included in Figure 2. Compound **2'** exhibits a one-electron oxidation at 0.49 V and a one-electron reduction couple at -0.72 V, and compound **3** exhibits a one-electron oxidation at 0.74 V and a one-electron reduction at -0.88 V. Since the oxidation potential of **3** is far more positive than that of **2'**, the first two one-electron oxidations ($+1/0$ and $+2/+1$) in **1a** are attributed to the $\{\text{Ru}_2(\text{DiMeOap})_4\}$ termini, and the third oxidation to the central $[\text{Ru}_2(m\text{-MeO-DMBA})_4]$ unit. Similarly, because of the more negative reduction potential of **3**, the first two one-electron reductions ($0/-1$ and $-1/-2$) in **1a** are attributed to the Ru_2 termini, and the third reduction to the central Ru_2 unit. Both the pair-wise oxidations and reductions of the $\{\text{Ru}_2(\text{DiMeOap})_4\}$ termini in **1a** are the hallmark of electronic couplings in the mixed-valent states ($+1$ and -1).^[1,14] The potential differences within the pair-wise couples are $\Delta E(+1) = E(+2/+1) - E(+1/0) = 91$ mV and $\Delta E(-1) = E(0/-1) - E(-1/-2) = 95$ mV. These values, though small in comparison to strongly coupled compounds such as the Creutz-Taube ion,^[14] are still significant considering the large donor-acceptor separation (18 Å). Determination of the intervalence charge-transfer transitions in **1a** ^{± 1} would facilitate the quantification of electronic coupling, but the spectroelectrochemical study of **1a** was unsuccessful because of its instability over the timescale required for electrolysis.

Additional evidence of significant delocalization comes from the comparison of the electronic absorption spectra of compounds **1a**, **2'**, and **3**, which are shown in Figure 3. The spectrum of **1a** features intense peaks at 931, 770, 618, and




Figure 3. Vis-NIR spectra of compounds **1a**, **2'**, and **3** recorded in THF, along with the composite spectrum $2\epsilon(2') + \epsilon(3)$.

479 nm, and is very different from those of **2'** and **3**. A composite spectrum was calculated as the sum of spectrum **3** and the double of spectrum **2'**, and is included in Figure 3 as well. Should the electronic coupling among three Ru_2 units be negligible, the composite spectrum would be in good agreement with that of **1a**. While the peaks at 770 and 480 nm in compound **1a** were “reproduced” in the composite, the peaks at 618 nm and 931 nm observed for **1a** were absent in the composite. The appearance of these “new” bands must be the consequence of extensive conjugation among three Ru_2 units through the butadiyn-diyl bridges.

The magnetic susceptibility of compounds **1a** and **2'** were measured in the temperature range of 1.8–300 K (Figure 4).^[15] The $\chi_m T$ characteristics for **2'** are consistent with a $S = 3/2$ ground state that undergoes zero-field splitting. The fitting of $\chi_m T$ data resulted in $g = 2.09$ and $|D| = 56.33$ cm^{-1} , which are




Figure 4. Plots of $\chi_m T$ versus T of compounds **1a** (●) and **2'** (○). The solid line represents the best theoretical fit of the data for **2'**.

in line with studies of related $\text{Ru}_2^{\text{II,III}}$ species.^[16] The $[\text{Ru}_2(\text{DMBA})_4(\text{C}\equiv\text{CR})_2]$ type compounds are generally diamagnetic ($S=0$).^[3] The magnetic properties of **1a** would be dominated by two $\{\text{Ru}_2(\text{DiMeOap})_4\}$ termini ($S=3/2$) with a room temperature $\chi_m T$ value around $3.75 \text{ cm}^3 \text{ K mol}^{-1}$ if the $-(\text{C}\equiv)_2\text{-Ru}_2(\text{DMBA})_4-(\text{C}\equiv\text{C})_2-$ fragment remains $S=0$ and functions as a weak mediator of spin coupling.^[17] However, compound **1a** exhibits a significantly higher $\chi_m T$ value of $5.67 \text{ cm}^3 \text{ K mol}^{-1}$ at 300 K, implying the presence of additional unpaired electrons in **1a**. Given the precedent of the $S=1$ state for Ru_2^{6+} compounds with different axial ligands^[18] and other examples of spin-admixed diruthenium species,^[19] it is possible that the unusually large $\chi_m T$ value of **1a** arises from the contribution of an $S=1$ state of the central Ru_2^{6+} moiety. Hence, the spin distribution in **1a** may be assigned as $(3/2)-(1)-(3/2)$. The theoretical $\chi_m T$ value of this model is 5.28 ($g=2.09$ for $\text{Ru}_2^{5+}(S=3/2)$ and g is 2.18 for $\text{Ru}_2^{6+}(S=1)$), and is in agreement with the experimental value at 300 K.^[18] On the other hand, the $\chi_m T$ value of **1a** at 1.8 K is $2.17 \text{ cm}^3 \text{ K mol}^{-1}$, which is twice of that of **2'**. This result suggests that the magnetic centers of **1a** at low temperature consist of two $\text{Ru}_2^{5+}(S=3/2)$ centers only. The spin state of central Ru_2^{6+} moiety changes from $S=1$ to $S=0$ with decreasing temperature (spin transition). The temperature dependence of the $\chi_m T$ values of **1a** is a combined effect of the zero-field splitting, antiferromagnetic interaction between the Ru_2^{5+} and Ru_2^{6+} centers, and the spin transition at the Ru_2^{6+} center, which is too complicated to be modeled.

To rationalize the unusual magnetism of **1a**, DFT/B3LYP calculations were performed.^[20] The computed ground state of **1a** is a singlet state that results from weak antiferromagnetic interaction between two $S=3/2$ Ru_2^{5+} termini (Figure 5). The computed coupling constant (J) between the




Figure 5. Relative energies of four energy states with possible spin alignment of **1a**. S denotes the calculated spin angular momentum.

two Ru_2^{5+} moieties (-0.18 cm^{-1}) is so small that the ferromagnetic $(3/2)-(3/2)$ and antiferromagnetic $(3/2)-(-3/2)$ states are nearly degenerate. The two Ru_2^{5+} termini are thus independent, which is consistent with the observed $\chi_m T$ at 1.8 K. Two excited states, $(3/2)-(1)-(3/2)$ and $(3/2)-(-1)-(3/2)$ derived from the spin flip at the Ru_2^{6+} center, lie at approximately 0.50 and 0.44 eV above the ground state. The computed coupling constant between the Ru_2^{5+} and Ru_2^{6+} centers is -72.26 cm^{-1} , implying moderate antiferromagnetic interaction.

To further understand the spin flip in **1a**, a companion calculation was performed for the model compound **4**, $[\text{Ru}_2(\text{DMBA})_4(\text{C}_4\text{H}_2)]$, which retains the central Ru_2 unit of **1a** with two Ru_2^{5+} termini being replaced by hydrogen atoms.^[20] The two unpaired electrons in **4** ($S=1$) occupy the π^* and δ^* orbitals with a 1.01 eV energy difference. Correspondingly, the SOMO and SOMO-3 in **1a** exhibit the π^* and δ^* characters of the central Ru_2^{6+} moiety, respectively. Comparing the π^* orbital of **4** with that of **1a**, shows the latter exhibits a pronounced contribution from two Ru_2^{5+} -butadiynyl fragments (Figure 6). Clearly, the presence of




Figure 6. π^* orbitals of a) **4** (SOMO) and b) **1a** (SOMO-3).

Ru_2^{5+} termini results in an enhanced π - π antibonding interaction between the butadiynyl ligand and the Ru_2^{6+} core, which destabilizes the $\pi^*(\text{Ru}_2^{6+})$ in **1a** and reduces the $\pi^*(\text{SOMO-3})-\delta^*(\text{SOMO})$ gap from 1.01 (in **4**) to 0.72 eV. The actual $\pi^*-\delta^*$ gap could be much smaller than the computed one, which would facilitate the spin transition at elevated temperature. Moreover, the spin population of the central Ru atoms in **1a** is 0.77, which is slightly higher than that in **4** (0.72). The increase in the spin population of the central Ru atoms may qualitatively indicate that the unpaired electrons in **1a** delocalize from the terminal Ru_2^{5+} moieties to the central Ru_2^{6+} unit.^[21]

In summary, we report very unusual trimeric compounds, **1**, based on Ru_2 species linked by the butadiyn-diyl bridges, for which both the voltammetric and spectroscopic data are consistent with an extensive delocalization over a span of 20 Å. A spin transition was also observed for **1a** and rationalized on the basis of DFT calculations by the existence of low-lying triplet states that are very close in energy to the singlet ground state.

Received: August 22, 2009

Revised: November 11, 2009

Keywords: delocalization · mixed-valent compounds · polyynes · ruthenium · spin transitions

- [1] F. Paul, C. Lapinte, *Coord. Chem. Rev.* **1998**, *178–180*, 431–509.
- [2] a) S. Szafert, J. A. Gladysz, *Chem. Rev.* **2003**, *103*, 4175–4206; b) H. Lang, *Angew. Chem.* **1994**, *106*, 569–572; *Angew. Chem. Int. Ed. Engl.* **1994**, *33*, 547–550.
- [3] T. Ren, *Organometallics* **2005**, *24*, 4854–4870.
- [4] N. Hagihara, K. Sonogashira, S. Takahashi, *Adv. Polym. Sci.* **1980**, *40*, 149–179.
- [5] A. E. Dray, F. Wittmann, R. H. Friend, A. M. Donald, M. S. Khan, J. Lewis, B. F. G. Johnson, *Synth. Met.* **1991**, *41*, 871–874.
- [6] a) N. Le Narvor, L. Toupet, C. Lapinte, *J. Am. Chem. Soc.* **1995**, *117*, 7129–7138; b) M. Brady, W. Weng, Y. Zou, J. W. Seyler, A. J. Amoroso, A. M. Arif, M. Bohme, G. Frenking, J. A. Gladysz, *J. Am. Chem. Soc.* **1997**, *119*, 775–788; c) M. I. Bruce, P. J. Low, K. Costuas, J.-F. Halet, S. P. Best, G. A. Heath, *J. Am. Chem. Soc.* **2000**, *122*, 1949–1962; d) G.-L. Xu, G. Zou, Y.-H. Ni, M. C. DeRosa, R. J. Crutchley, T. Ren, *J. Am. Chem. Soc.* **2003**, *125*, 10057; e) S. Rigaut, C. Olivier, K. Costuas, S. Choua, O. Fadhel, J. Massue, P. Turek, J.-Y. Saillard, P. H. Dixneuf, D. Touchard, *J. Am. Chem. Soc.* **2006**, *128*, 5859–5876.
- [7] a) T. L. Schull, J. G. Kushmerick, C. H. Patterson, C. George, M. H. Moore, S. K. Pollack, R. Shashidhar, *J. Am. Chem. Soc.* **2003**, *125*, 3202–3203; b) A. S. Blum, T. Ren, D. A. Parish, S. A. Trammell, M. H. Moore, J. G. Kushmerick, G.-L. Xu, J. R. Deschamps, S. K. Pollack, R. Shashidhar, *J. Am. Chem. Soc.* **2005**, *127*, 10010–10011; c) B. Kim, J. M. Beebe, C. Olivier, S. Rigaut, D. Touchard, J. G. Kushmerick, X.-Y. Zhu, C. D. Frisbie, *J. Phys. Chem. C* **2007**, *111*, 7521–7526; d) A. K. Mahapatro, J. Ying, T. Ren, D. B. Janes, *Nano Lett.* **2008**, *8*, 2131–2136.
- [8] K.-T. Wong, J.-M. Lehn, S.-M. Peng, G.-H. Lee, *Chem. Commun.* **2000**, 2259–2260.
- [9] a) G.-L. Xu, R. J. Crutchley, M. C. DeRosa, Q.-J. Pan, H.-X. Zhang, X. Wang, T. Ren, *J. Am. Chem. Soc.* **2005**, *127*, 13354–13363; b) B. Xi, G.-L. Xu, P. E. Fanwick, T. Ren, *Organometallics* **2009**, *28*, 2338–2341.
- [10] B. Xi, G.-L. Xu, J.-W. Ying, H.-L. Han, A. Cordova, T. Ren, *J. Organomet. Chem.* **2008**, *693*, 1656–1663.
- [11] G.-L. Xu, C. G. Jablonski, T. Ren, *J. Organomet. Chem.* **2003**, *683*, 388–397.
- [12] Compound **1a** was prepared from the reaction between $[\text{Ru}_2(m\text{-MeO-DMBA})_4(\text{NO}_3)_2]$ (0.84 g, 0.080 mmol) and **2** (0.188 g, 0.16 mmol) in THF/ Et_3NH (3:1, v:v) at room temperature for 1 h. The blue solid obtained upon solvent removal was recrystallized from THF/hexanes to afford 0.200 g (77 %) **1a** as a brown powder. Data: MS-MALDI (m/z , based on ^{101}Ru): 3259 [M^+]; elemental analysis (%) calcd for $\text{C}_{152}\text{H}_{156}\text{N}_{24}\text{O}_{20}\text{Ru}_6$: C 56.25, H 4.84, N 10.36; Found C 56.01, H 4.88, N 9.85. Electrochemistry, $E_{1/2}/\text{V}$, $\Delta E_p/\text{V}$, $i_{\text{backward}}/i_{\text{forward}}$: $\pm 3/\pm 2$, 0.686, 0.063, 1.029; $\pm 2/\pm 1$, 0.328, 0.057, 0.696; $\pm 1/0$, 0.237, 0.062, 1.167; $0/-1$, -0.881, 0.095, 0.939; $-1/-2$, -0.976, 0.117, 0.703; $-2/-3$, -1.292, 0.074, 0.137. Preparation of **1b** is similar to that of **1a**.
- [13] Crystallographic data for **1a**: space group $C2/c$, $a = 56.642(3)$ Å, $b = 16.6840(8)$ Å, $c = 16.0750(8)$ Å, $\beta = 95.001(3)^\circ$, $V = 15133(1)$ Å³, $Z = 4$, $\rho_{\text{calcd}} = 1.438$ Mg m⁻³, $\mu = 5.329$ mm⁻¹. X-ray intensity data of **1a** were measured at 100 K on a Bruker AXS SMART APEX II CCD-based diffractometer using $\text{Cu}_{K\alpha}$ ($\lambda = 1.54178$ Å), and the structure was solved using direct method; 34497 reflections collected, 12505 unique reflections ($R_{\text{int}} = 0.0936$), data/restraints/parameters 12505/0/933, final R indices ($I > 2\sigma(I)$) $R1 = 0.0977$ and $wR2 = 0.248$. CCDC 745048 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [14] W. Kaim, G. K. Lahiri, *Angew. Chem.* **2007**, *119*, 1808–1828; *Angew. Chem. Int. Ed.* **2007**, *46*, 1778–1796.
- [15] Magnetic susceptibility measurements for both compounds **1a** and **2'** were obtained with the use of a Quantum Design MPMS-XL7 SQUID magnetometer at temperature range of 1.8–300 K.
- [16] a) V. M. Miskowski, M. D. Hopkins, J. R. Winkler, H. B. Gray in *Inorganic Electronic Structure and Spectroscopy*, Vol. 2 (Eds.: E. I. Solomon, A. B. P. Lever), Wiley, New York, **1999**, pp. 343–402; b) F. A. Cotton, T. Ren, *Inorg. Chem.* **1995**, *34*, 3190–3193.
- [17] Y. Shi, G. T. Yee, G. Wang, T. Ren, *J. Am. Chem. Soc.* **2004**, *126*, 10552–10553.
- [18] F. A. Cotton, C. A. Murillo, J. H. Reibenspies, D. Villagran, X. P. Wang, C. C. Wilkinson, *Inorg. Chem.* **2004**, *43*, 8373–8378.
- [19] M. C. Barral, S. Herrero, R. Jiménez-Aparicio, M. R. Torres, F. A. Urbanos, *Angew. Chem.* **2005**, *117*, 309–311; *Angew. Chem. Int. Ed.* **2005**, *44*, 305–307.
- [20] Single point calculations on compounds **1a** and **4** were carried out using the DFT formalism with the spin unrestricted option as implemented in the Gaussian 03 program with the B3LYP functional. 6-31G basis sets were used to describe the H atoms. 6-31G* basis sets were used to describe the C, N, and O atoms. For more details, see the Supporting Information.
- [21] I. P. C. Liu, M. Bénard, H. Hasanov, I.-W. P. Chen, W.-H. Tseng, M.-D. Fu, M.-M. Rohmer, C.-h. Chen, G.-H. Lee, S.-M. Peng, *Chem. Eur. J.* **2007**, *13*, 8667–8677.