



Total Synthesis of (–)-Dactylolide**

Sang Young Yun, Eric C. Hansen, Ivan Volchkov, Eun Jin Cho, Wai Yip Lo, and Daesung Lee*

Since its discovery in 2001 by Riccio and co-workers,^[1] dactylolide **1** has attracted considerable synthetic interest.^[2] Several total syntheses of both natural (+)- and unnatural (–)-dactylolide have been reported. Extensive interest in dactylolide stems from not only its distinctive structural features, including a highly unsaturated 18-membered macrolactone with an embedded 2,6-disubstituted tetrahydropyran moiety, but also its biological activity toward several tumor cell lines such as L1210 and SK-OV-3 (63 % and 40 % inhibition, respectively, at 3.2 $\mu\text{g mL}^{-1}$).^[1] Furthermore, a closely related natural product zampanolide **2**,^[2b] has shown significantly more potent biological activity by blocking cell cycle in G2M transition through the stabilization of microtubules.^[3–5] From a standpoint of the biosynthetic origin, it is interesting to note that the absolute configuration of the identical macrolide substructures contained in these two natural compounds is opposite.^[6]



Recently, our research group has developed synthetic methods for efficient construction of trisubstituted Z-vinyl-boronates and 4-exomethylene-*cis*-2,6-disubstituted tetrahydropyrans. As an application to showcase these synthetic methods, herein we describe a concise route to (–)-dactylolide. Our retrosynthesis shown in Scheme 1 focuses on harnessing the power of transition-metal-catalyzed reactions to effect C–C and C–O bond formations. Often, the products from these reactions serve as the direct substrate for the next reaction,^[7] thus minimizing the need for functional group transformations and protection/deprotection steps. We envision the macrocycle of (–)-**1** will be achieved through a late-



Scheme 1. Retrosynthesis of (–)-dactylolide (*ent*-**1**). PG = protecting group.

stage ring-closing metathesis (RCM) reaction^[8] to form the C16–C17 double bond from **3** or relay metathesis substrate **4**.^[9] To access **3** and **4**, we planned to use a Suzuki–Miyaura^[10] coupling of iodoacrylate **5** and **6** and cyclic boronic acid half ester **7**. This is in turn accessed through the ruthenium-catalyzed Alder–ene reaction (RCAER)^[11] to form the C5–C6 bond with the terminal double bond of **8** and alkynyl boronate **9** and subsequent rhodium-catalyzed allylic [1,3]-transposition.^[12] The pyran subunit in **8** would be installed using a tandem RCAER and palladium-catalyzed ring closure.^[13]


The synthesis of tetrahydropyran **8** commenced with the RCAER between ethyl carbonate **10** and homopropargylic alcohol **11**^[14] and led to 1,4-diene **12** (Scheme 2). Treatment of **12** with palladium catalyst in the presence of Trost's chiral (+)-DPPBA^[15] ligand afforded 2,6-*cis*-disubstituted tetrahydropyran **13** (11:1 diastereomeric ratio). This transformation was best achieved using a one-pot procedure,^[16] and gave **13** in 72 % yield. Removal of the pivalate group and oxidation of the resulting alcohol gave aldehyde **14**. Leighton allylation^[17] and subsequent TBS protection afforded homoallyl silyl ether moiety of **8**. The RCAER of alkynyl boronate **9** occurred selectively with the least hindered double bond at C7 of **8**, and afforded vinyl boronate **15** (*cis/trans* = 5:1). The 1,3-transposition of the allyl alcohol of **15** with rhodium oxide in ether^[18] afforded a relatively unstable cyclic boronic acid half ester **7**.

Subsequently, Suzuki coupling partner 2-iodoacrylate **5** was prepared from TBS-protected (*S*)-glycidol **16** (Scheme 3). The opening of the epoxide ring with 2-propenylmagnesium chloride and a subsequent Mitsunobu reaction^[19] of **17** with 2-iodoacrylic acid **18**^[20] provided ester **5**. The Suzuki coupling between **5** and **7** delivered **3** (79 %). An attempted RCM with alcohol **3** did not yield the expected 18-membered lactone,


[*] Dr. S. Y. Yun, Dr. E. C. Hansen, I. Volchkov, Dr. E. J. Cho, W. Y. Lo, Prof. D. Lee
Department of Chemistry
University of Illinois at Chicago
845 West Taylor Street, Chicago, IL 60607 (USA)
Fax: (+1) 312-996-0431
E-mail: dsunglee@uic.edu

[**] We acknowledge the UIC for financial support of this research. We are grateful to Prof. Amos B. Smith III for providing us with NMR and IR spectra of **29** and dactylolide.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/ange.201001681>.




Scheme 2. a) 7 mol % $[\text{RuCp}(\text{CH}_3\text{CN})_3]\text{PF}_6$, acetone, RT; b) 3 mol % $[\text{Pd}_2(\text{dba})_3]\cdot\text{CHCl}_3$, 9 mol % (+)-DPPBA, CH_2Cl_2 , -20°C to RT, 70% over two steps; c) DIBAL, CH_2Cl_2 , -78°C , 96%; d) IBX, DMSO, 95%; e) (S,S)-Leighton reagent, CH_2Cl_2 , -10°C , 85% (8:1 d.r.); f) TBSO, imidazole, DMF, 98%; g) 10 mol % $[\text{RuCp}(\text{CH}_3\text{CN})_3]\text{PF}_6$, acetone, RT, 65% (78% brsm), h) 10 mol % Re_2O_7 , 65%. brsm = based on recovered starting material, Cp = cyclopentadienyl, dba = trans,trans-dibenzylideneacetone, DIBAL = diisobutylaluminum hydride, DMF = *N,N*-dimethylformamide, DMSO = dimethyl sulfoxide, DPPBA = 1,2-diaminocyclohexane-*N,N'*-bis(2'-diphenyl-phosphinobenzoyl), IBX = 2-iodoxybenzoic acid, Piv = pivoyl, TBS = tert-butyldimethylsilyl.




Scheme 3. a) 2-Propenylmagnesium chloride, -30°C , THF, 82%; b) DIAD, PPh_3 , THF, 0°C , 84%; c) **7**, $[\text{Pd}(\text{PPh}_3)_4]$, TIOEt , THF/ H_2O (3:1), 79%; d) DMP, CH_2Cl_2 , 89%. DIAD = diisopropyl azodicarboxylate, DMP = Dess–Martin periodinane, THF = tetrahydrofuran.

but instead generated a six-membered lactone through ring closure at C17–C2. Believing that the local conformation of the ketone at C7 should be similar to that of dactylolide, alcohol **3** was oxidized to α,β -unsaturated ketone **19** (89%).

With the pivotal RCM substrate **19** in hand, we explored various reaction conditions to maximize the formation of lactone **21** (Scheme 4). Treatment of **19** with the second-generation Grubbs catalyst **24** in the presence of benzoquinone^[21] delivered the desired macrolactone **21** in 45% yield. Presumably, the RCM reaction occurred via the alkylidene intermediate **20**. Other catalysts **25** and **26** as well as the first-generation Grubbs catalyst resulted in lower yield or no



27 (C19 epimer of **19**) $\xrightarrow{\text{a) catalyst, CH}_2\text{Cl}_2}$ **28** (C19 epimer of **21**)
 catalyst = **24** 35% (isolated)
25 32% (34% recovered **27**)
26 0% (<5% conversion)




Scheme 4. a) 10 mol % **24–26**, 10 mol % benzoquinone, CH_2Cl_2 , 65°C , 45%. Cy = cyclohexyl, Mes = mesityl.

reaction. We expected that the efficiency of the ring closure should be improved by generating an alternative alkylidene **23** because its ring closure with the less hindered double bond at C16 should be more facile. To test this hypothesis, we prepared an RCM substrate **22** that contains a relay metathesis tether. However, **22** did not cyclize directly into **21**, instead only to **19** by cleaving off the relay tether. This cleavage results from a faster methylene transfer from **22** to alkylidene **23** rather than its macrocyclic RCM. Based on this behavior and different modes of RCM with **3** and **19**, we surmised that the conformation of long linear substrates plays a more important role than the steric hindrance of the reacting double bond. To gain more insight into these divergent RCM behaviors, we examined RCM of diastereomeric substrate **27** (C19 epimer of **19**; Scheme 4). Interestingly, the RCM of **27** was similar to that of **21**, and gave macrolactone **28** (C19 epimer of **21**) with similar efficiency.^[22] This outcome suggests that the RCM behavior of these substrates cannot be predicted simply on the basis of steric hindrance or local conformation.

Completion of (–)-dactylolide **ent-1** was accomplished by removal of the TBS group from **21**, thus generating **29** with an alcohol group at the C20—an intermediate reported by Smith and Safonov^[2a] in their synthesis—and its subsequent oxidation with Dess–Martin periodinane (Scheme 5). The spectroscopic data of both **29** and **ent-1** are identical to those reported in the literature.

In conclusion, we have achieved a concise total synthesis of (–)-dactylolide. Several distinctive features in this syn-



Scheme 5. a) 1 N HCl, MeOH, 95 %; b) DMP, pyridine, CH₂Cl₂, 90 %.

thesis include: 1) a one-pot synthesis of *cis*-2,6-disubstituted-4-methylene tetrahydropyran through RCAER and palladium-catalyzed ring closure, 2) an Alder–ene reaction of alkynyl boronates and subsequent rhenium-catalyzed allylic transposition, and 3) a macrocyclic RCM to form an 18-membered lactone with concomitant installation of a trisubstituted double bond. The effectiveness and versatility of transition-metal-catalyzed transformations demonstrated in this synthetic maneuver can be further exploited in the synthesis of other complex molecular frameworks.

Received: March 20, 2010

Published online: May 10, 2010

Keywords: Alder–ene reactions · allylic transposition · natural products · ring-closing metathesis · total synthesis

- [1] A. Cutignano, I. Bruno, G. Bifulco, A. Casapullo, C. Debitus, L. Gomez-Paloma, R. Riccio, *Eur. J. Org. Chem.* **2001**, 775–778.
- [2] a) A. B. Smith III, I. G. Safonov, *Org. Lett.* **2002**, 4, 635–637; b) T. R. Hoye, M. Hu, *J. Am. Chem. Soc.* **2003**, 125, 9576–9577; c) F. Ding, M. P. Jennings, *Org. Lett.* **2005**, 7, 2321–2324; d) D. L. Aubele, S. Wan, P. E. Floreancig, *Angew. Chem.* **2005**, 117, 3551–3554; *Angew. Chem. Int. Ed.* **2005**, 44, 3485–3488; e) C. C. Sanchez, G. E. Keck, *Org. Lett.* **2005**, 7, 3053–3056; f) I. Louis, N. L. Hungerford, E. J. Humphries, M. D. McLeod, *Org. Lett.* **2006**, 8, 1117–1120; g) J. Uenishi; T. Iwamoto J. Tanaka, *Org. Lett.* **2009**, 11, 3262–3265.
- [3] T. Higa, J. Tanaka, *Tetrahedron Lett.* **1996**, 37, 5535–5538.
- [4] A. B. Smith III, I. G. Safonov, R. M. Corbett, *J. Am. Chem. Soc.* **2001**, 123, 12426–12427.
- [5] J. J. Field, A. J. Singh, A. Kanakkanthara, T. Halafihi, P. T. Northcote, J. H. Miller, *J. Med. Chem.* **2009**, 52, 7328–7332.
- [6] A. B. Smith, I. G. Safonov, R. M. Corbett, *J. Am. Chem. Soc.* **2002**, 124, 11102–11113.
- [7] a) T.-L. Ho, *Tandem Organic Reactions*, Wiley, New York, **1992**; b) L. F. Tietze, *Chem. Rev.* **1996**, 96, 115–136; c) J.-C. Wasilke, S. J. Obrey, R. T. Baker, G. C. Bazan, *Chem. Rev.* **2005**, 105, 1001–1020.
- [8] a) R. H. Grubbs, S. Chang, *Tetrahedron* **1998**, 54, 4413–4449; b) *Handbook of Metathesis*, Vol. 2 (Ed.: R. H. Grubbs), Wiley-VCH, Weinheim, **2003**; c) R. R. Schrock, A. H. Hoveyda, *Angew. Chem.* **2003**, 115, 4740–4782; *Angew. Chem. Int. Ed.* **2003**, 42, 4592–4633; d) S. J. Connon, S. Blechert, *Angew. Chem.* **2003**, 115, 1944–1968; *Angew. Chem. Int. Ed.* **2003**, 42, 1900–1923; e) A. Deiters, S. F. Martin, *Chem. Rev.* **2004**, 104, 2199–2238; f) K. C. Nicolaou, P. G. Bulger, D. Sarlah, *Angew. Chem.* **2005**, 117, 4564–4601; *Angew. Chem. Int. Ed.* **2005**, 44, 4490–4527.
- [9] a) T. R. Hoye, C. S. Jeffrey, M. A. Tennakoon, J. Wang, H. Zhao, *J. Am. Chem. Soc.* **2004**, 126, 10210–10211; b) E. C. Hansen, D. Lee, *Org. Lett.* **2004**, 6, 2035–2038; for a review, see: c) D. J. Wallace, *Angew. Chem.* **2005**, 117, 1946–1949; *Angew. Chem. Int. Ed.* **2005**, 44, 1912–1915.
- [10] a) J. Uenishi, J.-M. Beau, R. W. Armstrong, Y. Kishi, *J. Am. Chem. Soc.* **1987**, 109, 4756–4758; b) S. A. Frank, H. Chen, R. K. Kunz, M. J. Schnaderbeck, W. R. Roush, *Org. Lett.* **2000**, 2, 2691–2694; for a review, see: c) N. Miyaoura, A. Suzuki, *Chem. Rev.* **1995**, 95, 2457–2483.
- [11] a) E. C. Hansen, D. Lee, *J. Am. Chem. Soc.* **2005**, 127, 3252–3253; for a review, see: b) B. M. Trost, M. U. Frederiksen, M. T. Rudd, *Angew. Chem.* **2005**, 117, 6788–6825; *Angew. Chem. Int. Ed.* **2005**, 44, 6630–6666.
- [12] a) S. Bellemin-Lapponnaz, H. Gisie, J. P. Le Ny, J. A. Osborn, *Angew. Chem.* **1997**, 109, 1011–1013; *Angew. Chem. Int. Ed. Engl.* **1997**, 36, 976–978; b) S. Bellemin-Lapponnaz, H. Gisie, J. P. Le Ny, J. A. Osborn, *Tetrahedron Lett.* **2000**, 41, 1549–1552.
- [13] E. C. Hansen, D. Lee, *Tetrahedron Lett.* **2004**, 45, 7151–7155.
- [14] G. Pattenden, M. A. Gonzalez, P. B. Little, D. S. Millan, A. T. Plowright, J. A. Tornos, T. Ye, *Org. Biomol. Chem.* **2003**, 1, 4173–4208.
- [15] B. M. Trost, M. R. Machacek, A. Aponick, *Acc. Chem. Res.* **2006**, 39, 747–760.
- [16] B. M. Trost, M. R. Machacek, B. D. Faulk, *J. Am. Chem. Soc.* **2006**, 128, 6745–6754.
- [17] J. W. A. Kinnaird, P. Y. Ng, K. Kubota, X. Wang, J. L. Leighton, *J. Am. Chem. Soc.* **2002**, 124, 7920–7921.
- [18] E. C. Hansen, D. Lee, *J. Am. Chem. Soc.* **2006**, 128, 8142–8143.
- [19] O. Mitsunobu, *Synthesis* **1981**, 1–28.
- [20] R. Takeuchi, K. Tanabe, S. Tanaka, *J. Org. Chem.* **2000**, 65, 1558–1561.
- [21] S. H. Hong, D. P. Sanders, C. W. Lee, R. H. Grubbs, *J. Am. Chem. Soc.* **2005**, 127, 17160–17161.
- [22] With catalyst **24**, no starting material **27** was recovered. However, 34 and > 95 % of **27** was recovered intact with catalyst **25** and **26**, respectively.