

Synthesis of Polyarylated Methanes through Cross-Coupling of Tricarbonylchromium-Activated Benzylolithiums**

Genette I. McGrew, Jesada Temaismithi, Patrick J. Carroll, and Patrick J. Walsh*


Polyarylated methanes are attracting considerable attention due to their growing importance in developing medicinal agents for cancer^[1] and vascular disease,^[2,3] as well as in leuco dye precursors,^[4,5] photochromic agents, and applications in materials science.^[6,7] Most syntheses of polyarylmethanes involve Friedel–Crafts-type (F-C) electrophilic aromatic substitution reactions,^[5,6,8–13] although there are some limited exceptions.^[4,14,15] Recently, modified F-C processes,^[9,16] including a novel copper-catalyzed aza-Friedel–Crafts variant,^[17] have been used in the preparation of diversely substituted triarylmethanes. The F-C approach, however, is limited by both reactivity and selectivity: the nucleophile must be electron-rich and unhindered for adequate reactivity and the selectivity is controlled by the relative directing abilities of the substituents. Thus, triarylmethanes with certain electron withdrawing groups and those with *meta* substitution, are largely inaccessible. A complementary and general route to polyarylated methanes that enables the synthesis of currently inaccessible members of this important structural class is needed.

A strategy that would circumvent the limitations of the F-C reaction is based on benzylic anion synthons. Traditionally, reagents for this synthon are based on metals such as boron, tin or zinc to temper reactivity.^[18–20] An alternative approach to attenuate the reactivity of benzylic nucleophiles is

η^6 -coordination to a metal fragment such as $\{\text{Cr}(\text{CO})_3\}$.^[21–24] Along these lines, Kalinin and co-workers generated $[(\eta^6\text{-C}_6\text{H}_5\text{CH}_2\text{Li})\text{Cr}(\text{CO})_3]$, but were unable to effect palladium-catalyzed cross-coupling reactions. After transmetalation to zinc the resulting $[(\eta^6\text{-C}_6\text{H}_5\text{CH}_2\text{ZnCl})\text{Cr}(\text{CO})_3]$ underwent palladium catalyzed single coupling with aryl halides to furnish diarylmethanes in low yield (average 37%).^[25] The intermediate transmetalation to zinc prevents the realization of the tricarbonylchromium group's full potential: to activate more than one benzylic C–H bonds and open the door to multiple functionalizations through sequential deprotonation/coupling events.^[26]

Herein we disclose a general, high-yielding cross-coupling method between $\{\text{Cr}(\text{CO})_3\}$ -activated toluene derivatives and aryl bromides to afford a broad range of di- and triarylmethanes that are difficult or impossible to access by known methods.

Our initial studies used tricarbonylchromium-complexed diphenylmethane (**1**, Scheme 1) and 4-bromotoluene as coupling partners. Use of alkoxide bases at room temperature



Scheme 1. Coupling of diphenylmethane tricarbonylchromium **1** with aryl bromide.

resulted in no product (Table 1, entries 1 and 2). In contrast, $\text{LiN}(\text{SiMe}_3)_2$ combined with $[\text{PdCl}_2(\text{PPh}_3)_2]$ (3 mol %) afforded 70 % isolated yield of the coupled product **2a** after 20 h at room temperature (Table 1, entry 3) and 91 % yield in

Table 1: Optimization of reaction conditions (Scheme 1).

	Catalyst	mol %	Base	<i>t</i> [h] ^[a]	Yield [%] ^[b]	SM [%] ^[c]
1	$[\text{PdBr}_2(\text{PPh}_3)_2]$	10	LiOtBu	24	— ^[d]	—
2	$[\text{PdBr}_2(\text{PPh}_3)_2]$	5	NaOtBu	24	— ^[d]	67
3	$[\text{PdCl}_2(\text{PPh}_3)_2]$	3	$\text{LiN}(\text{SiMe}_3)_2$	20	70 ^[d]	trace
4	$[\text{PdCl}_2(\text{PPh}_3)_2]$	3	$\text{LiN}(\text{SiMe}_3)_2$	0.75	91	0
5	$[\text{PdCl}_2(\text{PPh}_3)_2]$	3	LDA	12	50	trace
6	$[\text{PdCl}_2(\text{PPh}_3)_2]$	5	$\text{NaN}(\text{SiMe}_3)_2$	18	26	38
7	$[\text{PdCl}_2(\text{PPh}_3)_2]$	5	$\text{KN}(\text{SiMe}_3)_2$	20	9	48
8	$[\text{NiCl}_2(\text{PPh}_3)_2]$	5	$\text{LiN}(\text{SiMe}_3)_2$	20	0	29
9	$[\text{Pd}(\text{PPh}_3)_4]$	5	$\text{LiN}(\text{SiMe}_3)_2$	17	67	12
10	$[\text{PdCl}_2(\text{dppf})]$	5	$\text{LiN}(\text{SiMe}_3)_2$	19	78	14

[a] Conducted at 55–60 °C in THF solvent, except where otherwise noted.

[b] Yields of isolated products. [c] Recovered starting material. [d] Reaction conducted at room temperature.

45 min at 60 °C (Table 1, entry 4). On the other hand, stronger amide bases such as lithium diisopropylamide (LDA), $\text{NaN}(\text{SiMe}_3)_2$, or $\text{KN}(\text{SiMe}_3)_2$ (Table 1, entries 5–7), or related catalysts, including $[\text{Cl}_2\text{Pd}(\text{dppf})]$ (Table 1, entries 8–10) were less effective. The optimized conditions in entry 4, Table 1, were used to determine the substrate scope (Table 2).

The diphenylmethane complex **1** readily undergoes cross-coupling reactions with a variety of aryl bromides in 81–94 % isolated yield of triarylmethane complexes **2a–2k** (Table 2).

[*] G. I. McGrew, J. Temaismithi, Dr. P. J. Carroll, Prof. P. J. Walsh
UPenn/Merck HTE Laboratories, Department of Chemistry
University of Pennsylvania
231 S. 34th St, Philadelphia, PA 19104 (USA)
Fax: (+1) 215-573-6743
E-mail: pwalsh@sas.upenn.edu
Homepage: <http://titanium.chem.upenn.edu/walsh/index.html>

[**] This research was supported by the National Institutes of Health, General Medical Sciences (GM058101), and National Science Foundation (CHE-0848467). J.T. is a Vagelos Scholar.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/ange.201000957>.


Table 2: Single coupling of **1** with ArBr (Scheme 1).^[a]

Entry	Ar =	Yield ^[b]	Entry	Ar =	Yield
2a		91	2f		91
2b		92	2g		94
2c		90	2h		91
2d		83	2j		81
2e		88	2k		89

[a] Conducted at 55–60 °C using 1.5 equiv LiN(SiMe₃)₂, 1.5 equiv ArBr, 3–5 mol % [PdCl₂(PPh₃)₂], and THF as solvent. TIPS = triisopropylsilyl.

Electron-withdrawing (**2d**, **2e**), -donating (**2f–2h**), and *ortho*-substituted aryl bromides (**2h–2k**) were all successful coupling partners. Of medicinal relevance, our method enabled installation of a 4-substituted indole (**2k**).^[27] In contrast, F-C chemistry strongly favors reaction of indole at the 3-position (or 2-position if the 3-position is blocked).^[5, 11–13, 17] Polyarylmethane derivatives bearing privileged indoles have attracted interest due to their widespread occurrence in biologically active compounds.

Next we examined reactions of [(η⁶-toluene)Cr(CO)₃] (**3**, Scheme 2). Although these reactions did not proceed well at room temperature, conversion to triarylmethane products

**Scheme 2.** Cross-coupling reactions with [(η⁶-toluene)Cr(CO)₃] (**3**).


5a–f (Table 3) was observed upon heating at 60 °C. Note that reaction with diarylmethane intermediate **4** is faster than the first coupling with **3**, probably due to the greater acidity of the benzylic C–H position of **4** over **3**.^[28] In contrast, bulky 2,6-disubstituted aryl bromides afforded monocoupled products **4g** and **4h**, likely due to steric hindrance.

Having demonstrated that two couplings could occur on a single methyl group, we wondered if one tricarbonylchromium center could activate multiple methyl groups in complexes of xylene and mesitylene. We were concerned that after the first methyl had undergone two arylations, the newly formed triarylmethane moiety would be deprotonated and the resulting anion would inhibit further deprotonation and coupling at the remaining methyl groups. As shown in Scheme 3 and Table 4, this was not a serious issue. Subjecting [(η⁶-*p*-xylene)Cr(CO)₃] (**6**) to cross-coupling conditions with the bulky 1-bromo-2,4,6-triisopropylbenzene furnished dicoupled **7** in 73% yield. Employing 4-bromofluorobenzene gave tetracoupled **8** in 70% yield. Furthermore, the mesitylene

Table 3: Double and single couplings of **3** with ArBr.^[a]

Entry	Ar =	Yield ^[b]	Entry	Ar =	Yield
5a		90	5e		86
5b		83	5f		88
5c		78	4g		86
5d		82	4h		81

[a] Conducted at 55–60 °C using 2.8 equiv LiN(SiMe₃)₂, 3 equiv ArBr, 4–8 mol % [PdCl₂(PPh₃)₂], and THF as solvent. [b] Yields of isolated products.

**Scheme 3.** Cross-coupling at multiple benzylic reaction sites. TMS = trimethylsilyl.

complex **9** was coupled with 4-bromoanisole (Scheme 3) resulting in formation of the hexacoupled product **10** in 43% yield (over 86% for each C–C bond-forming event). The structure of the hexacoupled product is illustrated in Figure 1.^[29] In the case of [(η⁶-*o*-xylene)Cr(CO)₃] (**11**), bulky 1-bromo-2,6-dimethylbenzene as coupling partner resulted in single arylation of each benzylic methyl to give the dicoupled product (**12**), in 85% yield. Use of the less sterically demanding aryl bromide 4-bromo-*N,N*-dimethylaniline, however, led to either unsymmetrical dicoupled product (**13**, in which both aryl groups couple to a single

Table 4: Coupling reaction at multiple arylation sites.^[a]

Starting complex	Product ^[b]	Starting complex	Product
	 Ar = 2,4,6-C ₆ H ₂ (iPr) ₃		 Ar = 2,6-C ₆ H ₃ Me ₂
	 Ar = <i>p</i> -C ₆ H ₄ -F		 Ar = 4-C ₆ H ₄ -NMe ₂
	 Ar = 4-C ₆ H ₄ -OMe		 Ar = 4-C ₆ H ₄ -NMe ₂

[a] Conducted at 50–60 °C using 4–11 equiv LiN(SiMe₃)₂, 3–10 equiv ArBr, 10–20 mol % [PdCl₂(PPh₃)₂], and THF as solvent. [b] Yields of isolated products.





Figure 1. ORTEP diagram (thermal ellipsoids at 50% probability) of hexa-arylated product (**10**).

methyl), or unsymmetrical tricoupled product (**14**) by varying the amount of base and reaction time. Steric barriers to both deprotonation and transmetalation likely account for the absence of tetraarylated (symmetrical) coupling product. These polyarylation reactions highlight a significant advantage of this approach over others: the possibility of multiple coupling reactions.^[16]


We also examined the possibility of coupling in the presence of α -heteroatoms and β -hydrogens using tricarbonylchromium-coordinated benzyl ether (**15**), *N,N*-dimethylbenzylamine (**16**), and ethylbenzene (**17**). Benzyl ether complex **15** underwent coupling employing conditions similar to those in Scheme 2 to afford monocoupled products **18a–e** in 58–80% yield (Scheme 4). Surprisingly, further reaction, whether in the presence of the original aryl bromide or with a different aryl bromide after isolation, was found to give triarylmethyl ether complexes **19a–f**. *N,N*-Dimethylbenzylamine complex **16** underwent single coupling to form diaryl-



Scheme 4. Coupling in the presence of heteroatoms and β -H atoms. [a] Product obtained directly from **15**. [b] From isolated **18b**. [c] From isolated **18a**. [d] 6% doubly coupled product was also isolated.

amines **20a–d** in 68–82% yield. No doubly coupled products were observed, possibly due to larger size of the dimethylamino group over the *n*-propyl ether in **15**. It is noteworthy that diarylmethylamines are important pharmacophores and are present in numerous medications.^[30] Ethylbenzene complex **17** underwent single-coupling to generate the 1,1-diarylethane derivative **21** in 70% yield, despite the presence of β -hydrogens in the intermediate palladium complex. Interestingly, this reaction also produced 6% of the triarylmethane quaternary product **22**.

To exploit of the ability of $\{\text{Cr}(\text{CO})_3\}$ to shield one face of the coordinated arene,^[21] we also examined cross-coupling reactions with complexes of indane (**23**) and tetrahydronaphthalene (**24**) (Scheme 5). By adjusting the conditions, either




Scheme 5. Access to 1-arylindanes and 1,3-*cis*-diarylindanes. [a] 40:60 toluene/THF was used instead of pure THF to improve selectivity for monoarylation. [b] From isolated **25b**.

mono- or diarylated product could be obtained, though yields of **25** and **26** were diminished due to partial conversion to diarylated product. In the case of the diarylation reactions to form **27** and **28**, *cis*-diaryl products were obtained in all cases (determined by ¹H NMR and X-ray diffraction of **27b**,^[29] see Supporting Information).

To demonstrate the practical advantages to our method, double coupling of indane tricarbonylchromium (**23**) with PhBr, followed by exposure to light and air provided 79% isolated yield of *cis*-1,3-diphenylindane (**29**, Scheme 6). Existing syntheses of 1,3-diaryl indanes provide *cis/trans* mixtures.^[31] The *cis*-1,3-diarylindane moiety is present in potent nonpeptide endothelin receptor antagonists.^[2,3] Performing the coupling of $[(\eta^6\text{-toluene})\text{Cr}(\text{CO})_3]$ (**3**) with 3-bromoanisole on a 5 mmol scale, followed by exposure to air and light afforded 1.4 g PhCH(3-C₆H₄-OMe)₂ (**30**) in 93% yield, a product inaccessible by standard F-C procedures with anisole.^[32]

In summary, we have introduced an efficient method for benzylic coupling reactions leading to polyarylated methanes. This method is complementary to F-C approaches and enables the synthesis of polyarylmethanes that are not accessible through electrophilic aromatic substitution reac-



Scheme 6. One-pot coupling and direct decomplexation of $[(\eta^6\text{-arene})\text{Cr}(\text{CO})_3]$ complexes. [a] Reaction is exposed to light prior to purification. [b] Single (*cis*-) isomer observed. [c] Reaction was conducted on a 5 mmol scale.

tions alone. Activation of the benzylic protons of $[(\eta^6\text{-arene})\text{Cr}(\text{CO})_3]$ allows a base of moderate strength, $\text{LiN}(\text{SiMe}_3)_2$, to be used for the in situ generation of the benzyllithiums,^[33] which directly participate in a palladium-catalyzed coupling reaction. Coordinated complexes can undergo multiple arylations, furnishing polyarylated products including unsymmetrically substituted triarylmethanes with quaternary carbon centers. Currently, efforts are underway to broaden the application of our method to the generation of quaternary polyarylmethanes and their enantioselective syntheses.

Received: February 16, 2010

Revised: May 4, 2010

Published online: July 2, 2010

Keywords: C–C coupling · chromium · organolithium · palladium · triarylmethanes

- [1] R. Palchaudhuri, V. Nesterenko, P. J. Hergenrother, *J. Am. Chem. Soc.* **2008**, *130*, 10274.
- [2] J. D. Elliott, M. A. Lago, R. D. Cousins, A. Gao, J. D. Leber, K. F. Erhard, P. Nambi, N. A. Elshourbagy, C. Kumar, *J. Med. Chem.* **1994**, *37*, 1553.
- [3] E. H. Ohlstein, P. Nambi, S. A. Douglas, R. M. Edwards, M. Gellai, A. Lago, J. D. Leber, R. D. Cousins, A. M. Gao, J. S. Frazee, C. E. Peishoff, J. W. Bean, D. S. Eggleston, N. A. Elshourbagy, C. Kumar, J. A. Lee, T. L. Yue, C. Loudon, D. P. Brooks, J. Weinstock, G. Feuerstein, G. Poste, R. R. Ruffolo, J. G. Gleason, J. D. Elliott, *Proc. Natl. Acad. Sci. USA* **1994**, *91*, 8052.
- [4] R. Muthyala, A. R. Katritzky, X. Lan, *Dyes Pigm.* **1994**, *25*, 303.
- [5] A. R. Katritzky, V. Gupta, C. Garot, C. V. Stevens, M. F. Gordeev, *Heterocycles* **1994**, *38*, 345.
- [6] D. F. Duxbury, *Chem. Rev.* **1993**, *93*, 381.
- [7] M. S. Shchepinov, V. A. Korshun, *Chem. Soc. Rev.* **2003**, *32*, 170.
- [8] V. Nair, S. Thomas, S. C. Mathew, K. G. Abhilash, *Tetrahedron* **2006**, *62*, 6731.
- [9] S. Lin, X. Lu, *J. Org. Chem.* **2007**, *72*, 9757.
- [10] S. K. Das, Shagufta, G. Panda, *Tetrahedron Lett.* **2005**, *46*, 3097.
- [11] J. S. Yadav, B. V. S. Reddy, S. Sunitha, *Adv. Synth. Catal.* **2003**, *345*, 349.
- [12] C. Ramesh, J. Barerjee, R. Pal, B. Das, *Adv. Synth. Catal.* **2003**, *345*, 557.
- [13] V. Nair, K. G. Abhilash, N. Vidya, *Org. Lett.* **2005**, *7*, 5857.
- [14] For a complementary procedure involving vicarious nucleophilic substitution of nitroarenes see: A. R. Katritzky, D. Toader, *J. Org. Chem.* **1997**, *62*, 4137.
- [15] A. R. Katritzky, X. Lan, J. N. Lam, *Chem. Ber.* **1991**, *124*, 1809.
- [16] Y.-Z. Li, B.-J. Li, X.-Y. Lu, S. Lin, Z.-J. Shi, *Angew. Chem.* **2009**, *121*, 3875; *Angew. Chem. Int. Ed.* **2009**, *48*, 3817.
- [17] J. Esquivias, R. G. Arrayás, J. C. Carretero, *Angew. Chem.* **2006**, *118*, 645; *Angew. Chem. Int. Ed.* **2006**, *45*, 629.
- [18] G. A. Molander, N. Ellis, *Acc. Chem. Res.* **2007**, *40*, 275.
- [19] B. Liégault, J.-L. Renaud, C. Bruneau, *Chem. Soc. Rev.* **2008**, *37*, 290.
- [20] D. Imao, B. W. Glasspoole, V. S. Laberge, C. M. Crudden, *J. Am. Chem. Soc.* **2009**, *131*, 5024.
- [21] M. Uemura in *Organic Reactions*, Vol. 67 (Ed.: L. E. Overman), Wiley, New York, **2006**, p. 217.
- [22] C. A. Merlic, J. C. Walsh, D. J. Tantillo, K. N. Houk, *J. Am. Chem. Soc.* **1999**, *121*, 3596.
- [23] A. Pfletschinger, T. K. Dargel, J. W. Bats, H.-G. Schmalz, W. Koch, *Chem. Eur. J.* **1999**, *5*, 537.
- [24] E. P. Küding in *Topics in Organometallic Chemistry*, Vol. 7, Springer, Berlin, **2004**.
- [25] V. N. Kalinin, I. A. Cherepanov, S. K. Moiseev, *J. Organomet. Chem.* **1997**, *536–537*, 437.
- [26] A. R. E. Brewer, A. F. Drake, S. E. Gibson, J. T. Rendell, *Org. Lett.* **2007**, *9*, 3487.
- [27] R. J. Sundberg, *Indoles*, Academic Press, San Diego, **1996**.
- [28] When ^1H NMR deprotonation studies were conducted in THF with 1 equiv $\text{LiN}(\text{SiMe}_3)_2$, **1** exhibited approximately 50% deprotonation, whereas **3** was only 10% deprotonated.
- [29] CCDC 775771 (**10**) and 765716 (**27b**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [30] K.-i. Yamada, K. Tomioka, *Chem. Rev.* **2008**, *108*, 2874.
- [31] B. Lantaño, J. Aguirre, M. L. Finkielstein, E. Alesso, E. Brunet, G. Y. Moltrasio, *Synth. Commun.* **2004**, *34*, 625.
- [32] Alternatively, the coordinated arene can be purified first and then simply decomplexed by exposure to light and air. A solution of coordinated arene **2a** exposed to light and air provided triphenylmethane in 98% isolated yield.
- [33] For cross-coupling reactions with organolithium reagents see: S.-I. Murahashi, *J. Organomet. Chem.* **2002**, *653*, 27.