


Platinum and Palladium Complexes Containing Cationic Ligands as Catalysts for Arene H/D Exchange and Oxidation**

Marion H. Emmert, J. Brannon Gary, Janette M. Villalobos, and Melanie S. Sanford*

The direct functionalization of C–H bonds has frequently been deemed a “Holy Grail” of organometallic chemistry.^[1] A seminal example of this transformation was the demonstration by Shilov and co-workers that platinum(II) salts catalyze the direct oxidation of alkanes into their corresponding alcohols and alkyl halides.^[2] Subsequent work in this area has focused on surveying diverse ligands for these reactions in an effort to enhance reactivity and selectivity, slow catalyst decomposition, and replace platinum(IV)-based oxidants with more cost-effective alternatives.^[3–5] In a key development, chemists at Catalytica identified [bpymPtCl₂] (bpym = bipyrimidine) as a pre-catalyst for the oxidation of CH₄ into CH₃OSO₃H in fuming H₂SO₄ (Scheme 1).^[5] The reaction




Scheme 1. Oxidative functionalization of methane with pre-catalyst **1**.

medium is believed to play several key roles in this system, including acting as a solvent, oxidant, and catalyst activator. Computational studies suggest that the active catalyst (**1**-H⁺) is formed by in-situ protonation of the ligand backbone, which limits oxidative catalyst degradation and renders the platinum center highly electrophilic and reactive for C–H bond cleavage.^[6]

Despite this initial success, it remains challenging to develop new generations of Group 10 metal catalysts that display higher turnover frequencies and operate in different, less corrosive media.^[3,4] A key goal of our efforts has been to identify ligands that mimic the desirable properties of protonated bipyrimidine but are stable in the absence of

strong acids.^[7] Herein, we describe the application of dicationic-bipyridine-based ligands in platinum- and palladium-catalyzed arene H/D exchange and oxidation reactions.

We considered dicationic ligands of general structure **2** (Scheme 2) for several key reasons. First, they contain



Scheme 2. Synthesis of platinum and palladium complexes **3a–c**. cod = cycloocta-1,5-diene, dmsO = dimethyl sulfoxide.

electron-withdrawing quaternized nitrogen substituents, which should render coordinated metal centers highly electrophilic.^[8] Second, they are bidentate, sp² N-donors, which should allow a direct comparison with other bipyrimidine and bipyridine ligand systems. Third, the quaternized nitrogen atoms are not susceptible to decomposition by dealkylation or deprotonation, which has been problematic in related systems.^[5,7] Fourth, substituents can easily be added to the pyridinium ring to tune the solubility of these dications. Finally, ligands **2a** and **2b** are readily available in four steps from commercially available 2,2'-bipyridine (bpy) in 33 % and 24 % overall yield, respectively (for details, see the Supporting Information).^[9]

Platinum(II) complexes of **2a** and **2b** were synthesized by reaction of these ligands with [(dmsO)₂PtCl₂] in methanol at 60 °C to afford **3a** and **3b** in 66 % and 52 % yield, respectively, after recrystallization (Scheme 2). The analogous palladium(II) complex **3c** was prepared in a similar fashion by the reaction of **2a** with [(cod)PdCl₂] in dichloromethane at room temperature (84 % yield; Scheme 2). All of these complexes were fully characterized by ¹H, ¹³C, and ¹⁹F NMR spectroscopy and elemental analysis.

We first evaluated **3a–c** as catalysts for H/D exchange^[10] between [D₄]acetic acid and benzene using an assay developed in our laboratory.^[10d] Under our standard conditions, (2 mol % [M], 4 mol % AgBF₄, 1 equiv C₆H₆ in 25 equiv of [D₄]AcOH, 150 °C), these complexes showed very high catalytic activity, with turnover frequencies (TOFs) of 0.1 s^{–1} (**3a,b**) and 0.05 s^{–1} (**3c**) after 15 minutes at 150 °C. For comparison, [dtbpyPtCl₂] (dtbpy = 4,4'-di-*tert*-butylbipyridine)

[*] Dr. M. H. Emmert, J. B. Gary, Dr. J. M. Villalobos, Prof. M. S. Sanford
Department of Chemistry, University of Michigan
930 N. University Ave, Ann Arbor, MI 48109 (USA)
Fax: (+1) 734-647-4865
E-mail: mssanfor@umich.edu

[**] We thank the NSF for support of this work through the Center for Enabling New Technologies through Catalysis. M.H.E. thanks the DFG (German Research Council) for a postdoctoral fellowship (EM 219/1-1). J.B.G. thanks the NSF for a predoctoral fellowship.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201002351>.

idine), [dtbpyPdCl₂], and bipyrimidine catalyst **1** provided TOFs of 0.0002, 0.002, and 0.003 s⁻¹, respectively, under identical conditions.^[10d] The catalysts were also compared on the basis of turnover numbers (TONs). Complexes **3a–c** achieved the statistical maximum TON of 242 after 24 hours,^[11,12] which is also superior to the results with [dtbpyPtCl₂], [dtbpyPdCl₂], and **1** (TON = 144 ± 12, 90 ± 18, and 94 ± 13 under analogous conditions).^[10d] Lowering the catalyst loading of **3a** to 0.1 mol% resulted in turnovers of 3273 ± 110 after 48 hours at 150 °C, which demonstrates the high activity and stability of this species, even upon prolonged heating at elevated temperatures. Finally, we decreased the reaction temperature to 100 °C. As shown in Figure 1, [dtbpyPtCl₂], [dtbpyPdCl₂], and **1** promoted very little H/D exchange under these conditions, whilst **3a–c** maintained high activity.




Figure 1. Turnover numbers for H/D exchange between benzene and [D₄]AcOH at 100 °C, catalyzed by **1**, **3a–c**, [dtbpyPtCl₂], and [dtbpyPdCl₂] after 2 h (white), 24 h (light gray), and 48 h (dark gray). Conditions: catalyst (2 mol%, 5 μmol), benzene (22.3 μL, 0.250 mmol), AgBF₄ (1.9 mg, 10 μmol), [D₄]AcOH (0.37 mL, 6.5 mmol, 25 equiv relative to benzene).

There are two possible explanations for the high H/D exchange activity of **3a–c**. The first is that the cationic ligands enhance the reactivity of the coordinated metal center towards arene C–H cleavage to generate metal σ-aryl intermediates (through an organometallic mechanism). An alternative possibility is that the Lewis acidic metal centers promote proton catalysis through an electrophilic aromatic substitution (Ar–S_E) pathway. To gain insight into the mechanism for H/D exchange, we examined the site-selectivity of D incorporation with catalyst **3a** for a variety of substituted arenes at partial (ca. 25%) conversion. Table 1 compares the observed selectivity (as determined by ¹H NMR spectroscopy) to the product ratio for electrophilic bromination^[13,14] (Table 1, entries 1–3); furthermore, the selectivity of DCl/CF₃CO₂D-catalyzed H/D exchange with bromobenzene is shown in entry 4. In all cases, the site selectivity with **3a** differed dramatically from that observed in Ar–S_E. This is strongly suggestive of an organometallic pathway for the platinum/palladium-catalyzed H/D exchange reactions.

Table 1: Comparison of site-selectivity for arene H/D exchange^[a,b] at 25% conversion with site-selectivity obtained in electrophilic aromatic bromination.

Entry	R	H/D exchange ortho/meta/para (catalyst)	Ar–S _E ortho/meta/para
1	Et	1.70:1.33:1 (3a) ^[a]	1.72:0.06:1 ^[c]
2	CO ₂ Et	1.26:2:1.33 (3a) ^[a]	0:2:0 ^[d]
3	Br	2.92:1.19:1 (3a) ^[a]	0.48:0.01:1 ^[c]
4	Br	1.15:0.28:1 (DCl/THF) ^[b]	0.48:0.01:1 ^[c]

[a] Conditions: **3a** (3.0 mg, 2.5 μmol, 0.5 mol%), AgBF₄ (1.9 mg, 5.0 μmol, 1.0 mol%), RPh (0.50 mmol, 1.0 equiv), [D₄]AcOH (0.71 mL, 25 equiv), 150 °C. [b] Conditions: BrPh (26.3 μL, 39.3 mg, 0.250 mmol, 1.00 equiv), [D₁]TFA (0.51 mL, 0.719 g, 6.25 mmol, 25.0 equiv), DCl in D₂O (35%, 0.05 mL), 38 h, 150 °C. [c] See Ref. [13]. [d] See Ref. [14]. TFA = trifluoroacetic acid, THF = tetrahydrofuran.

The substrate scope of H/D exchange reactions catalyzed by **3a** was also investigated. As summarized in Figure 2, naphthalene, veratrole, 1,2-dichlorobenzene, bromobenzene, ethylbenzoate, (H₃C)₃CCH₂Ph, *sec*-butylbenzene, cumene,




Figure 2. Substrate scope of H/D exchange catalyzed by **3a**; numbers in small font are %D incorporation. Conditions: **3a** (3.0 mg, 2.5 μmol, 2.0 mol%), substrate (0.125 mmol, 1.00 equiv), AgOTf (5.0 μmol, 4.0 mol%), [D₄]AcOH (0.36 mL, 0.40 g, 6.25 mmol, 50 equiv), 48 h (naphthalene, veratrole, PhBr) or 168 h (other substrates). Tf = trifluoromethanesulfonyl.

and trifluorotoluene all underwent extensive (48–98%) aromatic H/D exchange with CD₃CO₂D. Significant (2–8%) deuteration was also observed at unactivated sp³ C–H sites along the alkyl chains. In an interesting contrast, substrates that do not bear a tethered aromatic group (e.g. cyclooctane, 2,2,3,3-tetramethylbutane, and methane) did not show H/D exchange reactivity with catalyst **3a** under analogous conditions. This observation suggests that the arene moiety plays a role in directing the catalyst to the unactivated sp³ sites, likely via cyclometalation^[15] or π-coordination.^[16] The observation of aliphatic H/D exchange provides further evidence to support an organometallic mechanism, as this side reaction is expected to be negligible under proton catalysis.

The H/D-exchange experiments probed the reactivity of **3a–c** in C–H bond cleavage, which is only the first step of a

potential C–H functionalization process. Thus, it was important to determine whether the high reactivity and stability of these new complexes for arene H/D exchange correlated to catalytic activity in arene oxidation. To test this, we conducted preliminary studies of the effect of ligand **2a** on the Pd(OAc)₂-catalyzed acetoxylation of arenes with PhI(OAc)₂.^[17,18] As illustrated in Figure 3, the use of Pd(OAc)₂/




Figure 3. Pd^{II} catalyzed acetoxylation of benzene. Catalysts [Pd]: (◆) Pd(OAc)₂/**2a** 2:1; (☆) Pd(OAc)₂/bpy (2:1); (×) Pd(OAc)₂; (□) Pd(OAc)₂/bpym (2:1).

2a for the oxidation of benzene resulted in a significantly enhanced reaction rate compared to the best catalysts reported to date for this transformation (Pd(OAc)₂^[17] or Pd(OAc)₂/bpy 2:1^[18]). Similarly enhanced turnover frequencies were obtained in the C–H acetoxylation of naphthalene, 1,2-dichlorobenzene, chlorobenzene, bromobenzene, ethylbenzoate, and α,α -trifluorotoluene using Pd(OAc)₂/**2a** (for full details, see the Supporting Information). These initial results show that ligand **2a** can be used to generate robust, highly active palladium C–H oxidation catalysts.

In conclusion, the application of cationic pyridinium substituted ligands of general structure **2** for Group 10 C–H functionalization catalysis has been described. Both platinum and palladium complexes of these ligands display high catalytic activity for arene H/D exchange; furthermore, the combination of Pd(OAc)₂/**2a** shows enhanced activity for arene acetoxylation, compared with the best previously reported catalysts.^[17,18] Further applications of late transition metal complexes of **2a** and **2b** in C–H functionalization reactions are currently underway in our laboratory.

Received: April 20, 2010

Revised: June 1, 2010

Published online: July 15, 2010

Keywords: C–H activation · H/D exchange · ligand design · oxidation · platinum

- [1] B. A. Arndtsen, R. G. Bergman, T. A. Mobley, T. H. Peterson, *Acc. Chem. Res.* **1995**, *28*, 154–162.
- [2] a) N. F. Goldshleger, V. V. Eskova, A. E. Shilov, A. A. Shteinmann, *Zh. Fiz. Khim.* **1972**, *46*, 1353–1364; b) A. E. Shilov, G. B. Shulpin, *Chem. Rev.* **1997**, *97*, 2879–2932.
- [3] A. F. Heyduk, H. A. Zhong, J. A. Labinger, J. E. Bercaw, *Activation and Functionalization of C–H Bonds*, ACS Symp. Ser. **2004**, *885*, pp. 250–264.
- [4] For examples, see: a) M. Muehlhofer, T. Strassner, W. A. Herrmann, *Angew. Chem.* **2002**, *114*, 1817–1819; *Angew. Chem. Int. Ed.* **2002**, *41*, 1745–1747; b) I. Bar-Nahum, A. M. Khenkin, R. Neumann, *J. Am. Chem. Soc.* **2004**, *126*, 10236–10237; c) D. Meyer, M. A. Taïge, A. Zeller, K. Hohlfield, S. Ahrens, T. Strassner, *Organometallics* **2009**, *28*, 2142–2149; d) G. S. Chen, J. A. Labinger, J. E. Bercaw, *Organometallics* **2009**, *28*, 4899–4901; e) R. Palkovits, M. Antonietti, P. Kuhn, A. Thomas, F. Schüth, *Angew. Chem.* **2009**, *121*, 7042–7045; *Angew. Chem. Int. Ed.* **2009**, *48*, 6909–6912; f) J. E. Kreutz, A. Shukhaev, W. Du, S. Druskin, O. Daugulis, R. F. Ismagilov, *J. Am. Chem. Soc.* **2010**, *132*, 3128–3132.
- [5] R. A. Periana, D. J. Taube, S. Gamble, H. Taube, T. Satoh, H. Fujii, *Science* **1998**, *280*, 560–564.
- [6] a) J. Kua, X. Xu, R. A. Periana, W. A. Goddard, *Organometallics* **2002**, *21*, 511–525; b) X. Xu, J. Kua, R. A. Periana, W. A. Goddard, *Organometallics* **2003**, *22*, 2057–2068; c) A. Paul, C. B. Musgrave, *Organometallics* **2007**, *26*, 793–809; d) M. Ahlquist, R. A. Periana, W. A. Goddard, *Chem. Commun.* **2009**, 2373–2373.
- [7] J. M. Villalobos, A. J. Hickman, M. S. Sanford, *Organometallics* **2010**, *29*, 257–262.
- [8] For example, the Hammett σ_{para} value for NMe₃⁺ is 0.96.
- [9] P. P. Lainé, I. Ciofini, P. Ochsenbein, E. Amouyal, C. Adamo, F. Bedioui, *Chem. Eur. J.* **2005**, *11*, 3711–3727; and the references therein.
- [10] For examples, see: a) K. J. H. Young, S. K. Meier, J. M. Gonzales, J. Oxgaard, W. A. Goddard, R. A. Periana, *Organometallics* **2006**, *25*, 4734–4737; b) V. R. Ziatdinov, J. Oxgaard, O. A. Mironov, K. J. H. Young, W. A. Goddard, R. A. Periana, *J. Am. Chem. Soc.* **2006**, *128*, 7404–7405; c) G. Gerdes, P. Chen, *Organometallics* **2004**, *23*, 3031–3036; d) A. J. Hickman, J. M. Villalobos, M. S. Sanford, *Organometallics* **2009**, *28*, 5316–5322.
- [11] All reported TONs and TOFs are corrected for the background H/D exchange reaction of AgCl, which is formed as a by-product under the reaction conditions.
- [12] For TONs obtained after 2 h and 24 h at 2, 1, 0.5, 0.25, and 0.1 mol % catalyst loading (**3a**), see the Supporting Information.
- [13] H. Gilow, *J. Chem. Educ.* **1977**, *54*, 450–452.
- [14] S. Rozen, M. Brand, *J. Chem. Soc. Chem. Commun.* **1987**, 752–753.
- [15] T. G. Driver, M. W. Day, J. A. Labinger, J. E. Bercaw, *Organometallics* **2005**, *24*, 3644–3654.
- [16] N. Ito, H. Esaki, T. Maesawa, E. Imamiya, T. Maegawa, H. Sajiki, *Bull. Chem. Soc. Jpn.* **2008**, *81*, 278–286.
- [17] T. Yoneyama, R. H. Crabtree, *J. Mol. Catal. A* **1996**, *108*, 35–40.
- [18] a) L. Eberson, L. Jönsson, *J. Chem. Soc. Chem. Commun.* **1974**, 885–886; b) L. Eberson, L. Jönsson, *Acta Chem. Scand. Ser. B* **1976**, *30*, 361–364.