



Auf diesen drei Seiten weisen wir auf wichtige aktuelle Beiträge in unseren Schwesterzeitschriften hin. Wenn Sie die Seiten online lesen, dann können Sie

die Artikel mit einem Klick direkt aufrufen, ansonsten sind sie durch Eingabe der DOIs über Wiley Online Library leicht online zugänglich.




Fluorescent Nanosensors

T. Doussineau,* A. Schulz, A. Lapresta-Fernandez, A. Moro, S. Körsten, S. Trupp, G. J. Mohr*

On the Design of Fluorescent Ratiometric Nanosensors

Small but well-grounded! Recently developed nanoparticle-based sensors (see figure) appear as new powerful tools to provide quantitative information about different species (ions, metabolites, biomolecules) in biosamples through ratiometric measurements. This article describes some of the methodologies developed so far in the design of such nanosensors.



Chem. Eur. J.
DOI: 10.1002/chem.201000829




Nanotubes

L. Meng,* C. Fu, Z. Fei,* Q. Lu,* P. J. Dyson

Photochemical Behavior of High Quantum Yield SWNTs Functionalized with Anthracene Salts

I hope you like bamim too: Anthracene derivatives attached to functionalized single-walled carbon nanotubes through Coulombic interactions undergo a partial (and reversible) photodimerization process upon irradiation with UV light of appropriate wavelengths.



Chem. Asian J.
DOI: 10.1002/asia.201000236




Oligomers


A. A. Reinke, G. A. Abulwerdi, J. E. Gestwicki*

Quantifying Prefibrillar Amyloids in vitro by Using a "Thioflavin-Like" Spectroscopic Method

In a glass darkly: The formation of fibrillar amyloids is linked to many neurodegenerative diseases. However, recent models suggest an important role for prefibrillar amyloids. Here, we develop a facile, inexpensive assay for specifically detecting prefibrillar amyloids in opaque plates. Results obtained with this method suggest that prefibrillar amyloids are depleted during the aggregation process.



ChemBioChem
DOI: 10.1002/cbic.201000358





ChemPhysChem
DOI: 10.1002/cphc.201000210

DNA Hybridization

M. Gębala, W. Schuhmann*

Controlled Orientation of DNA in a Binary SAM as a Key for the Successful Determination of DNA Hybridization by Means of Electrochemical Impedance Spectroscopy

The composition of the monolayer, the ionic strength, the surface coverage with single-stranded DNA (ssDNA), and the electrode pretreatment are prerequisites for reliably detecting DNA hybridization using electrochemical impedance spectroscopy. Co-assembly of mercaptopropionic acid is proposed as a means to control the electrode/electrolyte interface (see figure).





ChemMedChem
DOI: 10.1002/cmdc.201000184

Neuroprotection

P. Conti,* A. Pinto, L. Tamborini, U. Madsen, B. Nielsen, H. Bräuner-Osborne, K. B. Hansen, E. Landucci, D. E. Pellegrini-Giampietro, G. De Sarro, E. Donato Di Paola, C. De Micheli

Novel 3-Carboxy- and 3-Phosphonopyrazoline Amino Acids as Potent and Selective NMDA Receptor Antagonists: Design, Synthesis, and Pharmacological Characterization

Calming the storm: New NMDA receptor antagonists and their pharmacological activity toward various types and subtypes of glutamate receptors are described. Among these derivatives, two compounds exhibit promising in vitro and in vivo pharmacological activity as neuroprotective and anticonvulsant agents.





ChemSusChem
DOI: 10.1002/cssc.200900293

Carbon Dioxide Capture

S. Zhou, X. Chen, T. Nguyen, A. K. Voice, G. T. Rochelle*

Aqueous Ethylenediamine for CO₂ Capture

Aqueous ethylenediamine (EDA) is investigated as a solvent for CO₂ capture from flue gas. Similar to monoethanolamine, EDA can be used up to 120 °C in a stripper without significant thermal degradation. The results indicate that EDA is a suitable choice for capturing CO₂.



ChemCatChem
DOI: 10.1002/cctc.201000112

Biocatalysis

M. Anbar, R. Lamed, E. A. Bayer*

Thermostability Enhancement of *Clostridium thermocellum* Cellulosomal Endoglucanase Cel8A by a Single Glycine Substitution

Thermostable mutant enzymes: The X-ray crystal structure of the *Clostridium thermocellum* Cel8A cellulase depicts the amino acid substitutions which enable the protein to withstand elevated temperatures while maintaining wild-type levels of activity. The mutant enzyme may be used in combination with other selected glycoside hydrolases to efficiently degrade lignocellulosic materials used in the biofuel industry.






Spirocyclic Silicon Clusters

T. M. Klapötke,* S. K. Vasisht,* P. Mayer

Spirocycle $(\text{Si}t\text{Bu}_3)_6\text{Si}_9\text{Cl}_2$: The First of Its Kind among Group 14 Elements

An Si_4 cage stabilizes itself in a heavily loaded environment in tetrasilatetrahedranide KR^*Si_4 ($\text{R}^* = \text{Si}t\text{Bu}_3$), but steric congestion in $(\text{R}^*\text{Si}_4)_2\text{SiCl}_2$ results in rearrangement leading to cage opening into less strained tricyclopentasilane rings in $(\text{R}^*\text{Si}_4\text{Cl})_2\text{Si}$, and the molecule acquires a spiro[4.4]nonasilane structure.



Eur. J. Inorg. Chem.
DOI: 10.1002/ejic.201000498




Molecules for Switchable Monolayers

J. Kubitschke, C. Näther, R. Herges*

Synthesis of Functionalized Triazatriangulenes for Application in Photo-Switchable Self-Assembled Monolayers

The synthesis of functionalized triazatriangulenes is presented and the *trans/cis* isomerization of azobenzene derivatives in solution is investigated. The described attachment of various molecular functions to the TATA platform represents a modular system for the formation of switchable SAMs.



Eur. J. Org. Chem.
DOI: 10.1002/ejoc.201000650

New Journal

Heterogeneous, Homogeneous and BioCatalysis

www.chemcatchem.org

FREE ONLINE ACCESS

In 2010 for all users from institutions that have registered

Ask your librarian to register for complimentary online access TODAY

www.interscience.wiley.com/newjournals

A Journal of

ChemPubSoc Europe

A journal of

Founding Societies:

WILEY-VCH