



Auf diesen drei Seiten weisen wir auf wichtige aktuelle Beiträge in unseren Schwesterzeitschriften hin. Wenn Sie die Seiten online lesen, dann können Sie

die Artikel mit einem Klick direkt aufrufen, ansonsten sind sie durch Eingabe der DOIs über Wiley Online Library leicht online zugänglich.




Heterocycles

M. D. Hill*

Recent Strategies for the Synthesis of Pyridine Derivatives

Closing the ring: Modification of traditional condensation strategies continues to be a recurrent theme in contemporary literature. Advancements in transition-metal-catalyzed cyclization and cross-coupling procedures offer new routes to functionalized pyridine derivatives (see scheme).



Chem. Eur. J.

DOI: 10.1002/chem.201001100




Carbaboration

C. Chen, F. Eweiner, B. Wibbeling, R. Fröhlich, S. Senda, Y. Ohki, K. Tatsumi, S. Grimme, G. Kehr, G. Erker*


Exploring the Limits of Frustrated Lewis Pair Chemistry with Alkynes: Detection of a System that Favors 1,1-Carbaboration over Cooperative 1,2-P/B-Addition

Born of frustration: The boryl part of a frustrated Lewis pair, attached at the zirconocene framework, reacted with a terminal alkyne by 1,1-carbaboration of the triple bond. The Lewis base plays a negligible role in this reaction, as shown on a phosphorous-free example.



Chem. Asian J.

DOI: 10.1002/asia.201000189




Biosynthesis

J. Zeng, J. Zhan*


A Novel Fungal Flavin-Dependent Halogenase for Natural Product Biosynthesis

A novel halogenating agent: A fungal halogenase Rdc2 from *Pochonia chlamydosporia* has been reconstituted in *E. coli* and biochemically characterized as the dedicated halogenase in radicicol biosynthesis. The enzyme displays broad substrate specificity towards molecules such as dihydroresorcylic acid (**1**) to generate mono- and dihalogenated (X = Br or Cl) derivatives, thus representing a promising biocatalyst for natural product biosynthesis.



ChemBioChem

DOI: 10.1002/cbic.201000439





ChemPhysChem
DOI: 10.1002/cphc.201000374

Gas-phase reactions

I. Barnes,* G. Solignac, A. Mellouki, K. H. Becker

Aspects of the Atmospheric Chemistry of Amides

Active space: The gas-phase reactions of six amides, formamide, *N*-methyl formamide, *N,N*-dimethyl formamide, acetamide, *N*-methyl acetamide and *N,N*-dimethyl acetamide with the atmospheric oxidants OH radicals and Cl atoms, but in a number of cases also with NO₃ radicals and ozone, are presented and discussed (see graphic).




ChemMedChem
DOI: 10.1002/cmdc.201000297

Drug Design

A. Fischer, C. Schmidt, S. Lachenicht, D. Grittner, M. Winkler, T. Wrobel, A. Rood, H. Lemoine, W. Frank, M. Braun*

Synthesis of Benzofuran, Benzothiophene, and Benzothiazole-Based Thioamides and their Evaluation as K_{ATP} Channel Openers

A recipe for success: Combine a hydrogen-bond donor and acceptor with sulfur atoms that provide a region of high electron density and lipophilic, bulky substituents in an appropriate manner, and you will obtain novel K_{ATP} channel openers with remarkable selectivity toward different isoforms of sulfonylurea receptors.




ChemSusChem
DOI: 10.1002/cssc.201000209

Biorenewables

T. Thananattachon, T. B. Rauchfuss*

Efficient Route to Hydroxymethylfurans from Sugars via Transfer Hydrogenation

Tandem catalysis, relying on formic acid as an acid catalyst and as a source of hydrogen, provides a promising route to highly pure furanyl-methanols. The new approach exploits (i) the use of DMSO to mediate highly efficient routes to furfurals and (ii) the ability of transfer hydrogenation catalysts to effect the hydrogenation, with good tolerance for DMSO.




ChemCatChem
DOI: 10.1002/cctc.201000081

Supported Catalysts

H. Wang, X. Li,* Y. M. Wang, P. Wu

Pt Nanoparticles Supported on Highly Dispersed Alumina Coated inside SBA-15 for Enantioselective Hydrogenation

Everything's aluminated: A series of Al₂O₃@SBA-15 composites with different Al₂O₃ loadings are prepared by a solvent-free solid-state grinding method. The composites retain the meso-structure of SBA-15 and the alumina is uniformly coated inside the mesopores. Furthermore, the Al₂O₃@SBA-15 composites serve as remarkable supports for Pt nanoparticle catalysts in the enantioselective hydrogenation of ethyl pyruvate.






Flexible Copper(II) MOFs


K. Sumida, M. L. Foo, S. Horike, J. R. Long*

Synthesis and Structural Flexibility of a Series of Copper(II) Azolate-Based Metal–Organic Frameworks

The reaction of $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$ with three novel ligands, 2-methyl-1,4-benzeneditrazolate (MeBDT^{2-}), 4,4'-biphenylditrazolate (BPD^{2-}), and 2,3,5,6-tetrafluoro-1,4-benzeneditrazolate (TFBD^{2-}), affords three metal–organic frameworks with a common network topology but a significantly different flexibility, highlighting the significant impact the ligand component can have on the dynamic properties of the material.



Eur. J. Inorg. Chem.
DOI: 10.1002/ejic.201000490




Natural Product Synthesis

S. Beaumont, E. A. Ilardi, N. D. C. Tappin, A. Zakarian*

Marine Toxins with Spiroimine Rings: Total Synthesis of Pinnatoxin A

This microreview provides a compilation of synthetic approaches and total syntheses of pinnatoxin A, a single representative member of a fascinating group of potent marine toxins that share a spiroimine subunit as a unifying structural element. The literature is surveyed up to early 2010.



Eur. J. Org. Chem.
DOI: 10.1002/ejoc.201000842

New Journal

Heterogeneous, Homogeneous and BioCatalysis

www.chemcatchem.org

FREE ONLINE ACCESS
In 2010 for all users from institutions that have registered
Ask your librarian to register for complimentary online access TODAY
www.interscience.wiley.com/newjournals

A Journal of

ChemPubSoc Europe

A journal of

Founding Societies:

WILEY-VCH