


Catalytic Asymmetric Dihydroxylation of Enamides and Application to the Total Synthesis of (+)-Tanikolide**

Benoit Gourdet and Hon Wai Lam*

Asymmetric transformations of 1,1-disubstituted alkenes provide important building blocks for chemical synthesis, but are often plagued with low stereoselectivities because it can be difficult for a chiral reagent or catalyst to discriminate between the enantiotopic faces of these substrates.^[1] Among the available methods, asymmetric dihydroxylation (AD) stands out as one of the more successful, and can provide high enantioselectivities in certain cases (Scheme 1, **1**→**2**).^[2,3] However, low enantioselectivities are usually observed when the two alkene substituents are of similar steric demand.^[4,5]




Scheme 1. Asymmetric dihydroxylation (AD) routes to tertiary-alcohol-containing terminal 1,2-diols **2**.

A possible solution to this problem is to employ β,β' -disubstituted enol derivatives **3** as the substrates, where discrimination of the enantiotopic faces is expected to be more straightforward (Scheme 1, **3**→**2**). An additional benefit of this approach is that asymmetric dihydroxylation results in chiral α -hydroxyaldehydes **4**, which are themselves valuable compounds, and which can be reduced to 1,2-diols **2** if required. To ensure high enantioselectivity in the dihydroxylation event, the substrate **3** must be obtained in high stereoisomeric purity.^[6] Unfortunately, existing methods to prepare these compounds typically proceed with poor *E/Z* stereoselectivity,^[7–9] and the resulting geometric isomers can be difficult to separate.

Partial solutions to these problems have been described recently.^[10] Ready and co-workers have developed stereocontrolled syntheses of β,β' -disubstituted enol esters and enol

silanes involving carbocupration^[10a] or carboalumination^[10b] of terminal alkynes, oxidation of the resulting alkenylmetal species using a metal *tert*-butyl peroxide, and trapping of the resulting enolate with an acylating or silylating agent.^[10] Furthermore, asymmetric dihydroxylation of enol benzoates containing methyl substitution on the alkene was demonstrated.^[10b] Nevertheless, improvements with these approaches can be envisaged. In the carbocupration procedure,^[10a,11] the organocopper reagents are generated from organolithium or Grignard reagents, which pose restrictions on the functional groups that may be present in the organometallic reagent. Although functionalized organocopper reagents may be obtained from the corresponding organozinc halides, these reagents are poorly reactive towards unactivated terminal alkynes.^[12] In addition, because only alkylcopper reagents exhibit sufficient reactivity in alkyne carbocupration, the introduction of important groups such as (hetero)aryl substituents is usually not possible. In the carboalumination procedure,^[10b,13] only methyl groups can be transferred. Therefore, full exploration of asymmetric dihydroxylation of enol derivatives **3** to access chiral α -hydroxyaldehydes **4** and diols **2** is compromised by these limitations.

Our research group has recently developed rhodium-catalyzed carbometalation reactions that offer solutions to many of these drawbacks.^[14] Instead of providing enol derivatives **3**, these reactions furnish β,β' -disubstituted enamides **6**^[15] from the corresponding ynamides **5**^[16] (Scheme 2). Collectively, these processes enable the introduction of alkyl,



Scheme 2. Asymmetric dihydroxylation of enamides prepared by rhodium-catalyzed carbometalation of ynamides.

alkenyl, aryl, heteroaryl, benzyl, and alkynyl groups, and the presence of sensitive functional groups such as esters^[14a–c] and ketones^[14c] on the organometallic reagent is permitted. Accordingly, asymmetric dihydroxylation of enamides **6** should provide access to a much wider range of chiral products than is possible using comparable methods.^[10b] To our knowledge, there are only limited reports of Sharpless asymmetric dihydroxylation of enamides, where cyclic substrates were oxidized with modest ($\leq 77\%$ *ee*) enantioselectivities.^[17] Herein, we report highly enantioselective dihy-

[*] B. Gourdet, Dr. H. W. Lam
School of Chemistry, University of Edinburgh
Joseph Black Building, The King's Buildings
West Mains Road, Edinburgh EH9 3JJ (UK)
Fax: (+44) 131-650-6453
E-mail: h.lam@ed.ac.uk
Homepage: <http://homepages.ed.ac.uk/hlam/index.html>

[**] This work was supported by the European Commission (Project No. MEST-CT-2005-020744) and the University of Edinburgh. We thank the EPSRC National Mass Spectrometry Service Centre at the University of Wales, Swansea, for providing high-resolution mass spectra.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201004328>.

droxylations of β,β' -disubstituted enamides, and the application of this method in the total synthesis of (+)-tanikolide.^[18]


The enamide employed in the initial part of this study are shown in Table 1, and were prepared according to one of the

Table 1: Catalytic asymmetric dihydroxylation of enamides.^[a]

Entry	Enamide	Product	Yield [%] ^[b]	ee [%] ^[c]
1	6a R = Me	2a	72	90 (–94) ^[d]
2	6b R = Et	2b	84	87 (–78) ^[d]
3	6c R = 2-thienyl	2c	61	94
4	6d Ar = Ph	2d	68	95
5	6e Ar = 3-EtO ₂ CC ₆ H ₄	2e	70	97
6	6f Ar = Ph	2f	83	97
7	6g Ar = 4-MeC ₆ H ₄	2g	82	97
8	6h Ar = 2-thienyl	2h	86	94
9	6i n-Hex	2i	68	96


[a] Reactions were conducted using **6a–6i** (0.20 mmol) and AD-mix- β (280 mg; 0.4 mol% of Os and 1 mol% of (DHQD)₂PHAL) in *t*BuOH (2 mL) and H₂O (1 mL) for 7–24 hours, and subsequent addition of NaBH₄ (6 equiv) and stirring for an additional 1 hour. [b] Yield of isolated product. [c] Determined by HPLC analysis on a chiral stationary phase. [d] Values in parentheses refer to *ee* values obtained from asymmetric dihydroxylation of the corresponding 1,1-disubstituted alkene using the same chiral ligand (DHQD)₂PHAL that is present in AD-mix- β , see Ref. [2c]. TBS = *tert*-butyldimethylsilyl.


rhodium-catalyzed ynamide carbometalation procedures we have described recently.^[14] The enamides **6a**, **6b**, and **6d–6g** have been reported previously,^[14] whereas enamides **6c**, **6h**, and **6i** are new compounds.^[19] Highly enantioselective dihydroxylation of these substrates was readily accomplished using commercially available AD-mix- β without the addition of MeSO₂NH₂.^[3] After dihydroxylation was complete, addition of NaBH₄ to the reaction mixture resulted in formation of the 1,2-diols **2**. The sense of enantiofacial selectivity by AD-mix- β in these reactions is consistent with the Sharpless mnemonic.^[3] Enamides **6a–6c** containing a phenyl substituent at the R¹ position were effective substrates, with a 2-thienyl group at R² (entry 3) providing a higher enantioselectivity than a methyl (entry 1) or an ethyl group (entry 2). Enamides **6d–6i** containing aliphatic groups (simple alkyl, oxygenated alkyl) at R¹ and (hetero)aryl groups (phenyl, *p*-tolyl, 3-carboethoxyphenyl, 2-thienyl) at R² proved to be especially competent substrates, providing products in generally good yields with uniformly high enantioselectivities (entries 4–9). Comparison of the present method with asymmetric dihydroxylation of 1,1-disubstituted alkenes is informative. While diol **2a** was obtained from enamide **6a** with slightly lower enantioselectivity (90% *ee*; entry 1) and as the opposite enantiomer compared with asymmetric dihydroxylation of α -methylstyrene under comparable reaction conditions (–94% *ee*),^[2c] the present method afforded diol **2b** with improved selectivity (87% *ee*; entry 2) compared with that obtained from the corresponding 1,1-disubstituted alkene (–78% *ee*).^[2c] Interestingly, diol **2d** was obtained favoring the *R* enantiomer from both enamide **6d** and alkene **1d**, but the enantioselectivity was far superior in the case of the enamide (95% *ee*; entry 4) compared with the alkene [41% *ee*; Eq. (1)].



The reaction of β,β' -diarylenamide **6j**^[19] with AD-mix- β was sluggish and did not proceed to completion, but dihydroxylation using increased loadings of K₂OsO₂(OH)₄ and the chiral ligand (DHQD)₂PHAL (hydroquinidine 1,4-phthalazinediyl diether), which is present in AD-mix- β , gave improved results and afforded diol **2j** in 98% *ee* [Eq. (2)].

Scheme 3 illustrates application of this method to the concise preparation of α -hydroxyaldehyde **4k**, a known intermediate in the synthesis of (*S*)-oxybutynin (**7**),^[20,21] which is widely used in the treatment for urinary problems.^[22] In this case, rhodium-catalyzed carbocyclization of ynamide **5a**






Scheme 3. Synthesis of α -hydroxyaldehyde **4k**, a precursor to (S)-oxybutynin (**7**). THF = tetrahydrofuran.

with Cy₂Zn (prepared from CyMgBr and ZnCl₂; Cy = cyclohexyl) according to our reported method^[14a,b] did provide enamide **6k**, but in low yield owing to the formation of (Z)-3-(2-styryl)oxazolidin-2-one as a major side product resulting from undesired hydrazination.^[14b] However, improved results were obtained from copper-catalyzed carbomagnesiation of **5a** using CyMgBr and [Cu(acac)₂] (acac = acetylacetonate), which provided **6k** in 74% yield.^[23] In similar fashion to **6j** [Eq. (2)], asymmetric dihydroxylation of **6k** with AD-mix- β was sluggish, but dihydroxylation using increased loadings of K₂OsO₂(OH)₄ and (DHQD)₂PHAL afforded α -hydroxyaldehyde **4k** in 80% yield and 91% ee. It should be noted that dihydroxylation of α -cyclohexylstyrene using (DHQD)₂PHAL provides the corresponding diol as the opposite *R* enantiomer in only 57% ee^[2c,24]—once again illustrating that use of β,β' -disubstituted enamides in place of 1,1-disubstituted alkenes can prove advantageous in asymmetric dihydroxylation reactions.

As a further exemplification of asymmetric enamide dihydroxylation, we applied this method to a concise total synthesis of the antifungal natural product (+)-tanikolide (**14**).^[18,25] First, bromoalkyne **9** was prepared from commercially available tridecyne **8** using NBS/AgNO₃ (Scheme 4).^[26] By using the method of Hsung and co-workers,^[27] copper-catalyzed coupling of **9** with oxazolidin-2-one then afforded ynamide **10**. Rhodium-catalyzed carbozincation of **10** using commercially available 4-ethoxy-4-oxobutylzinc bromide^[14a,b] proceeded smoothly and provided enamide **11** in 56% yield. Unfortunately, dihydroxylation of **11** using AD-mix- β was poorly selective, and after reduction of the initially formed aldehyde with NaBH₄, diol **12** was produced in 58% ee. However, use of (DHQD)₂AQN (hydroquinidine anthraquinone-1,4-diyl diether)^[28] as the chiral ligand in place of (DHQD)₂PHAL led to **12** being obtained in a much improved 88% ee.^[29] Basic hydrolysis of **12**, and subsequently heating the resulting acid **13** in CDCl₃ at reflux led to smooth lactonization and provided (+)-tanikolide (**14**). Overall, the route to (+)-tanikolide (**14**) presented in Scheme 4 is competitive with previous syntheses.^[25]

In summary, the asymmetric dihydroxylation of β,β' -disubstituted enamides has been described. By drawing upon



Scheme 4. Total synthesis of (+)-tanikolide (**14**). cod = cycloocta-1,5-diene, NBS = N-bromosuccinimide.

the diversity of β,β' -disubstituted enamide derivatives available using recently reported rhodium-catalyzed ynamide carbometallation protocols,^[14] and offering often superior enantioselectivity compared with asymmetric dihydroxylation of the corresponding 1,1-disubstituted alkenes, this strategy serves as a versatile method for the preparation of important 1,2-dioxygenated compounds. Many of these tertiary-alcohol-containing building blocks **2** and **4** would be difficult to access efficiently using alternative methods. Application of this process to the concise syntheses of the natural product (+)-tanikolide (**14**) and α -hydroxyaldehyde **4k**, an intermediate en route to (S)-oxybutynin (**7**), has been demonstrated.

Received: July 15, 2010

Published online: September 30, 2010

Keywords: asymmetric catalysis · dihydroxylation · enamides · total synthesis · ynamides

[1] For a useful discussion, see: S. P. Thomas, V. K. Aggarwal, *Angew. Chem.* **2009**, *121*, 1928–1930; *Angew. Chem. Int. Ed.* **2009**, *48*, 1896–1898.

[2] For example, see: a) K. J. Hale, S. Manaviyar, S. A. Peak, *Tetrahedron Lett.* **1994**, *35*, 425–428; b) H.; Yasui, H. Yorimitsu, K. Oshima, *Chem. Lett.* **2007**, *36*, 32–33; c) S. B. King, M. Taniguchi, K. P. M. Vanhessche, K. B. Sharpless, *J. Org. Chem.* **1995**, *60*, 3940–3941; d) K. P. M. Vanhessche, K. B. Sharpless, *J. Org. Chem.* **1996**, *61*, 7978–7979; e) X. Wang, M. Zak, M. Maddess, P. O'Shea, R. Tillyer, E. J. J. Grabowski, P. J. Reider, *Tetrahedron Lett.* **2000**, *41*, 4865–4869.

[3] For reviews, see: a) H. C. Kolb, M. S. VanNieuwenhze, K. B. Sharpless, *Chem. Rev.* **1994**, *94*, 2483–2547; b) M. C. Noe, M. A.

- Letavic, S. L. Snow, *Org. React.* **2005**, *66*, 109–625; c) A. B. Zaitsev, H. Adolfsson, *Synthesis* **2006**, 1725–1756.
- [4] Alternative catalytic asymmetric syntheses of terminal 1,2-dioxygenated compounds containing a tertiary alcohol at the 2-position usually involve addition of carbon nucleophiles to α -ketoesters. For selected examples, see: a) F. Wang, X. Liu, Y. Zhang, L. Lin, X. Feng, *Chem. Commun.* **2009**, 7297–7299; b) H.-F. Duan, J.-H. Xie, X.-C. Qiao, L.-X. Wang, Q.-L. Zhou, *Angew. Chem.* **2008**, *120*, 4423–4425; *Angew. Chem. Int. Ed.* **2008**, *47*, 4351–4353; c) K. Zheng, B. Qin, X. Liu, X. Feng, *J. Org. Chem.* **2007**, *72*, 8478–8483; d) T. Ooi, K. Fukumoto, K. Maruoka, *Angew. Chem.* **2006**, *118*, 3923–3926; *Angew. Chem. Int. Ed.* **2006**, *45*, 3839–3842; e) J.-R. Kong, M.-Y. Ngai, M. J. Krische, *J. Am. Chem. Soc.* **2006**, *128*, 718–719; f) L. C. Akullian, M. L. Snapper, A. H. Hoveyda, *J. Am. Chem. Soc.* **2006**, *128*, 6532–6533; g) L. C. Wieland, H. Deng, M. L. Snapper, A. H. Hoveyda, *J. Am. Chem. Soc.* **2005**, *127*, 15453–15456; h) B. Jiang, Z. Chen, X. Tang, *Org. Lett.* **2002**, *4*, 3451–3453; i) D. A. Evans, S. W. Tregay, C. S. Burgey, N. A. Paras, T. Vojtkovsky, *J. Am. Chem. Soc.* **2000**, *122*, 7936–7943; j) D. A. Evans, C. S. Burgey, M. C. Kozlowski, S. W. Tregay, *J. Am. Chem. Soc.* **1999**, *121*, 686–699.
- [5] For reviews detailing the catalytic asymmetric synthesis of tertiary cyanohydrins, which in many cases may be converted into 1,2-dioxygenated compounds by manipulation of the nitrile group, see: a) J.-M. Brunel, I. P. Holmes, *Angew. Chem.* **2004**, *116*, 2810–2837; *Angew. Chem. Int. Ed.* **2004**, *43*, 2752–2778; b) M. Hatano, K. Ishihara, *Synthesis* **2008**, 1647–1675; c) M. North, D. L. Usanov, C. Young, *Chem. Rev.* **2008**, *108*, 5146–5226.
- [6] This scenario contrasts with that for asymmetric dihydroxylation of ketone-derived enol ethers and enol silanes, where *E/Z* stereoisomeric purity of the substrate is often inconsequential, see: T. Hashiyama, K. Morikawa, K. B. Sharpless, *J. Org. Chem.* **1992**, *57*, 5067–5068.
- [7] For preparation of enol esters from α -substituted aldehydes, see: a) C. P. Casey, C. R. Cyr, *J. Am. Chem. Soc.* **1973**, *95*, 2240–2247; b) H. H. Wasserman, L. S. Keller, *Tetrahedron Lett.* **1974**, *15*, 4355–4358; c) T. Mukaiyama, M. Murakami, M. Yamaguchi, *Chem. Lett.* **1980**, 529–532; d) S. Kojima, S. Maki, T. Hirano, Y. Ohmiya, H. Niwa, *Tetrahedron Lett.* **2000**, *41*, 4409–4413.
- [8] For preparation of enol silanes from α -substituted aldehydes, see: a) R. C. Hartley, I. C. Richards, S. Warren, *J. Chem. Soc. Perkin Trans. 1* **1996**, 359–376; b) J. Hydrio, P. Van de Weghe, J. Collin, *Synthesis* **1997**, 68–72; c) Y. Ishino, Y. Kita, H. Maekawa, T. Ohno, Y. Yamasaki, T. Miyata, I. Nishiguchi, *Tetrahedron Lett.* **1999**, *40*, 1349–1352; d) M. H. Becker, P. Chua, R. Downham, C. J. Douglas, N. K. Garg, S. Hiebert, S. Jaroch, R. T. Matsuoka, J. A. Middleton, F. W. Ng, L. E. Overman, *J. Am. Chem. Soc.* **2007**, *129*, 11987–12002.
- [9] For preparation of enol ethers by Wittig reactions of ketones, see: a) C. Nevado, D. J. Cárdenas, A. M. Echavarren, *Chem. Eur. J.* **2003**, *9*, 2627–2635; b) M. G. Kulkarni, S. I. Davawala, A. K. Doke, D. S. Pendharkar, *Synthesis* **2004**, 2919–2926; c) P. K. Yong, A. Banerjee, *Org. Lett.* **2005**, *7*, 2485–2487; d) M. E. Bos, C. Loncaric, C. Wu, W. D. Wulff, *Synthesis* **2006**, 3679–3705; e) K. Kobayashi, Y. Shirai, T. Nagaoka, H. Konishi, *Synth. Commun.* **2009**, *39*, 2866–2881.
- [10] a) D. Zhang, J. M. Ready, *Org. Lett.* **2005**, *7*, 5681–5683; b) J. R. DeBergh, K. M. Spivey, J. M. Ready, *J. Am. Chem. Soc.* **2008**, *130*, 7828–7829.
- [11] a) J. F. Normant, M. Bourgain, *Tetrahedron Lett.* **1971**, *12*, 2583–2586; for relevant reviews covering carbocupration of alkynes, see: b) J. F. Normant, A. Alexakis, *Synthesis* **1981**, 841–870; c) B. H. Lipshutz, S. Sengupta, *Org. React.* **1992**, *41*, 135–631.
- [12] a) S. A. Rao, P. Knochel, *J. Am. Chem. Soc.* **1991**, *113*, 5735–5741; b) P. Knochel, R. D. Singer, *Chem. Rev.* **1993**, *93*, 2117–2188.
- [13] a) D. E. Van Horn, E. Negishi, *J. Am. Chem. Soc.* **1978**, *100*, 2252–2254; b) E. Negishi, D. E. Van Horn, T. Yoshida, *J. Am. Chem. Soc.* **1985**, *107*, 6639–6647; c) P. Wipf, S. Lim, *Angew. Chem.* **1993**, *105*, 1095–1097; *Angew. Chem. Int. Ed. Engl.* **1993**, *32*, 1068–1071.
- [14] a) B. Gourdet, H. W. Lam, *J. Am. Chem. Soc.* **2009**, *131*, 3802–3803; b) B. Gourdet, M. E. Rudkin, C. A. Watts, H. W. Lam, *J. Org. Chem.* **2009**, *74*, 7849–7858; c) B. Gourdet, D. L. Smith, H. W. Lam, *Tetrahedron* **2010**, *66*, 6026–6031.
- [15] For recent reviews of enamide chemistry, see: a) D. R. Carbery, *Org. Biomol. Chem.* **2008**, *6*, 3455–3460; b) R. Matsubara, S. Kobayashi, *Acc. Chem. Res.* **2008**, *41*, 292–301.
- [16] For reviews of ynamide chemistry, see: a) K. A. DeKorver, H. Li, A. G. Lohse, R. Hayashi, Z. Lu, Y. Zhang, R. P. Hsung, *Chem. Rev.* **2010**, *110*, 5064–5106; b) G. Evano, A. Coste, K. Jouvin, *Angew. Chem.* **2010**, *122*, 2902–2921; *Angew. Chem. Int. Ed.* **2010**, *49*, 2840–2859; c) A. R. Katritzky, R. Jiang, S. K. Singh, *Heterocycles* **2004**, *63*, 1455–1475; d) J. A. Mulder, K. C. M. Kurtz, R. P. Hsung, *Synlett* **2003**, 1379–1390; e) C. A. Zificsak, J. A. Mulder, R. P. Hsung, C. Rameshkumar, L.-L. Wei, *Tetrahedron* **2001**, *57*, 7575–7606.
- [17] a) C. H. Sugisaki, P. J. Carroll, C. R. D. Correia, *Tetrahedron Lett.* **1998**, *39*, 3413–3416; < lit b > M. Sunose, K. M. Anderson, A. G. Orpen, T. Gallagher, S. J. F. MacDonald, *Tetrahedron Lett.* **1998**, *39*, 8885–8888; c) S. Sukemoto, M. Oshige, M. Sato, K. Mimura, H. Nishioka, H. Abe, T. Harayama, Y. Takeuchi, *Synthesis* **2008**, 3081–3087.
- [18] Isolation of (+)-tanikolide: I. P. Singh, K. E. Milligan, W. H. Gerwick, *J. Nat. Prod.* **1999**, *62*, 1333–1335.
- [19] See the Supporting information for details of the synthesis of enamides **6c**, and **6h–6j**.
- [20] See, for example: a) P. Gupta, P. A. Fernandes, P. Kumar, *Tetrahedron Lett.* **2003**, *44*, 4231–4232; b) S. Masumoto, M. Suzuki, M. Kanai, M. Shibasaki, *Tetrahedron* **2004**, *60*, 10497–10504; c) S. Roy, A. Sharma, N. Chattopadhyay, S. Chattopadhyay, *Tetrahedron Lett.* **2006**, *47*, 7067–7069; d) B. M. Trost, J. Xu, M. Reichle, *J. Am. Chem. Soc.* **2007**, *129*, 282–283.
- [21] For the conversion of (S)-2-cyclohexyl-2-phenylglycolic acid (from oxidation of **4k** with NaClO₂) into (S)-oxybutynin (**7**), see: R. P. Bakale, J. L. Lopez, F. X. McConville, C. P. Vandenbossche, C. H. Senanayake, U.S. Patent 6140529, **1999**.
- [22] a) A. C. Diokno, J. Lapides, *J. Urol.* **1972**, *108*, 307–309; b) I. M. Thompson, R. Lauvetz, *Urology* **1976**, *8*, 452–454.
- [23] These reaction conditions are related to those described previously for copper-catalyzed carbomagnesiation of ynamides, see: a) H. Chechik-Lankin, S. Livshin, I. Marek, *Synlett* **2005**, 2098–2100; b) H.; Yasui, H. Yorimitsu, K. Oshima, *Chem. Lett.* **2007**, *36*, 32–33; c) H. Yasui, H. Yorimitsu, K. Oshima, *Bull. Chem. Soc. Jpn.* **2008**, *81*, 373–379; for stoichiometric carbocupration of ynamides, see: J. J. Prakash Das, H. Chechik, I. Marek, *Nature Chem.* **2009**, *1*, 128–132.
- [24] The asymmetric dihydroxylation of α -cyclohexylstyrene using (DHDO)₂PHAL was also reported in Ref. [20a]. However, the enantioselectivity of this reaction was not reported, and the resulting material required two recrystallizations to obtain **4k** in 92% ee.
- [25] For previous total syntheses of tanikolide, see: a) R. M. Kanada, T. Taniguchi, K. Ogasawa, *Synlett* **2000**, 1019–1021; b) J. Krauss, *Nat. Prod. Lett.* **2001**, *15*, 393–399; c) H. Tanaka, Y. Kozuki, K. Ogasawara, *Tetrahedron Lett.* **2002**, *43*, 4175–4178; d) M.-Y. Chang, C.-L. Lin, S.-T. Chen, *J. Chin. Chem. Soc.* **2001**, *48*, 787–794; e) R. Zhang, Z. Wang, F. Wei, Y. Huang, *Synth. Commun.* **2002**, *32*, 2187–2194; f) H. Mizutani, M. Watanabe, T. Honda, *Tetrahedron* **2002**, *58*, 8929–8936; g) M. Carda, S. Rodríguez, E.

- Castillo, A. Bellido, S. Díaz-Oltra, J. A. Marco, *Tetrahedron* **2003**, *59*, 857–864; h) H. Zhai, Q. Chen, J. Zhao, S. Luo, X. Jia, *Tetrahedron Lett.* **2003**, *44*, 2893–2894; i) A. E. Koumbis, K. M. Dieti, M. G. Vikentiou, J. K. Gallos, *Tetrahedron Lett.* **2003**, *44*, 2513–2516; j) J. M. Schomaker, B. Borhan, *Org. Biomol. Chem.* **2004**, *2*, 621–624; k) H. Arasaki, M. Iwata, M. Makfida, Y. Masaki, *Chem. Pharm. Bull.* **2004**, *52*, 848–852; l) T. Ohgiya, S. Nishiyama, *Tetrahedron Lett.* **2004**, *45*, 8273–8275; m) Q. Chen, H. Deng, J. Zhao, Y. Lu, M. He, H. Zhai, *Tetrahedron* **2005**, *61*, 8390–8393; n) T. Ohgiya, K. Nakamura, S. Nishiyama, *Bull. Chem. Soc. Jpn.* **2005**, *78*, 1549–1554; o) T. Yajima, C. Saito, H. Nagano, *Tetrahedron* **2005**, *61*, 10203–10205; p) Y. Kita, S. Matsuda, E. Fujii, M. Horai, K. Hata, H. Fujioka, *Angew. Chem.* **2005**, *117*, 6007–6010; *Angew. Chem. Int. Ed.* **2005**, *44*, 5857–5860; q) F. Wu, R. Hong, J. Khan, X. Liu, L. Deng, *Angew. Chem.* **2006**, *118*, 4407–4411; *Angew. Chem. Int. Ed.* **2006**, *45*, 4301–4305; r) H. Fujioka, S. Matsuda, M. Horai, E. Fujii, M. Morishita, N. Nishiguchi, K. Hata, Y. Kita, *Chem. Eur. J.* **2007**, *13*, 5238–5248; s) C. Zhang, N. Hosoda, M. Asami, *Tetrahedron: Asymmetry* **2007**, *18*, 2185–2189; t) P. Vichare, A. Chattopadhyay, *Tetrahedron: Asymmetry* **2008**, *19*, 598–602.
- [26] H. Hofmeister, K. Annen, H. Laurent, R. Wiechert, *Angew. Chem.* **1984**, *96*, 720–722; *Angew. Chem. Int. Ed. Engl.* **1984**, *23*, 727–729.
- [27] Y. Zhang, R. P. Hsung, M. R. Tracey, K. C. M. Kurtz, E. L. Vera, *Org. Lett.* **2004**, *6*, 1151–1154.
- [28] (DHQD)₂AQN has been shown to provide superior levels of enantioselection in the dihydroxylation of many alkenes, including those containing only aliphatic substituents, see: H. Becker, K. B. Sharpless, *Angew. Chem.* **1996**, *108*, 447–449; *Angew. Chem. Int. Ed. Engl.* **1996**, *35*, 448–451.
- [29] The *ee* value of diol **12** was determined by HPLC on a chiral stationary phase of the benzoyl ester derivative of the primary alcohol. See the Supporting Information for details.