


Convergent Total Synthesis of (+)-TMC-151C by a Vinylogous Mukaiyama Aldol Reaction and Ring-Closing Metathesis**

Ryosuke Matsui, Kentaro Seto, Yuna Sato, Takahiro Suzuki, Atsuo Nakazaki, and Susumu Kobayashi*

Antibiotic agents of the TMC-151 family were originally isolated from the fungus *Gliocladium catenulatum* Gilman & Abbott TC 1280 by a research group at the pharmaceutical company Tanabe Seiyaku in 1999 as novel polyketides containing β -D-mannoside and D-mannitol groups.^[1] A related TMC-171 family of antibiotics and TMC-154 were also isolated.^[2] Of these compounds, TMC-151C (**1**) shows the most significant cytotoxicity against a wide range of tumor cell lines, including HCT-116, B16, and HeLa cells. TMC-151C displays several interesting structural features. Notably, 1) the polyketide moiety contains three contiguous *anti* homoallylic alcohol motifs in the C1–C13 segment and three methyl-substituted asymmetric carbon centers in the C14–C20 segment, and 2) D-mannose is attached to the C13 hydroxy group as β -D-mannoside, which is still problematic in terms of chemical synthesis (Scheme 1). This combination of significant biological activity and structural complexity encouraged us to attempt the total synthesis of (+)-TMC-151C (**1**).



Scheme 1. Structure of (+)-TMC-151C and the vinylogous Mukaiyama aldol reaction (VMAR). TMC-151C contains three units that should be amenable to synthesis by the VMAR: C2–C5, C6–C9, and C10–C13. TBS = *tert*-butyldimethylsilyl.


[*] R. Matsui, K. Seto, Y. Sato, Dr. T. Suzuki, Dr. A. Nakazaki, Prof. Dr. S. Kobayashi
Faculty of Pharmaceutical Sciences, Tokyo University of Science
2641 Yamazaki, Noda-shi, Chiba 278-8510 (Japan)
Fax: (+81) 4-7121-3671
E-mail: kobayash@rs.noda.tus.ac.jp
Homepage: <http://www.rs.noda.tus.ac.jp/kobalab/index.html>

[**] This research was supported in part by a Grant-in-Aid for Scientific Research (B; KAKENHI No.18390010) from the Japan Society for the Promotion of Science.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201006230>.

We previously developed a highly stereoselective vinylogous Mukaiyama aldol reaction (VMAR) that enabled remarkable remote asymmetric induction by using the vinylketene silyl N,O-acetals **2** and **3** (Scheme 1).^[3] From a synthetic point of view, this method can directly afford the *anti*- δ -hydroxy- α,γ -dimethyl α,β -unsaturated carbonyl unit which is present in many naturally occurring compounds. Indeed, the VMAR has been utilized successfully in natural product syntheses by many research groups,^[4] including our own.^[5] Given the presence of three *anti* homoallylic alcohol motifs (C2–C5, C6–C9, and C10–C13 units) in (+)-**1**, we reasoned that the total synthesis of **1** would provide a representative example of the VMAR. Herein, we report the first total synthesis of (+)-TMC-151C (**1**).


We initially attempted a straightforward linear approach consisting of three iterative VMARs (Scheme 2). The carbohydrate units would then be combined at a later stage in the synthesis. According to the established protocol, the first VMAR of the known chiral aldehyde **4**^[6] with the vinylketene silyl N,O-acetal **3** in the presence of TiCl_4 afforded the



Scheme 2. Attempted linear approach: a) TiCl_4 (1.0 equiv), aldehyde **4** (1.5 equiv), CH_2Cl_2 , $-78 \rightarrow -40^\circ\text{C}$, 91 % (d.r. >20:1); b) TESOTf , 2,6-lutidine, CH_2Cl_2 , 0°C , quantitative; c) LiBH_4 , MeOH , Et_2O , 0°C , 98 %; d) MnO_2 , CH_2Cl_2 , room temperature, 99 %; e) TiCl_4 (4.0 equiv), **3** (5.0 equiv), CH_2Cl_2 , $-78 \rightarrow -20^\circ\text{C}$, 51 % (d.r. 13:1); f) TESOTf , 2,6-lutidine, CH_2Cl_2 , 0°C , 93 %; g) LiBH_4 , MeOH , Et_2O , 0°C , 94 %; h) MnO_2 , CH_2Cl_2 , room temperature, quantitative; i) TiCl_4 (5.0 equiv), **3** (5.0 equiv), CH_2Cl_2 , $-78 \rightarrow -20^\circ\text{C}$, complex mixture. TES = triethylsilyl, Tf = trifluoromethanesulfonyl.


corresponding C12–C13 *anti* aldol adduct **5** in 91 % yield with high diastereoselectivity. TES protection of the secondary alcohol, reductive removal of the chiral auxiliary with LiBH_4 , and oxidation of the resulting primary alcohol with MnO_2 provided the second VMAR substrate **6**. An excess amount of TiCl_4 and the *N,O*-acetal **3** were required for this second VMAR owing to the low reactivity of enal **6**. Under these conditions, the desired aldol adduct **7** was formed in 51 % yield with good diastereoselectivity (d.r. 13:1; the minor diastereomer was separated from **7** by preparative TLC). The aldol adduct **7** was transformed into enal **8** by the same sequence of reactions used for the conversion of **5** into **6**. However, despite repeated efforts, the desired third VMAR product **9** could not be obtained from **8**. In each case, a complex mixture of products was generated, probably as a result of the Lewis acidity of TiCl_4 .

The lack of success with the third VMAR in the linear approach prompted us to investigate an alternative synthetic route. In this context, we recently developed an unusually *E*-selective silicon-tethered ring-closing metathesis (RCM) of silylene acetal **10**, prepared from an allylic alcohol and a homoallylic alcohol (Scheme 3).^[7] Our novel convergent strategy, which is based on this methodology, is outlined



Scheme 3. Formation of an eight-membered ring by *E*-selective silicon-tethered ring-closing metathesis (RCM).


in Scheme 4. (+)-TMC-151C (**1**) was divided at the C6–C7 olefin into two fragments: the left-hand fragment **12** and the right-hand fragment **13**. We envisaged that **12** and **13** could be connected by using this *E*-selective silicon-tethered RCM methodology. We reasoned that the use of this methodology could result in a short and convergent synthetic route. The left-hand fragment **12** was synthesized from the VMAR product **5** by β -mannosylation followed by asymmetric crotylation. To dispense with the protection–deprotection steps, we introduced a β -mannosyl group into **5**. Although β -mannoside formation is often regarded as one of the most difficult glycosylation reactions, several excellent methodologies have been developed to facilitate this transforma-



Scheme 4. Convergent retrosynthetic analysis based on the VMAR and RCM.

tion.^[8] The right-hand fragment **13** was synthesized by a VMAR with methacrolein (**14**) and esterification with a D-mannitol derivative.


The synthesis of the left-hand fragment **12** began with a β -selective mannosylation of **5** (Scheme 5). After extensive studies in which we examined a variety of different approaches, we found that β -mannosylation of **5** with the mannosyl donor **15** by the method developed recently by Crich and Karatholuvhu^[8g] afforded the desired product **16** in 82 % yield almost as a single isomer. The stereostructure of **16** was determined by NOE experiments.^[9] Reductive removal of the chiral auxiliary in **16** with LiBH_4 , followed by the



Scheme 5. Synthesis of the left-hand fragment **12**: a) **15** (Ar = *p*-CF₃C₆H₄), BSP, TiF_4 , TTBP, 4 Å M.S., CH_2Cl_2 , -60°C , 82 % (β : α > 20:1); b) LiBH_4 , MeOH, Et_2O , 0°C , 97 %; c) Na, liquid NH_3 , THF–*t*BuOH, $-70 \rightarrow -20^\circ\text{C}$, 75 %; d) TESOTf, 2,6-lutidine, CH_2Cl_2 , -78°C , quantitative; e) AcOH– H_2O , THF, room temperature, 47 % (89 % based on recovered starting material); f) MnO_2 , CH_2Cl_2 , room temperature, 97 %; g) (*E*)-(R,R)-crotyl boronate, 4 Å M.S., toluene, -78°C , 89 % (d.r. 4.7:1). Bn = benzyl, BSP = 1-benzenesulfinylpiperidine, M.S. = molecular sieves, TTBP = 2,4,6-tri-*tert*-butylpyrimidine.

removal of all protecting groups under Birch conditions,^[8g] provided the pentaol **17**. Following the protection of all five hydroxy groups of **17** as their TES ethers, selective cleavage of the primary allylic TES ether with AcOH–H₂O–THF afforded the corresponding allylic alcohol, which was oxidized with MnO₂ in 97 % yield. A Roush asymmetric *anti* crotylation^[10] then gave the desired left-hand fragment **12** in 89 % yield with d.r. 4.7:1 (the minor diastereomer was separated by silica-gel column chromatography).

Next, we focused on the synthesis of the right-hand fragment **13** (Scheme 6). The synthesis commenced with a VMAR between the N,O-acetal **3** and methacrolein (**14**). Under standard conditions (TiCl₄ in CH₂Cl₂, 2.0 equivalents of **14**), the aldol adduct **18** was formed in low yield (23 %), presumably as a result of polymerization of **14**. During an extensive survey of reaction conditions, we improved the yield of




Scheme 6. Synthesis of the right-hand fragment **13**: a) TiCl₄, methacrolein (**14**; 4.0 equiv), toluene, –78 → –50 °C, 65 % (d.r. > 20:1); b) PMBOC(NH)CCl₃, CSA (cat.), CH₂Cl₂, 0 °C → RT, 68 %; c) LiOH·H₂O, H₂O₂, THF–H₂O, 0 °C, 83 %; d) Piv₂O, Et₃N, CH₂Cl₂, 0 °C; e) **20**, DMAP, toluene, room temperature, 60 % (over 2 steps); f) DDQ, CH₂Cl₂–buffer (pH 7.5), 0 °C, 93 %. PMB = *p*-methoxybenzyl, CSA = camphorsulfonic acid, DDQ = 2,3-dichloro-5,6-dicyano-1,4-benzoquinone, DMAP = *N,N*-4-dimethylaminopyridine, Piv = pivaloyl.

18 to 65 % with greater than 20:1 diastereoselectivity by using toluene as the solvent and 4.0 equivalents of **14**. The major by-products were

1,4-addition adducts (15 % yield) as a mixture of diastereomers. After protection of the secondary alcohol with PMBOC(NH)CCl₃, the chiral auxiliary was removed with LiOH and H₂O₂ to provide the carboxylic acid **19**. Esterification of the unsaturated acid **19** (as the mixed anhydride formed by treatment with Piv₂O) with the alcohol **20**^[9] afforded the corresponding ester in 60 % overall yield for the two steps. Cleavage of the PMB ether gave the right-hand fragment **13**.

With both fragments in hand, we proceeded to the final stage of the total synthesis of (+)-**1** (Scheme 7). Our silicon-tethering method with Et₂NPh₂SiCl/Et₃N/DMAP^[7] was used successfully for the connection of **12** and **13** to afford silylene acetal **21** in 54 % yield (81 % from **12** as calculated on the basis of recovered starting material). The crucial silicon-



Scheme 7. Completion of the total synthesis of (+)-TMC-151C (**1**): a) **13** (1.5 equiv), Et₂NPh₂SiCl, Et₃N, DMAP (cat.), then **12**, DMAP, CH₂Cl₂, 0 °C → RT, 54 % (81 % from **12** on the basis of recovered starting material); b) Hoveyda–Grubbs second-generation catalyst (20 mol %), *p*-benzoquinone, xylene, reflux, 87 % (*E/Z* > 20:1); c) HF–pyridine, pyridine, then aqueous HF, THF–MeCN, 0 °C → RT, 54 %.

tethered RCM^[11,12] of **21** under the previously reported conditions^[7] (with the Hoveyda–Grubbs second-generation catalyst^[13] in the presence of *p*-benzoquinone^[14] in xylene at reflux) provided the desired *E* olefin **22** in 87 % yield with high stereoselectivity (*E/Z* > 20:1). The configuration of the *E* olefin **22** was determined by NOE experiments.^[9] Finally, desilylation by the sequential treatment of **22** with HF–pyridine and aqueous HF completed the total synthesis of (+)-TMC-151C.^[15] The spectroscopic data (¹H NMR, ¹³C NMR, and IR spectra, HRMS) of synthetic (+)-**1** were identical to those of natural (+)-**1**.^[1a]

In conclusion, we have completed the first total synthesis of (+)-TMC-151C by a highly convergent synthetic route. Characteristic features of the present synthesis include the construction of the C1–C5 and C9–C13 units by vinylogous Mukaiyama aldol reactions and the use of a silicon-tethered diene for the stereoselective formation of the C6–C7 double bond by our recently developed *E*-selective RCM reaction to give an eight-membered ring. This synthetic strategy should provide efficient access to a range of related naturally occurring polyketides containing pent-2-ene-1,5-diol units.

Received: October 5, 2010

Published online: December 22, 2010

Keywords: aldol reaction · natural products · polyketides · ring-closing metathesis · total synthesis

- [1] a) J. Kohno, M. Nishio, M. Sakurai, K. Kawano, H. Hiramatsu, N. Kameda, N. Kishi, T. Yamashita, T. Okuda, S. Komatsubara, *Tetrahedron* **1999**, *55*, 7771–7786; b) J. Kohno, M. Nishio, N. Kishi, T. Okuda, S. Komatsubara, *J. Antibiot.* **2000**, *53*, 1301–1304; c) T. Okuda, J. Kohno, N. Kishi, Y. Asai, M. Nishio, S. Komatsubara, *Mycoscience* **2000**, *41*, 239–253.
[2] J. Kohno, Y. Asai, M. Nishio, M. Sakurai, K. Kawano, H. Hiramatsu, N. Kameda, N. Kishi, T. Okuda, S. Komatsubara, *J. Antibiot.* **1999**, *52*, 1114–1123.

- [3] a) S.-i. Shirokawa, M. Kamiyama, T. Nakamura, M. Okada, A. Nakazaki, S. Hosokawa, S. Kobayashi, *J. Am. Chem. Soc.* **2004**, *126*, 13604–13605; b) M. Shinoyama, S.-i. Shirokawa, A. Nakazaki, S. Kobayashi, *Org. Lett.* **2009**, *11*, 1277–1280; c) M. Yamaoka, A. Nakazaki, S. Kobayashi, *Tetrahedron Lett.* **2010**, *51*, 287–289.
- [4] a) S. Hosokawa, T. Ogura, H. Togashi, K. Tatsuta, *Tetrahedron Lett.* **2005**, *46*, 333–337; b) S. Hosokawa, K. Yokota, K. Imamura, Y. Suzuki, M. Kawarasaki, K. Tatsuta, *Tetrahedron Lett.* **2006**, *47*, 5415–5418; c) S. Hosokawa, S. Kuroda, K. Imamura, K. Tatsuta, *Tetrahedron Lett.* **2006**, *47*, 6183–6186; d) X. Jiang, B. Liu, S. Lebreton, J. K. De Brabander, *J. Am. Chem. Soc.* **2007**, *129*, 6386–6387; e) K. C. Nicolaou, R. Guduru, Y.-P. Sun, B. Banerji, D. Y.-K. Chen, *Angew. Chem.* **2007**, *119*, 6000–6004; *Angew. Chem. Int. Ed.* **2007**, *46*, 5896–5900; f) K. C. Nicolaou, Y.-P. Sun, R. Guduru, B. Banerji, D. Y.-K. Chen, *J. Am. Chem. Soc.* **2008**, *130*, 3633–3644; g) A. Schmauder, S. Müller, M. E. Maier, *Tetrahedron* **2008**, *64*, 6263–6269; h) S. Hosokawa, K. Yokota, K. Imamura, Y. Suzuki, M. Kawarasaki, K. Tatsuta, *Asian J. Chem.* **2008**, *3*, 1415–1421; i) B. H. Lipshutz, B. Amorelli, *J. Am. Chem. Soc.* **2009**, *131*, 1396–1397; j) L. Wang, J. Gong, L. Deng, Z. Xiang, Z. Chen, Y. Wang, J. Chen, Z. Yang, *Org. Lett.* **2009**, *11*, 1809–1812; k) S. Hosokawa, Y. Mukaeda, R. Kawahara, K. Tatsuta, *Tetrahedron Lett.* **2009**, *50*, 6701–6704; l) S. Hosokawa, K. Matsushita, S. Tokimatsu, T. Toriumi, Y. Suzuki, K. Tatsuta, *Tetrahedron Lett.* **2010**, *51*, 5532–5536; for reviews, see: m) K. Tatsuta, S. Hosokawa, *Chem. Rev.* **2005**, *105*, 4707–4729; n) K. Tatsuta, S. Hosokawa, *Chem. Rec.* **2006**, *6*, 217–233; o) S. Hosokawa, K. Tatsuta, *Mini-Rev. Org. Chem.* **2008**, *5*, 1–18.
- [5] a) T. Nakamura, S.-i. Shirokawa, S. Hosokawa, A. Nakazaki, S. Kobayashi, *Org. Lett.* **2006**, *8*, 677–679; b) S.-i. Shirokawa, M. Shinoyama, I. Ooi, S. Hosokawa, A. Nakazaki, S. Kobayashi, *Org. Lett.* **2007**, *9*, 849–852; c) M. Yamaoka, Y. Fukatsu, A. Nakazaki, S. Kobayashi, *Tetrahedron Lett.* **2009**, *50*, 3849–3852; d) M. Yamaoka, A. Nakazaki, S. Kobayashi, *Tetrahedron Lett.* **2009**, *50*, 6764–6768.
- [6] E.-i. Negishi, Z. Tan, B. Liang, T. Novak, *Proc. Natl. Acad. Sci. USA* **2004**, *101*, 5782–5787.
- [7] R. Matsui, K. Seto, K. Fujita, T. Suzuki, A. Nakazaki, S. Kobayashi, *Angew. Chem.* **2010**, DOI: 10.1002/ange.201004746; *Angew. Chem. Int. Ed.* **2010**, DOI: 10.1002/anie.201004746.
- [8] For examples of β -mannosylation, see: a) Y. Ito, T. Ogawa, *Angew. Chem.* **1994**, *106*, 1843–1845; *Angew. Chem. Int. Ed. Engl.* **1994**, *33*, 1765–1767; b) Y. Ito, H. Ando, M. Wada, T. Kawai, Y. Ohnishi, Y. Nakahara, *Tetrahedron* **2001**, *57*, 4123–4132; c) A. A.-H. Abdel-Rahman, S. Jonke, E. S. H. El Ashry, R. R. Schmidt, *Angew. Chem.* **2002**, *114*, 3100–3103; *Angew. Chem. Int. Ed.* **2002**, *41*, 2972–2974; d) S.-i. Tanaka, M. Takashina, H. Tokimoto, Y. Fujimoto, K. Tanaka, K. Fukase, *Synlett* **2005**, 2325–2328; e) D. Crich, P. Jayalath, *Org. Lett.* **2005**, *7*, 2277–2280; f) D. Crich, P. Jayalath, T. K. Hutton, *J. Org. Chem.* **2006**, *71*, 3064–3070; g) D. Crich, M. S. Karatholuvhu, *J. Org. Chem.* **2008**, *73*, 5173–5176; for a review, see: h) J. J. Gridley, H. M. I. Osborn, *J. Chem. Soc. Perkin Trans. 1* **2000**, 1471–1491.
- [9] See the Supporting Information for details.
- [10] a) W. R. Roush, K. Ando, D. B. Powers, A. D. Palkowitz, R. L. Halterman, *J. Am. Chem. Soc.* **1990**, *112*, 6339–6348; b) W. R. Roush, A. D. Palkowitz, K. Ando, *J. Am. Chem. Soc.* **1990**, *112*, 6348–6359.
- [11] For reviews on olefin metathesis, see: a) A. K. Chatterjee, T.-L. Choi, D. P. Sanders, R. H. Grubbs, *J. Am. Chem. Soc.* **2003**, *125*, 11360–11370; b) S. J. Connon, S. Blechert, *Angew. Chem.* **2003**, *115*, 1944–1968; *Angew. Chem. Int. Ed.* **2003**, *42*, 1900–1923; c) A. Gradillas, J. Pérez-Castells, *Angew. Chem.* **2006**, *118*, 6232–6247; *Angew. Chem. Int. Ed.* **2006**, *45*, 6086–6101; d) J. Cossy, S. Arseniyadis, C. Meyer, R. H. Grubbs, *Metathesis in Natural Product Synthesis: Strategies, Substrates and Catalysts*, Wiley-VCH, Weinheim, **2010**.
- [12] For previous examples of silicon-tethered ring-closing metathesis, see: a) P. A. Evans, V. S. Murthy, *J. Org. Chem.* **1998**, *63*, 6768–6769; b) T. R. Hoye, M. A. Promo, *Tetrahedron Lett.* **1999**, *40*, 1429–1432; c) A. Briot, M. Bujard, V. Gouverneur, S. P. Nolan, C. Mioskowski, *Org. Lett.* **2000**, *2*, 1517–1519; d) T. M. Gierasch, M. Chytil, M. T. Didiuk, J. Y. Park, J. J. Urban, S. P. Nolan, G. L. Verdine, *Org. Lett.* **2000**, *2*, 3999–4002; e) B. A. Harrison, G. L. Verdine, *Org. Lett.* **2001**, *2*, 2157–2159; f) P. Van de Weghe, D. Aoun, J.-G. Boiteau, J. Eustache, *Org. Lett.* **2002**, *4*, 4105–4108; g) P. A. Evans, J. Cui, G. P. Buffone, *Angew. Chem.* **2003**, *115*, 1776–1779; *Angew. Chem. Int. Ed.* **2003**, *42*, 1734–1737; h) E. M. Casey, P. Teesdale-Spittle, J. E. Harvey, *Tetrahedron Lett.* **2008**, *49*, 7021–7023; i) T. R. Hoye, J. Jeon, L. C. Kopel, T. D. Ryba, M. A. Tennakoon, Y. Wang, *Angew. Chem.* **2010**, *122*, 6287–6291; *Angew. Chem. Int. Ed.* **2010**, *49*, 6151–6155; for reviews on the use of silicon tethers, see: j) M. Bols, *Chem. Rev.* **1995**, *95*, 1253–1277; k) I. Fleming, A. Barbero, D. Walter, *Chem. Rev.* **1997**, *97*, 2063–2192; l) D. R. Gauthier, Jr., K. S. Zandi, K. J. Shea, *Tetrahedron* **1998**, *54*, 2289–2338; for more-recent examples of useful silicon tethering, see: m) M. Petit, G. Chouraqui, C. Aubert, M. Malacria, *Org. Lett.* **2003**, *5*, 2037–2040; n) J. B. Grimm, D. Lee, *J. Org. Chem.* **2004**, *69*, 8967–8970; o) J. Beignet, P. J. Jervis, L. R. Cox, *J. Org. Chem.* **2008**, *73*, 5462–5475.
- [13] a) S. B. Garber, J. S. Kingsbury, B. L. Gray, A. H. Hoveyda, *J. Am. Chem. Soc.* **2000**, *122*, 8168–8179; b) S. Gessler, S. Randl, S. Blechert, *Tetrahedron Lett.* **2000**, *41*, 9973–9976.
- [14] Isomerization of the terminal alkenes occurred to some extent during RCM without *p*-benzoquinone; for the effect of *p*-benzoquinone, see also: a) S. H. Hong, D. P. Sanders, C. W. Lee, R. H. Grubbs, *J. Am. Chem. Soc.* **2005**, *127*, 17160–17161; b) I. Larrosa, M. I. Da Silva, P. M. Gómez, P. Hannen, E. Ko, S. R. Lenger, S. R. Linke, A. J. P. White, D. Wilton, A. G. M. Barrett, *J. Am. Chem. Soc.* **2006**, *128*, 14042–14043.
- [15] Although the $[\alpha]_D$ value of synthetic (+)-**1** ($+4.4 \text{ deg cm}^3 \text{ g}^{-1} \text{ dm}^{-1}$, $c = 1.3 \times 10^{-3} \text{ g cm}^{-3}$ in MeOH) was lower than that of natural (+)-**1** ($+27 \text{ deg cm}^3 \text{ g}^{-1} \text{ dm}^{-1}$, $c = 1.0 \times 10^{-3} \text{ g cm}^{-3}$ in MeOH), the stereostructure and the purity of synthetic (+)-**1** were confirmed by NMR spectroscopy.