

Stable Iso-osmabenzenes from a Formal [3+3] Cycloaddition Reaction of Metal Vinylidene with Alkynols**

Qianyi Zhao, Lei Gong, Chunfa Xu, Jun Zhu,* Xumin He, and Haiping Xia*

The investigation on the synthesis and reactivity of metal vinylidene complexes has always been an active area,^[1] especially given their applications as well-known catalysts for metathesis and carbon–carbon and carbon–heteroatom coupling reactions.^[1b,e,2] Compared with other reactions of metal vinylidenes, the cyclization reactions are underdeveloped.


Previous studies showed that the vinylidene $M=C_{\alpha}$ bond participated in cycloadditions (see *a*) as key steps in carbon–carbon coupling reactions,^[3] and some cycloaddition products have also been isolated.^[4]

Interestingly, similar reactions of the metal vinylidene $C_{\alpha}=C_{\beta}$ bond (see *b*) have been realized as well.^[5] However, cyclizations of the skeleton of metal vinylidene ($M=C_{\alpha}=C_{\beta}$) as a unit (see *c*) remain scarce. In fact, the reported cyclization is a formal [3+2] cycloaddition reaction.^[5c] To the best of our knowledge, the [3+3] cyclization reactions of metal vinylidene to construct six-membered rings have not been reported.

Moreover, since the first isolation of metallabenzene was achieved by Roper and co-workers,^[7a] considerable research interest has been attracted in this field.^[6,7] Nevertheless, the metallabenzene analogue isometallabenzene, which can be seen as a metal-containing 1,2,4-cyclohexatriene, has not been thoroughly explored. Until now, only the 16e[−] isometallabenzene $[Os\{C=C(Ph)-CH(Ph)-CH=C(CH_2Ph)\}Cl-(P\text{Pr}_3)_2]$, prepared by double coupling reactions, was reported by Esteruelas and co-workers in 2004.^[8] Herein, we report an unprecedented formal [3+3] cycloaddition reaction between the hydride vinylidene complex $[OsHCl_2(=C=CH(PPh_3))(PPh_3)_2]$ (**2**) and alkynols under mild conditions to give stable 18e[−] iso-osmabenzenes. The origin of the unexpected stability of these iso-osmabenzene complexes and their isomerization into η^5 -cyclopentadienyl complexes through metalated cyclopentadiene intermediates is also described.


Stirring of the osmium hydride alkenylcarbyne complex $[OsHCl_2(=C-C(PPh_3)=CHPh)(PPh_3)_2]BF_4$ (**1**)^[9] in a mixture

of H_2O/CH_3OH (3:2 v/v) at reflux produced the osmium hydride vinylidene complex **2**, which was isolated as a light-yellow solid in 60% yield (Scheme 1). The other product, PhCHO, was detected by ¹H NMR (δ = 9.3 ppm; PhCHO) and GC analysis of the reaction mixture (see Figure S1 in the Supporting Information). Complex **2** was characterized by NMR spectroscopy and elemental analysis, and the structure was additionally confirmed by single-crystal X-ray diffraction (Figure S2).



Scheme 1. Preparation of osmium hydride vinylidene **2**.

Complex **2** is thermally stable in the solid state but highly reactive in solution. Treatment of **2** with $HC\equiv CCH(OH)Ph$ in dichloromethane or chloroform at room temperature led to a fast color change from light yellow to brown red. The unexpected iso-osmabenzene **3** was generated from a formal [3+3] cycloaddition and isolated in high yield (93%). Similarly, when **2** reacted with $HC\equiv CCH(OH)CH=CH_2$ or $HC\equiv CCH(OH)CH_2CH_3$ under the same reaction conditions, the other two iso-osmabenzenes, **4** (88% yield) and **5** (80% yield), were also formed (Scheme 2).



Scheme 2. Preparation of iso-osmabenzene **3–5**.

The structure of **3** was characterized by single-crystal X-ray diffraction analysis.^[10] As shown in Figure 1, the six-membered ring is almost planar. The deviations, in Å, from the best plane are 0.0410 (Os1), 0.0003 (C1), 0.0515 (C2), 0.0571 (C3), 0.0022 (C4), and 0.0491 (C5). The Os1=C1 bond

[*] Q. Zhao, L. Gong, C. Xu, Prof. Dr. J. Zhu, Prof. Dr. X. He, Prof. Dr. H. Xia
State Key Laboratory of Physical Chemistry of Solid Surfaces
College of Chemistry and Chemical Engineering
Xiamen University, Xiamen, 361005 (P.R. China)
Fax: (+86) 592-218-6628
E-mail: jun.zhu@xmu.edu.cn
hpxia@xmu.edu.cn

[**] This work was financially supported by the National Science Foundation of China (Nos. 20872123 and 20925208).

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/ange.201006442>.




Figure 1. X-ray crystal structure of **3** (ellipsoids shown at the 50% probability level). Some of the hydrogen atoms are omitted for clarity. Selected bond lengths [Å] and angles [deg]: Os1–Cl1 2.487(2), Os1–Cl2 2.492(3), Os1–P1 2.393(3), Os1–P2 2.398(3), Os–C1 1.781(11), Os1–C5 2.043(12), C1–C2 1.339(13), C2–C3 1.563(14), C3–C4 1.507(13), C4–C5 1.351(14); C1–Os1–C5 78.3(4), Os1–C1–C2 155.1(8), C1–C2–C3 114.2(9), C2–C3–C4 111.6(9), C3–C4–C5 128.6(10), C4–C5–Os1 131.4(8).

length (1.781(11) Å) is at the low-end of the typical Os=C=CRR' (1.78–1.90 Å)^[11] and Os=C_{carbene} (1.78–2.14 Å) bonds.^[12] The Os1–C5 bond length (2.043(12) Å) is within the range observed for typical Os–C_{vinyl} bonds.^[13] The C1=C2, C4=C5, C2–C3, and C3–C4 bond lengths are typical for carbon–carbon double and single bonds. The Os–C and C–C bond lengths within the six-membered ring indicate that the metallacycle has a localized nature. The Os1–C1–C2 angle of 155.1(8)° is comparable to that of Esteruelas' iso-osmabenzene (158.5(3)°)^[8] and Jia's osmabenzynes (148.7(3)–154.9(9)°)^[14] or osmanaphalynes (151.1(5)° and 155.0(3)°).^[15]


In keeping with the solid-state structure of **3**, the ¹H NMR analysis showed no signal for the carbene proton. Two proton signals for the Os–CH=CH unit were observed at δ = 8.3 and 4.7 ppm, respectively, with a coupling constant of 9.5 Hz, which indicates a *cis* geometry.

Mainly attributed to the chirality of the sp³-carbon atom C3, the ³¹P{¹H} NMR spectrum of **3** displayed the same characteristic AB spin system as did Esteruelas' isometallabenzene;^[8] the resonances were centered at δ = –3.4 (d, ²J(PP) = 360.5 Hz) and –7.9 ppm (d, ²J(PP) = 360.5 Hz) for each of the two OsPPh₃, respectively. The signal at δ = 3.9(s) ppm was assigned to CPh₃. Unfortunately, the poor solubility of **3** and **4** prevented ¹³C{¹H} NMR characterization. Isoosmabenzene **4** and **5** have similar NMR spectroscopic characteristics as **3**.

A plausible mechanism for the formation of **3**, **4**, and **5** is shown in Scheme 2. The insertion of terminal alkynes into Os–H bond should afford intermediate **A**. The subsequent dehydration and C–C coupling between the C_β of the vinylidene fragment and the hydroxy-linked carbon atom in **A** led to the formal [3+3] cycloaddition iso-osmabenzene products. It should be mentioned that the typical reaction of

metal hydride complexes is the insertion of terminal alkynes into their M–H bonds.^[16]

Solid samples of **3**, **4**, and **5** show air and moisture stabilities; they can be heated in air at 100°C for 5 hours without notable decomposition. In solution, complex **3** is thermally stable to some extent, and tolerant to bases such as Na₂CO₃ and NaOH; however **3** is sensitive to acids such as HCl and HBF₄. In contrast to **3**, complexes **4** and **5** are less stable. When **4** was continuously stirred in chloroform at room temperature, it rearranged slowly into the η⁵-cyclopentadienyl complex **7** after three days (Scheme 3). The




Scheme 3. Preparation of **6–8**.

isomerization of **5**, which led to **8**, was faster and accomplished in quantitative yield within two days. The solution of **3** behaved similarly to give **6**, but very slowly as the transformation took more than one week. By comparing the above-mentioned results, the stability of **3**, **4**, and **5** follows the sequence **3** > **4** > **5**, which is mainly influenced by the substituents on the sp³-carbon atom.

DFT calculations have been carried out to gain insight into the relative stabilities of **3**, **4**, and **5**. Compounds **3'**, **4'**, and **5'** are model complexes in which the PPh₃ ligand is replaced with PH₃. The computed relative free energies for **3'**, **4'**, and **5'** with respect to the model η⁵-cyclopentadienyl complexes **6'**, **7'**, and **8'** are 11.7, 13.5, and 19.6 kcal mol^{–1} (Scheme 4a), respectively, and are consistent with the experimental observations. For comparison, we also calculated the energy difference for the first iso-osmabenzene [Os{C=C(Ph)–CH(Ph)–CH=C(CH₂Ph)}Cl(PiPr₃)₂]^[8] (model complex **Os1'**) relative to the η⁵-Cp complex **Os1(Cp')** (38.2 kcal mol^{–1}, Scheme 4b).

To probe the origin of the relatively high stability of **3'**, we additionally constructed three more model complexes, namely **3-PH₃⁺**, **3-Cl[–]**, and **3-PH₃[–]**, to examine the role of the phosphonium substituent and different electron counts (Scheme 4c). Interestingly, when the phosphonium substituent is replaced by a hydrogen atom, the relative energy of **3-PH₃⁺** is increased from 11.7 kcal mol^{–1} to 26.3 kcal mol^{–1} compared to **3'**, indicating that the phosphonium substituent plays a crucial role in stabilizing iso-osmabenzene complexes. Moreover, when one Cl[–] or PH₃ ligand is deliberately removed from **3'** to form a 16e[–] iso-osmabenzene **3-Cl[–]** or



Scheme 4. Relative free energies (kcal mol⁻¹, in the solvent of chloroform) of iso-osmabenzene compared with Cp complexes.


3-PH₃', respectively, their stabilities relative to the η^5 -complexes are decreased to 34.5 and 25.4 kcal mol⁻¹, respectively, suggesting that the contribution to the stability from the 18e⁻ rule is not negligible. Taken together, both the phosphonium substituent at the C _{β} position and the 18e⁻ nature of **3** contribute to the high stability.

Additional studies showed that complexes **6**, **7**, and **8** could also be produced by heating **3**, **4**, and **5**, respectively, in chloroform for hours. By changing the counteranion of **8** to BPh₄⁻, complex **8a** was prepared and confirmed by X-ray diffraction (Figure S3). Furthermore, the complexes **6**, **7**, **8**, and **8a** were characterized by NMR spectroscopy and elemental analysis.

It is well known that metallabenzene can undergo carbene migratory insertion reactions into cyclopentadienyl complexes.^[17] A similar process can be detected in the isomerization of isometallabenzene (Scheme 3). The proposed mechanism shows that it may go through the metalated cyclopentadiene intermediate **B**. As a result of its instability, the 16e⁻ five-coordinated **B** immediately undergoes H-shift onto the C _{α} atom of the Os-C_{vinyl} followed by Os-C bond cleavage, giving the stable η^5 -cyclopentadienyl complexes **6**, **7**, and **8**. Indeed, the computed data for the metalated cyclopentadiene **B6'**, **B7'**, and **B8'** model complexes are less stable by 8.7, 10.0, and 14.0 kcal mol⁻¹ relative to the corresponding

η^5 -cyclopentadienyl model complexes **6'**, **7'**, and **8'**, respectively (Scheme 4a).

To capture the intermediate in this reaction, CO was introduced to stabilize **B**. A flask containing a solution of **3** was placed in an ice bath and under a CO atmosphere, and the mixture was stirred, thus leading to **9** (Scheme 5). Complex **9** was characterized by NMR spectroscopy, elemental analysis, and single-crystal X-ray diffraction (Figure 2).^[10] The forma-



Scheme 5. Preparation of **9**.





Figure 2. X-ray crystal structure of **9** (ellipsoids shown at the 50% probability level).

tion of **9** suggests that iso-osmabenzene rearranged via metalated cyclopentadiene intermediates.

At first glance, isometallabenzene **D** is similar to metallabenzene **C** and metallabenzynes **E**, particularly the resonance structure **E'** (Scheme 6). Compared with metallabenzynes **E**, metallabenzene **C** and isometallabenzene **D** have a lower degree of unsaturation. Although metallabenzene and metallabenzynes are familiar to us, isometallabenzene is extremely rare. In this regard, isolation of the isometallabenzene **3**, **4**, and **5** will definitely enrich the chemistry of six-membered metallacycles.



Scheme 6. Metallabenzene, isometallabenzene, and metallabenzynes.

In summary, isometallabenzene was isolated from the novel [3+3] cycloaddition reactions of a metal vinylidene complex with alkynols in high yields at room temperature.

This method provides a simple and efficient route to prepare stable isometallabenzenes. The origin of the unexpected stability was probed by using DFT calculations, which suggest that both the phosphonium substituent at the C_β position and the 18e[−] nature of the complex are the stabilizing factors. We also found that these isometallabenzenes could isomerize into η⁵-cyclopentadienyl complexes via a metalated cyclopentadiene intermediate. The new reaction pattern between metal vinylidene and alkynols represents a promising synthetic method to construct other metallacycles. Expansion of cyclizations of vinylidene complexes to heteroatom-containing metallacycles is ongoing.

Experimental Section

2: The osmium hydride-alkenylcarbyne complex [OsHCl₂(=C=C(PPh₃)=CHPh)(PPh₃)₂][BF₄] (**1**) (506 mg, 0.405 mmol) was stirred in a mixture of CH₃OH (20 mL) and H₂O (30 mL) under reflux for about 12 h to give a light-yellow suspension, which was collected by filtration, washed with methanol and diethyl ether, and then dried under vacuum. Yield: 261 mg, 60%. ¹H NMR (400.1 MHz, CDCl₃): δ = −8.0 (td, J(P,H) = 15.6 Hz, J(P,H) = 4.0 Hz, 1H, OsH), 7.7–6.6 ppm (m, 45H, PPh₃), 0.6 ppm (br, 1H, OsCCHPPh₃); ³¹P{¹H} NMR (162.0 MHz, CDCl₃): δ = 7.6 (d, J(P,P) = 4.9 Hz, OsPPh₃), −1.06 ppm (t, J(P,P) = 4.9 Hz, CPhPPh₃); elemental analysis (%) calcd for C₅₆H₄₇P₃Cl₂Os: C 62.62, H 4.41; found: C 62.32, H 4.31.

3: HC≡CCH(OH)Ph (55.4 μL, 0.447 mmol) was added to a solution of **2** (400 mg, 0.372 mmol) in CH₂Cl₂ or CHCl₃ (15 mL). The reaction mixture was stirred at room temperature for about 1 h at which point a brown-red solution was obtained. A red solid was collected after the solvent was evaporated to dryness under vacuum and the resulting residue was washed by diethyl ether and then dried under vacuum. Yield: 412 mg, 93%. ¹H NMR (500.2 MHz, CDCl₃): δ = 8.3 (dd, J(H,H) = 9.5 Hz, J(H,H) = 2.0 Hz, 1H, OsCHCHCH(Ph)), 7.5–6.6 (m, 50H, Ph), 5.2 (m, 1H, OsCHCHCH(Ph)), 4.7 ppm (m, OsCHCHCH(Ph)); ³¹P{¹H} NMR (202.5 MHz, CDCl₃): δ = 3.9 (s, CPhPPh₃), −3.4 (d, J(P,P) = 360.5 Hz, OsPPh₃), −7.9 ppm (d, J(P,P) = 360.5 Hz, OsPPh₃); elemental analysis (%) calcd for C₆₅H₅₃Cl₂P₃Os: C 65.71, H 4.50; found: C 65.70, H 4.33.

For details of the preparation of **4**, **5**, **6**, **7**, **8**, **8a**, and **9**, see the Supporting Information.

Received: October 14, 2010

Published online: January 5, 2011

Keywords: cycloaddition · density functional calculations · metallacycles · osmium · vinylidenes

- [1] For reviews, see: a) M. I. Bruce, *Chem. Rev.* **1991**, *91*, 197; b) M. C. Puerta, P. Valerga, *Coord. Chem. Rev.* **1999**, *193–195*, 977; c) M. A. Esteruelas, A. M. López, M. Oliván, *Coord. Chem. Rev.* **2007**, *251*, 795; d) C.-M. Che, C.-M. Ho, J.-S. Huang, *Coord. Chem. Rev.* **2007**, *251*, 2145; e) P. Dixneuf, C. Bruneau, *Metal Vinylidenes and Allenylidenes in Catalysis: From Reactivity to Applications in Synthesis*, Wiley-VCH, Weinheim, **2008**.
- [2] For reviews, see: a) C. Bruneau, P. H. Dixneuf, *Acc. Chem. Res.* **1999**, *32*, 311; b) H. Katayama, F. Ozawa, *Coord. Chem. Rev.* **2004**, *248*, 1703; c) H. Werner, *Coord. Chem. Rev.* **2004**, *248*, 1693; d) C. Bruneau, P. H. Dixneuf, *Angew. Chem.* **2006**, *118*, 2232; *Angew. Chem. Int. Ed.* **2006**, *45*, 2176.
- [3] For examples: a) C. Slugovc, K. Mereiter, R. Schmid, K. Kirchner, *J. Am. Chem. Soc.* **1998**, *120*, 6175; b) C. Slugovc, K.

- Mereiter, R. Schmid, K. Kirchner, *Organometallics* **1999**, *18*, 1011; c) M. Murakami, S. Hori, *J. Am. Chem. Soc.* **2003**, *125*, 4720; d) Y. J. Park, B.-I. Kwon, J.-A. Ahn, H. Lee, C.-H. Jun, *J. Am. Chem. Soc.* **2004**, *126*, 13892.
- [4] a) R. Beckhaus, I. Strauß, T. Wagner, P. Kiprof, *Angew. Chem.* **1993**, *105*, 281; *Angew. Chem. Int. Ed. Engl.* **1993**, *32*, 264; b) R. Beckhaus, I. Strauß, T. Wagner, *Angew. Chem.* **1995**, *107*, 738; *Angew. Chem. Int. Ed. Engl.* **1995**, *34*, 688; c) R. Beckhaus, J. Sang, T. Wagner, B. Ganter, *Organometallics* **1996**, *15*, 1176; d) M. Yamaguchi, Y. Arikawa, Y. Nishimura, K. Umakoshi, M. Onishi, *Chem. Commun.* **2009**, 2911.
- [5] For examples: a) P. Alvarez, E. Lastra, J. Gimeno, M. Bassetti, L. R. Falvello, *J. Am. Chem. Soc.* **2003**, *125*, 2386; b) P. Braña, J. Gimeno, J. A. Sordo, *J. Org. Chem.* **2004**, *69*, 2544; c) J. Ipaktschi, J. Mohseni-Ala, A. Dülmer, S. Steffens, C. Wittenburg, J. Heck, *Organometallics* **2004**, *23*, 4902; d) Y.-S. Yen, Y.-C. Lin, S.-L. Huang, Y.-H. Liu, H.-L. Sung, Y. Wang, *J. Am. Chem. Soc.* **2005**, *127*, 18037.
- [6] For reviews, see: a) G. Jia, *Acc. Chem. Res.* **2004**, *37*, 479; b) C. W. Landorf, M. M. Haley, *Angew. Chem.* **2006**, *118*, 4018; *Angew. Chem. Int. Ed.* **2006**, *45*, 3914; c) L. J. Wright, *Dalton Trans.* **2006**, 1821; d) J. R. Bleeke, *Acc. Chem. Res.* **2007**, *40*, 1035.
- [7] For examples: a) G. P. Elliott, W. R. Roper, J. M. Waters, *J. Chem. Soc. Chem. Commun.* **1982**, 811; b) U. Englert, F. Podewils, I. Schiffrs, A. Salzer, *Angew. Chem.* **1998**, *110*, 2196; *Angew. Chem. Int. Ed.* **1998**, *37*, 2134; c) C. E. F. Rickard, W. R. Roper, S. D. Woodgate, L. J. Wright, *Angew. Chem.* **2000**, *112*, 766; *Angew. Chem. Int. Ed.* **2000**, *39*, 750; d) U. Effertz, U. Englert, F. Podewils, A. Salzer, T. Wagner, M. Kaupp, *Organometallics* **2003**, *22*, 264; e) E. Álvarez, M. Paneque, M. L. Poveda, N. Rendón, *Angew. Chem.* **2006**, *118*, 488; *Angew. Chem. Int. Ed.* **2006**, *45*, 474; f) G. R. Clark, P. M. Johns, W. R. Roper, L. J. Wright, *Organometallics* **2008**, *27*, 451; g) A. F. Dalebrook, L. J. Wright, *Organometallics* **2009**, *28*, 5536; h) K. C. Poon, L. Liu, T. Guo, J. Li, H. H. Y. Sung, I. D. Williams, Z. Lin, G. Jia, *Angew. Chem.* **2010**, *122*, 2819; *Angew. Chem. Int. Ed.* **2010**, *49*, 2759; i) G. R. Clark, L. A. Ferguson, A. E. McIntosh, T. Söhnel, L. J. Wright, *J. Am. Chem. Soc.* **2010**, *132*, 13443.
- [8] P. Barrio, M. A. Esteruelas, E. Oñate, *J. Am. Chem. Soc.* **2004**, *126*, 1946.
- [9] B. Liu, H. Wang, H. Xie, B. Zeng, J. Chen, J. Tao, T. B. Wen, Z. Cao, H. Xia, *Angew. Chem.* **2009**, *121*, 5538; *Angew. Chem. Int. Ed.* **2009**, *48*, 5430.
- [10] CCDC 796181 (**3**) and CCDC 796183 (**9**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif. See the Supporting Information for details.
- [11] a) T. B. Wen, W. Y. Hung, Z. Y. Zhou, M. F. Lo, I. D. Williams, G. Jia, *Eur. J. Inorg. Chem.* **2004**, 2837; b) R. Castarlenas, M. A. Esteruelas, R. Lalrempuia, M. Oliván, E. Oñate, *Organometallics* **2008**, *27*, 795.
- [12] a) R. Castarlenas, M. A. Esteruelas, E. Oñate, *Organometallics* **2005**, *24*, 4343; b) T. Bolaño, R. Castarlenas, M. A. Esteruelas, E. Oñate, *Organometallics* **2007**, *26*, 2037; c) R. Castarlenas, M. A. Esteruelas, E. Oñate, *Organometallics* **2007**, *26*, 2129.
- [13] a) M. A. Esteruelas, F. J. Fernández-Alvarez, M. Oliván, E. Oñate, *J. Am. Chem. Soc.* **2006**, *128*, 4596; b) L. Zhang, L. Dang, T. B. Wen, H. H.-Y. Sung, I. D. Williams, Z. Lin, G. Jia, *Organometallics* **2007**, *26*, 2849.
- [14] a) T. B. Wen, Z. Y. Zhou, G. Jia, *Angew. Chem.* **2001**, *113*, 2005; *Angew. Chem. Int. Ed.* **2001**, *40*, 1951; b) T. B. Wen, S. M. Ng, W. Y. Hung, Z. Y. Zhou, M. F. Lo, L.-Y. Shek, I. D. Williams, Z. Lin, G. Jia, *J. Am. Chem. Soc.* **2003**, *125*, 884; c) T. B. Wen, W. Y. Hung, H. H. Y. Sung, I. D. Williams, G. Jia, *J. Am. Chem. Soc.* **2005**, *127*, 2856; d) W. Y. Hung, J. Zhu, T. B. Wen, K. P. Yu,

- H. H. Y. Sung, I. D. Williams, Z. Lin, G. Jia, *J. Am. Chem. Soc.* **2006**, *128*, 13742.
- [15] a) G. He, J. Zhu, W. Y. Hung, T. B. Wen, H. H.-Y. Sung, I. D. Williams, Z. Lin, G. Jia, *Angew. Chem.* **2007**, *119*, 9223; *Angew. Chem. Int. Ed.* **2007**, *46*, 9065; b) B. Liu, H. Xie, H. Wang, L. Wu, Q. Zhao, J. Chen, T. B. Wen, Z. Cao, H. Xia, *Angew. Chem.* **2009**, *121*, 5569; *Angew. Chem. Int. Ed.* **2009**, *48*, 5461.
- [16] a) M. A. Esteruelas, A. M. López, E. Oñate, *Organometallics* **2007**, *26*, 3260; b) R. Ghosh, T. J. Emge, K. Krogh-Jespersen, A. S. Goldman, *J. Am. Chem. Soc.* **2008**, *130*, 11317; c) Y. Jiang, O. Blacque, T. Fox, C. M. Frech, H. Berke, *Organometallics* **2009**, *28*, 4670.
- [17] a) J. Yang, W. M. Jones, J. K. Dixon, N. T. Allison, *J. Am. Chem. Soc.* **1995**, *117*, 9776; b) H.-P. Wu, S. Lanza, T. J. R. Weakley, M. M. Haley, *Organometallics* **2002**, *21*, 2824; c) H.-P. Wu, D. H. Ess, S. Lanza, T. J. R. Weakley, K. N. Houk, K. K. Baldrige, M. M. Haley, *Organometallics* **2007**, *26*, 3957; d) P. M. Johns, W. R. Roper, S. D. Woodgate, L. J. Wright, *Organometallics* **2010**, *29*, 5358.
-