


# Differentiating C–Br and C–Cl Bond Activation by Using Solvent Polarity: Applications to Orthogonal Alkyl–Alkyl Negishi Reactions\*\*

Niloufar Hadei, George T. Achonduh, Cory Valente, Christopher J. O'Brien, and Michael G. Organ\*

Orthogonal reactivity allows one to chemoselectively address one of two reactive centers—both of which are capable of undergoing the same chemical transformation, e.g., cross-couplings—within a bifunctional starting material or intermediate (Scheme 1). This is most commonly achieved by one




**Scheme 1.** General overview of orthogonal cross-coupling strategies.


of two methods: A) fine-tuning the reaction conditions (i.e., temperature, catalyst, additives, etc.) for each reactive center or B) protecting group chemistry. Method A is commonly achieved by taking advantage of the differences in bond enthalpies (C–I > C–Br > C–Cl),<sup>[1]</sup> and more recently has come to include the C<sub>aryl</sub>–O bonds (i.e., aryl carboxylates,<sup>[2]</sup> carbamates, carbonates, and sulfamates<sup>[3]</sup>) that can be efficiently coupled in the presence of a Ni (but not Pd) catalyst, lending itself to orthogonal reaction strategies.<sup>[4]</sup>

Method B has been applied in the form of masked boronic acids, including pinacol esters,<sup>[5]</sup> BF<sub>3</sub>K salts,<sup>[6]</sup> *N*-methylimidodiacetic acid (MIDA),<sup>[7]</sup> and 1,8-diaminonaphthalene (dan)-borane<sup>[8]</sup> derivatives. Strategies based on method B generally discount the possibility of one-pot reaction sequences due to the need for deprotection chemistry. However, bifunctional organodiborane linchpins possessing two of these boronic acid derivatives have been applied successfully in various orthogonal cross-coupling strategies to form aryl–aryl or aryl–vinyl motifs.<sup>[9]</sup> Largely absent from the literature are examples of orthogonal alkyl–alkyl cross-couplings of two unactivated alkyl fragments; to our knowledge, only one

Previous work (Kambe, 2007):



This work:



**Scheme 2.** An overview of known orthogonal alkyl–alkyl cross-couplings.

example<sup>[10]</sup> is known (Scheme 2, previous work).<sup>[11]</sup> Kambe and co-workers showed in a single example that 1-bromo-6-chlorohexane could undergo sequential Kumada–Tamao–Corriu cross-couplings using temperature as the orthogonal trigger in one-pot in the presence of a Cu<sup>II</sup> catalyst. Examples of this type are rare as specially designed, highly active catalysts are typically required to couple unactivated alkyl fragments efficiently,<sup>[12]</sup> with the trade-off that discrimination between C<sub>alkyl</sub>–Br and C<sub>alkyl</sub>–Cl bonds is now less chemoselective. For example, we have shown that NHC–Pd catalysts (NHC = N-heterocyclic carbene), generated in situ from either an imidazolium salt<sup>[13]</sup> or a pre-catalyst,<sup>[14]</sup> namely [Pd-PEPPSI-IPr] (1),<sup>[15]</sup> can effectively couple both unactivated primary C<sub>alkyl</sub>–Br and C<sub>alkyl</sub>–Cl bonds in the Negishi reaction at room temperature.<sup>[16]</sup> During the course of these studies, however, we discovered a unique property that allowed us to chemoselectively couple C<sub>alkyl</sub>–Br bonds in the presence of C<sub>alkyl</sub>–Cl bonds with alkylzinc reagents; that is, solvent polarity. This is a valuable attribute of this reaction, as it provides an opportunity for one-pot orthogonal alkyl–alkyl cross-couplings of bifunctional bromochloroalkanes by a solvent polarity “trigger” (Scheme 2). At its core, the chemoselectivity of these alkyl–alkyl Negishi cross-couplings depend on the ratio of dimethylimidazolidinone (DMI,  $\epsilon$  = 37.6) to tetrahydrofuran (THF,  $\epsilon$  = 7.5).

Preliminarily, a competition experiment (Table 1) was designed to evaluate the chemoselectivity in the alkyl–alkyl Negishi reaction. The *n*BuZnBr for these studies was

[\*] Dr. N. Hadei, G. T. Achonduh, Dr. C. Valente, Dr. C. J. O'Brien, Prof. M. G. Organ  
Department of Chemistry, York University  
4700 Keele Street, Toronto, Ontario M3J 1P3 (Canada)  
Fax: (+1) 416-736-5936  
E-mail: organ@yorku.ca

[\*\*] This work was supported by NSERC (Canada) and the ORDCF (Ontario).

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/ange.201100705>.

**Table 1:** Effect of the DMI:THF solvent ratio on the chemoselectivity in alkyl-alkyl Negishi cross-couplings of bromochloroalkanes.


$\text{Br-CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{Cl} \xrightarrow[\text{DMI:THF (variable), RT, 24 h}]{\text{1 (4 mol\%), LiBr (3.2 equiv), } n\text{BuZnBr (1.6 equiv)}^{[a]}} \text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{Cl} \quad \text{3}$ $\text{Br-CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{Br} \quad \text{4}$					
Entry	DMI:THF	2 [%] <sup>[b]</sup>	3 [%] <sup>[b]</sup>	4 [%] <sup>[b]</sup>	Apparent selectivity <sup>[19]</sup> (3:4)
1	1:2	9	61	5	12.2:1
2	1:1	—	61	12	5.1:1
3	2:1	—	58	20	2.9:1
4	3:1	—	49	19	2.6:1
5	1:0	10	42	17	2.5:1

[a]  $n\text{BuZnBr}$  (1.0 M in DMI) was prepared according to Huo's method.<sup>[17]</sup>

[b] Percent conversion was assessed by GCMS analysis using undecane as a calibrated internal standard.<sup>[20]</sup> Reactions were performed in duplicate and the average conversions are reported.

prepared as a solution in DMI by oxidative addition of activated Zn dust into  $n\text{BuBr}$  following a procedure reported by Huo;<sup>[17]</sup> this method of organozinc preparation is preferred as it requires relatively little synthetic effort, is efficient and produces no inorganic byproducts.<sup>[18]</sup> The optimal solvent ratio in the cross-coupling reaction involving 1-bromo-4-chlorobutane and  $n\text{BuZnBr}$  was found to be 1:2 DMI:THF, where a 12.2:1 apparent selectivity<sup>[19]</sup> ( $C_{\text{alkyl-Br}}:C_{\text{alkyl-Cl}}$ ) was obtained (Table 1, entry 1); increasing the proportion of DMI relative to THF was detrimental to the apparent selectivity (entries 2–5).<sup>[20]</sup> Thus, while the same highly active catalyst that is required in the cross-coupling of alkyl halide electrophiles also leads to a reduction in the intrinsic reactivity differences in  $C_{\text{alkyl-X}}$  bonds, excellent chemoselectivity can still be achieved with these bifunctional precursors through a simple adjustment of the solvent polarity.


A more detailed study was conducted subsequently to further delineate the observed solvent polarity effect (Figure 1). 1-Bromo-3-phenylpropane was subjected to Negishi reaction conditions with  $n\text{BuZnBr}$ , catalyst **1**, and  $\text{LiBr}$ <sup>[18]</sup> at room temperature for 2 h in various ratios of DMI:THF (■ data points). Using pure THF, only a trace amount of product was detected by GCMS analysis, however, the conversions steadily increased as the proportion of DMI was increased relative to THF, and plateaus at near quantitative conversion at ca. 1:1 DMI:THF solvent ratio. The analogous reaction, using 1-chloro-3-phenylpropane and  $\text{LiCl}$ , remained mostly dormant at this solvent ratio, and useful conversions were not observed until the proportion of DMI was increased to establish ca. 2:1 DMI:THF solvent ratio (● data points). To verify that the solvent polarity dependence was not an artifact of a Finkelstein reaction with  $\text{Br}^-$  (that arises from the use of  $n\text{BuZnBr}$ ), wherein substitution first takes place and is followed by efficient cross-coupling of the resultant alkyl bromide,  $n\text{BuZnCl}$  was prepared. The Grignard reagent  $n\text{BuMgCl}$  was first formed from  $n\text{BuCl}$  and Mg turnings, and subsequent transmetalation with  $\text{ZnCl}_2$  provided  $n\text{BuZnCl}$  free from any  $\text{Br}^-$ . Cross-



**Figure 1.** Alkyl-alkyl Negishi cross-couplings carried out at various ratios of DMI:THF. The plot reveals solvent polarity dependence for the coupling of alkyl bromides and chlorides. [a] For data points ranging from 0–59% DMI,  $n\text{BuZnBr}$  in THF was used (0.5 M, Riecke's method);<sup>[22]</sup> 60–100% DMI,  $n\text{BuZnBr}$  in DMI was used (1.0 M, Huo's method). [b]  $n\text{BuZnCl}$  (0.75 M in THF) was prepared by transmetalation of  $n\text{BuMgCl}$  with  $\text{ZnCl}_2$ . [c] Percent conversion was assessed by GCMS analysis using undecane as a calibrated internal standard. Reactions were performed in duplicate and the average conversions are reported. [d] The activation threshold refers to the point at which sufficient conversion is obtained; it is arbitrarily set at 30% for illustrative purposes.

coupling reactions carried out with 1-chloro-3-phenylpropane,  $\text{LiCl}$  and  $n\text{BuZnCl}$  in the presence of catalyst **1** (▲ data points) revealed a similar solvent polarity dependency, ruling out the Finkelstein mechanism. These control reactions also reveal that at a 1:2 DMI:THF solvent ratio, increasing the reaction time to 24 h had little effect on the percent conversion, indicating the reaction is truly switched off. At a 2:1 DMI:THF solvent ratio, a marked increase in percent conversion was obtained when increasing the reaction time from 2 to 24 h. Thus, the solvent polarity trigger is real and can be used as an on/off switch in the cross-coupling reaction of alkyl chlorides and bromides. These findings provide a unique, and to our knowledge, unexplored approach to orthogonal reactivity in bifunctional starting materials. A solvent polarity “trigger” permits the chemoselective cross-coupling of an  $C_{\text{alkyl-Br}}$  bond in the presence of a dormant  $C_{\text{alkyl-Cl}}$  bond, with the latter “activatable” simply by increasing the proportion of DMI in the reaction mixture.

Following this study, a general one-pot double alkyl-alkyl Negishi orthogonal reaction sequence (Scheme 3) was devised whereby a bifunctional bromochloroalkane was



**Scheme 3.** Substrate study for one-pot, orthogonal alkyl-alkyl Negishi cross-couplings that rely on a change in solvent polarity to achieve the orthogonality. Subsequent to the first cross-coupling,  $R^2ZnBr$  was added to the same pot as a solution in DMI (1.0 M) that effectively alters the DMI:THF solvent ratio that “activates” the  $C_{alkyl}-Cl$  bond to cross-coupling. Yields of isolated product are reported, unless indicated otherwise, along with the per-step average for the two cross-couplings. [a] Percent conversion was assessed by GCMS analysis using undecane as a calibrated internal standard.

dissolved in THF along with catalyst **1** and 3.2 equiv of LiBr. A solution of  $nBuZnBr$  (1.0 M in DMI) was added subsequently to achieve a final solvent ratio of 1:2 DMI:THF. After 24 h, the second alkylzinc reagent (1.0 M in DMI) was added to attain a final solvent ratio of 2:1 DMI:THF such that the dormant C–Cl bond was rendered active by the change in solvent polarity. For optimal results, the second cross-coupling required an additional 6.4 equiv of LiCl.<sup>[21]</sup> This one-pot orthogonal reaction sequence provided access to aliphatic derivatives **5–11** in good yields at room temperature.

In conclusion, orthogonal reactivity of bifunctional and unactivated bromochloroalkanes was possible in a double Negishi alkyl-alkyl cross-coupling strategy. The orthogonality was made possible by a relatively simple solvent polarity “trigger”, lending itself to a one-pot approach. To our knowledge, this is the first general strategy for orthogonal alkyl-alkyl cross-couplings, and the first to use solvent polarity as the orthogonal trigger. We are currently investigating the rationale behind the solvent polarity-mediated chemoselectivity of the alkyl-alkyl Negishi reaction.

Received: January 27, 2011

Revised: February 18, 2011

Published online: March 25, 2011

**Keywords:** alkyl-alkyl coupling · cross-coupling · Negishi reaction · orthogonal reactions · solvent polarity

- [1] For examples of orthogonal cross-couplings in one-pot natural product syntheses, see: a) H. Ghasemi, L. M. Antunes, M. G. Organ, *Org. Lett.* **2004**, *6*, 2913–2916; b) M. G. Organ, H. Ghasemi, *J. Org. Chem.* **2004**, *69*, 695–700.
- [2] a) B.-T. Guan, Y. Wang, B.-J. Li, D.-G. Yu, Z.-J. Shi, *J. Am. Chem. Soc.* **2008**, *130*, 14468–14470; b) K. W. Quasdorf, X. Tian, N. K. Garg, *J. Am. Chem. Soc.* **2008**, *130*, 14422–14423.
- [3] K. W. Quasdorf, M. Riener, K. V. Petrova, N. K. Garg, *J. Am. Chem. Soc.* **2009**, *131*, 17748–17749.
- [4] For a Minireview on  $C_{aryl}-O$  cross-couplings, see: L. J. Gooßen, K. Gooßen, C. Stanciu, *Angew. Chem.* **2009**, *121*, 3621–3624; *Angew. Chem. Int. Ed.* **2009**, *48*, 3569–3571.
- [5] N. Iwade, M. Sugimoto, *J. Am. Chem. Soc.* **2010**, *132*, 2548–2549.
- [6] G. A. Molander, D. L. Sandrock, *J. Am. Chem. Soc.* **2008**, *130*, 15792–15793.
- [7] a) E. P. Gillis, M. D. Burke, *J. Am. Chem. Soc.* **2007**, *129*, 6716–6717; b) E. P. Gillis, M. D. Burke, *J. Am. Chem. Soc.* **2008**, *130*, 14084–14085; c) S. J. Lee, K. C. Gray, J. S. Paek, M. D. Burke, *J. Am. Chem. Soc.* **2008**, *130*, 466–468; d) E. P. Gillis, M. D. Burke, *Aldrichimica Acta* **2009**, *42*, 17–27.
- [8] a) H. Noguchi, K. Hojo, M. Sugimoto, *J. Am. Chem. Soc.* **2007**, *129*, 758–759; b) H. Noguchi, T. Shioda, C.-M. Chou, M. Sugimoto, *Org. Lett.* **2008**, *10*, 377–380; c) N. Iwade, M. Sugimoto, *J. Organomet. Chem.* **2009**, *694*, 1713–1717; d) N. Iwade, M. Sugimoto, *Org. Lett.* **2009**, *11*, 1899–1902.
- [9] M. Tobisu, N. Chatani, *Angew. Chem.* **2009**, *121*, 3617–3620; *Angew. Chem. Int. Ed.* **2009**, *48*, 3565–3568.
- [10] Unactivated bromochloroalkanes were used previously in a two-pot orthogonal cross-coupling strategy, wherein the first cross-coupling was an aryl-alkyl or alkyl-alkyl Kumada–Tamao–Corriu reaction, and the subsequent reaction was a Sonogashira cross-coupling. The intrinsic reactivity of the C–Br bond relative to the C–Cl bond acted as the orthogonal “trigger”. See: O. Vechorkin, D. Barmaz, V. Proust, X. Hu, *J. Am. Chem. Soc.* **2009**, *131*, 12078–12079.
- [11] J. Terao, H. Todo, S. A. Begum, H. Kuniyasu, N. Kambe, *Angew. Chem.* **2007**, *119*, 2132–2135; *Angew. Chem. Int. Ed.* **2007**, *46*, 2086–2089.
- [12] For reviews on alkyl-alkyl cross-couplings, see: a) A. C. Frisch, M. Beller, *Angew. Chem.* **2005**, *117*, 680–695; *Angew. Chem. Int. Ed.* **2005**, *44*, 674–688; b) F. Glorius, *Angew. Chem.* **2008**, *120*, 8474–8476; *Angew. Chem. Int. Ed.* **2008**, *47*, 8347–8349; c) A. Rudolph, M. Lautens, *Angew. Chem.* **2009**, *121*, 2694–2708; *Angew. Chem. Int. Ed.* **2009**, *48*, 2656–2670.
- [13] a) N. Hadei, E. A. B. Kantchev, C. J. O’Brien, M. G. Organ, *Org. Lett.* **2005**, *7*, 3805–3807; b) N. Hadei, E. A. B. Kantchev, C. J. O’Brien, M. G. Organ, *J. Org. Chem.* **2005**, *70*, 8503–8507; c) C. J. O’Brien, E. A. B. Kantchev, G. A. Chass, N. Hadei, A. C. Hopkinson, M. G. Organ, D. H. Setiadi, T.-H. Tang, D.-C. Fang, *Tetrahedron* **2005**, *61*, 9723–9735.
- [14] M. G. Organ, S. Avola, I. Dubovyk, N. Hadei, E. A. B. Kantchev, C. J. O’Brien, C. Valente, *Chem. Eur. J.* **2006**, *12*, 4749–4755.
- [15] PEPPSI is an acronym for pyridine-enhanced precatalyst preparation, stabilization, and initiation; IPPr refers to the NHC ligand. [Pd-PEPPSI-IPPr] is commercially available through Aldrich (Product # 669032). For reviews on [Pd-PEPPSI] catalysts, see: a) M. G. Organ, G. A. Chass, D.-c. Fang, A. C. Hopkinson, C. Valente, *Synthesis* **2008**, 2776–2797; b) ref. [16].
- [16] C. Valente, M. E. Belowich, N. Hadei, M. G. Organ, *Eur. J. Org. Chem.* **2010**, 4343–4354.
- [17] S. Huo, *Org. Lett.* **2003**, *5*, 423–425.
- [18] Inorganic salts, such as LiBr or LiCl, have been found to have a significant effect on the alkyl-alkyl Negishi reaction. For studies relating to this topic, see: G. T. Achonduh, N. Hadei, C. Valente,

S. Avola, C. J. O'Brien, M. G. Organ, *Chem. Commun.* **2010**, 46, 4109–4111.

- [19] The term “apparent selectivity” is used as there is the opportunity for a Finkelstein reaction to occur with the  $C_{alkyl}-Cl$  bond and free  $Br^-$  to produce the corresponding alkyl bromide. This side reaction has the net effect of increasing the amount of **4**, thus leading to a decrease in the observed or apparent selectivity. Although this side reaction has not been fully quantified, our observations indicate that it is minor in its contribution to **4**.
- [20] The mass balance of **2** for each entry in Table 1 is attributed primarily to quenched (4-chlorobutyl)zinc bromide (volatility of 1-chlorobutane hinders quantification, b.p. 77–78°C), which is the product of disproportionation of  $nBuZnBr$  with **2**. Trace (<5 %) quantities of 1,8-dichlorooctane are observable in the GCMS chromatograms of each entry, suggesting that the intermediate (4-chlorobutyl)zinc bromide couples with **2** to some extent. Trace quantities of 1,4-dibromobutane (Finkelstein) and dodecane (coupling each site on **2**) are observed in some entries. For the GCMS chromatograms of each entry, see Figure S2 in the Supporting Information.
- [21] Whereas 3.2 equiv of LiBr or LiCl is sufficient for the first cross-coupling, the second cross-coupling requires 6.4 equiv to achieve optimal conversions. For optimization studies, see Table S1 in the Supporting Information.
- [22] LiX is a byproduct in Rieke's method for organozinc formation. When Rieke organozincs are utilized, the total effective equivalents of LiX are reported. For the preparation of Rieke organozincs, see: a) L. Zhu, R. M. Wehmeyer, R. D. Rieke, *J. Org. Chem.* **1991**, 56, 1445–1453; b) R. D. Rieke, M. V. Hanson, J. D. Brown, Q. J. Niu, *J. Org. Chem.* **1996**, 61, 2726–2730.