


Asymmetric Hydrocyanation of α,β -Unsaturated Ketones into β -Cyano Ketones with the $[\text{Ru}(\text{phgly})_2(\text{binap})]/\text{C}_6\text{H}_5\text{OLi}$ Catalyst System**

Nobuhito Kurono, Noriyuki Nii, Yusuke Sakaguchi, Masato Uemura, and Takeshi Ohkuma*

Catalytic asymmetric hydrocyanation of α,β -unsaturated ketones into the corresponding chiral β -cyano ketones is a challenging scientific endeavor. Four major hurdles must be cleared before this reaction can be realized: 1) use of HCN as a cyanide source,^[1] 2) high 1,4-addition selectivity over 1,2-addition, 3) sufficient enantioface selectivity and, 4) high catalytic activity (low catalyst loading). Recently, Shibasaki and co-workers reported pioneering studies on asymmetric 1,4-addition of cyanide to the conjugate enones catalyzed by chiral Gd and Sr compounds.^[2–5] A wide range of 1,4-adducts were obtained in high enantiomeric excess (*ee*), but two equivalents of a *tert*-C₄H₉(CH₃)₂SiCN/2,6-dimethylphenol system were required as a cyanide source to achieve the best catalyst performance.^[6] Furthermore, the substrate-to-catalyst molar ratio (S/C) of 10–200 in these reactions was relatively low.^[2]

Our research group recently reported the asymmetric cyanation of aldehydes and α -keto esters catalyzed by our original $[\text{Ru}(\text{phgly})_2(\text{binap})]/\text{Li}$ salt systems.^[7,8] The corresponding cyanated products were obtained in high *ee*. The spectroscopic analysis suggested that the bimetallic species $[\text{Li-Ru}(\text{phgly})_2(\text{binap})]^+$ acted as a chiral Lewis acidic catalyst. Herein, we describe the efficient asymmetric conjugate addition of HCN to α,β -unsaturated ketones catalyzed by the combined system of $[\text{Ru}(\text{phgly})_2(\text{binap})]$ and $\text{C}_6\text{H}_5\text{OLi}$. The reaction was carried out with an S/C of 200–1000 at -20 – 0°C to afford the β -cyano ketones in up to 98% *ee*.

1-Phenyl-2-buten-1-one (**1a**) was selected as a typical enone substrate to optimize the reaction conditions (Scheme 1 and Table 1). The $[\text{Ru}\{(\text{S})\text{-phgly}\}_2\{(\text{S})\text{-binap}\}]$ ((*S,S,S*)-**3a**) complex was prepared according to the method described in our previous report.^[7] The reaction of **1a** (1.0 mmol) and HCN prepared by mixing (CH₃)₃SiCN (1.5 mmol) and CH₃OH (1.5 mmol) was conducted in *tert*-C₄H₉OCH₃ (6 mL) with (*S,S,S*)-**3a** (20 μmol ,



Scheme 1. Asymmetric hydrocyanation of 1-phenyl-2-buten-1-one (**1a**) with **3** and $\text{C}_6\text{H}_5\text{OLi}$.

Table 1: Asymmetric hydrocyanation of 1-phenyl-2-buten-1-one (**1a**).^[a]

Entry	1a / 3a / PhOLi	Solvent	<i>T</i> [$^\circ\text{C}$]	<i>t</i> [h]	Yield [%] ^[b]	<i>ee</i> [%] ^[b]
1	500:1:1	<i>t</i> BuOMe	25	1	89	89
2	500:0:1	<i>t</i> BuOMe	25	1	35	— ^[c]
3	500:1:0	<i>t</i> BuOMe	25	1	< 1	n.d.
4	500:1:0.5	<i>t</i> BuOMe	25	1	53	90
5	500:1:2	<i>t</i> BuOMe	25	1	> 99	82
6	500:1:1	Et ₂ O	25	1	80	83
7	500:1:1	<i>n</i> -hexane	25	1	21	56
8	500:1:1	CH ₂ Cl ₂	25	1	< 1	n.d.
9	500:1:1	toluene	25	1	44	64
10	500:1:1	<i>t</i> BuOMe	0	5	99	93
11 ^[d]	500:1:1	<i>t</i> BuOMe	0	5	99	93
12 ^[e]	500:1:1	<i>t</i> BuOMe	0	5	98	90
13	1000:1:1	<i>t</i> BuOMe	0	5	98	90
14	2000:1:1	<i>t</i> BuOMe	0	5	< 1	n.d.
15	500:1:1	<i>t</i> BuOMe	-20	18	96	97

[a] Unless otherwise stated, the reactions were carried out using **1a** (1.0 mmol) and HCN (1.5 mmol) in solvent (6 mL) with (*S,S,S*)-**3a** (20 μmol in THF) and $\text{C}_6\text{H}_5\text{OLi}$ (20 μmol in THF) in the ratio given in the Table. HCN was prepared in situ from (CH₃)₃SiCN and CH₃OH in a 1:1 ratio. [b] Data for (*S*)-**2a** were determined by GC on a chiral stationary phase. [c] A racemic product was obtained. [d] Isolated HCN was used. [e] **3b** was used as a catalyst. n.d. = not determined.

S/C = 500) and $\text{C}_6\text{H}_5\text{OLi}$ (20 μmol in THF, 2.0 μmol) at 25°C for 1 hour and gave (*R*)-3-cyano-1-phenyl-1-butanone ((*R*)-**2a**) in 89% yield and 89% *ee* (Table 1, entry 1). Notably, no 1,2-addition product was observed. The reaction catalyzed only by $\text{C}_6\text{H}_5\text{OLi}$ gave racemic **2a** in 35% yield (Table 1, entry 2). No conversion was observed in the reaction with **3a** in the absence of $\text{C}_6\text{H}_5\text{OLi}$ (Table 1, entry 3). The use of a **3a**/ $\text{C}_6\text{H}_5\text{OLi}$ system in a 1:0.5 or 1:2 ratio afforded **2a** in 53%

[*] Dr. N. Kurono, N. Nii, Y. Sakaguchi, M. Uemura, Prof. Dr. T. Ohkuma
Division of Chemical Process Engineering
Faculty of Engineering, Hokkaido University
Sapporo, Hokkaido 060-8628 (Japan)
Fax: (+81) 11-706-6598
E-mail: ohkuma@eng.hokudai.ac.jp

[**] This work was supported by Grants-in-Aid from the Innovation Plaza Hokkaido in JST (No. 01-B07) and JSPS (No. 21350048). N.K. is the grateful recipient of a fellowship from the Global COE Program, "Catalysis as the Basis for Innovation in Materials Science" from MEXT (Japan). M.U. is also the grateful recipient of a JSPS Research Fellowship for Young Scientists. phgly = phenylglycinate, binap = 2,2'-bis(diphenylphosphanyl)-1,1'-binaphthyl.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/ange.201100939>.

yield and 90% *ee*, and >99% yield and 82% *ee*, respectively (Table 1, entries 4 and 5). These observations suggested that **3a** and C₆H₅OLi smoothly formed the 1:1 bimetallic species, [Li-Ru(phgly)₂(binap)]⁺,^[7,8] and the chiral species had a higher reactivity than that of achiral C₆H₅OLi alone. The solvent of choice was *tert*-C₄H₉OCH₃, while (C₂H₅)₂O gave a slightly less satisfactory result (Table 1, entry 6). The yield and enantioselectivity were significantly decreased in less polar solvents (Table 1, entries 7–9). The cyanation at 0°C was completed in 5 hours and afforded the adduct in 93% *ee* (Table 1, entry 10). The same result was obtained by using the isolated HCN prepared according to the method described in the literature,^[9] despite the very low catalyst loading (S/C = 500; Table 1, entry 11). This high reactivity at the low catalyst loading is the important point of difference between Shibasaki's system and the present system.^[2,6] We chose the HCN formed in situ for use in this study for operational and safety reasons. The [Ru(phgly)₂(tol-binap)] (**3b**) exhibited a similar efficiency (Table 1, entry 12).^[10] The high catalytic activity of the **3a**/C₆H₅OLi system allowed us to conduct the cyanation with an S/C of 1000 at 0°C (Table 1, entry 13). The reaction with an S/C of 2000 did not proceed (Table 1, entry 14). The excellent *ee* value of 97% was achieved in the reaction at –20°C, although the reaction took longer to achieve completion (Table 1, entry 15).^[11]


Thus, we selected the reaction conditions using **1a** and 1.5 equivalents of HCN prepared from (CH₃)₃SiCN and CH₃OH in *tert*-C₄H₉OCH₃ with **3a** and C₆H₅OLi (**3a**/C₆H₅OLi = 1:1) at an S/C of 500 at 0°C or –20°C when a higher *ee* value for the product was required (see the Experimental Section). The 1,4-adduct **2a** was quantitatively isolated by column chromatography on silica gel for a reaction carried out on a 3 mmol scale (Table 2, entries 1 and 2). Phenyl ketones (R¹ = C₆H₅ in Scheme 2) with alkyl substituents at the β position (R²), **1b–1g**, reacted with HCN catalyzed by the **3a**/C₆H₅OLi system under the standard conditions and afforded the corresponding β-cyano ketones in 90–96% *ee* (Table 2, entries 3–6, 9, and 10). Although the reactivity of **1c** and **1d** with long alkyl chains was somewhat lower, the enones substituted by secondary and tertiary alkyl groups, **1e–1g**, showed reactivity comparable to that of the methyl-substituted ketone **1a**. The cyanation of **1e** with an S/C of 1000 at 0°C or with an S/C of 500 at –20°C proceeded smoothly and gave **2e** in 92% *ee* and 98% *ee*, respectively (Table 2, entries 7 and 8). Chalcone (**1h**), a β-phenyl enone, was much less reactive than the corresponding β-alkyl substrates, but the cyanation product **2h** was obtained in 92% *ee* and 88% yield in the reaction with an S/C of 200 for 47 hours (Table 2, entry 11).^[12]

A series of substituted phenyl ketones, **1i–1o**, was applied to the asymmetric hydrocyanation. The 2'-Cl phenyl ketone **1i** was treated with **3a** under the typical reaction conditions and gave **2i** in 82% *ee* (Table 2, entry 12). A high enantioselectivity of 95% was achieved in the cyanation of the 3'-Cl substrate **1j** (Table 2, entry 13). The phenyl ketones with an electron-donating CH₃O group at the 3'- or 4'-position, **1k** and **1o**, showed lower reactivity (Table 2, entries 14 and 20). However, an excellent *ee* value of 98% was observed in the reaction of **1k**. The cyanation of substrates with an electron-

Table 2: Asymmetric hydrocyanation of α,β-unsaturated ketones **1**.^[a]

Entry	1	S/C ^[b]	T [°C]	t [h]	Yield [%] ^[c]	<i>ee</i> [%] ^[d]
1	1a	500	0	5	2a , 99	94
2	1a	500	–20	24	2a , 98	97
3	1b	500	0	3	2b , 98	95
4	1c	500	0	18	2c , 94	93
5	1d	500	0	18	2d , 97	90
6	1e	500	0	3	2e , 99	96
7	1e	1000	0	5	2e , 98	92
8	1e	500	–20	18	2e , 98	98
9	1f	500	0	5	2f , 98	95
10	1g	500	0	5	2g , 99	96
11	1h	200	0	47	2h , 88	92
12	1i	500	0	5	2i , 96	82
13	1j	500	0	5	2j , 97	95
14	1k	200	0	12	2k , 96	98
15	1l	500	0	3	2l , 96	95
16	1m	500	0	2	2m , 97	96
17	1m	1000	0	5	2m , 96	94
18	1m	500	–20	12	2m , 98	98
19	1n	500	0	3	2n , 97	95
20	1o	200	0	12	2o , 99	91
21	1p	500	0	12	2p , 99	93
22	1q	500	0	3	2q , 98	95
23	1r	500	0	5	2r , 97 ^[e]	95
24	1s	200 ^[f]	0	24	2s , 80 ^[g]	93

[a] Unless otherwise stated, reactions were conducted using **1** (3.0 mmol) and HCN (4.5 mmol) in *tert*-C₄H₉OCH₃ (18 mL) with a solid (S,S,S)-**3a** and C₆H₅OLi (60 mm in THF). **3a**/C₆H₅OLi = 1:1. HCN was prepared in situ from (CH₃)₃SiCN and CH₃OH in a 1:1 ratio. [b] Substrate-to-catalyst (**3a**) molar ratio. [c] Yield of isolated **2**. [d] Determined by GC or HPLC on a chiral stationary phase. [e] Contaminated by about 1% of an unidentified compound. [f] **3b** was used as a catalyst. [g] The yield determined by GC methods was 99%.




Scheme 2. Asymmetric hydrocyanation of α,β-unsaturated ketones **1** to the β-cyano ketones **2**.

withdrawing Cl, CF₃, or CH₃OCO group at the 4'-position, **1l–1n**, was faster than the reaction of unsubstituted ketone **1a** with maintaining high enantioselectivity (Table 2, entries 15, 16, and 19). The 4'-CF₃ ketone **1m** smoothly reacted with an S/C of 1000 at 0°C or with an S/C of 500 at –20°C and afforded **2m** in 94% *ee* and 98% *ee*, respectively (Table 2, entries 17 and 18).

The cyanation of 2'-naphthyl ketone **1p** under the standard conditions was completed in 12 hours and gave **2p** in 93% *ee* (Table 2, entry 21). The 2'-furyl and 2'-thienyl

ketones, **1q** and **1r**, were converted into the 1,4-adducts, **2q** and **2r**, with 95 % *ee* in both cases (Table 2, entries 22 and 23). The reaction of 3-hepten-2-one (**1s**), an aliphatic enone, with the **3a**/C₆H₅OLi system was slow, but the cyanation with an S/C of 200 at 0 °C for 24 hours afforded **2s** in 99 % yield and 93 % *ee* when **3b** was used instead of **3a** (Table 2, entry 24).^[11] The chiral Gd and Sr catalysts reported by Shibasaki exhibit wider applicability to the reaction of aliphatic and β,β-disubstituted enones.^[2]

The **3b**/C₆H₅OLi catalyst was applied to the regioselective cyanation of a dialkenyl ketone. When cyclohexenyl pentenyl ketone **4** was subjected to the cyanation conditions, the monocyanated product **5** (at the pentenyl group) was obtained in 96 % *ee* (Scheme 3). The regioselectivity was estimated to be greater than 99 %.



Scheme 3. Regioselective cyanation of dienone **4**.

The Ru complex **3a** was so robust that it was recovered by column chromatography on silica gel from the reaction mixture in the open air, and was reusable as a cyanation catalyst with the addition of fresh C₆H₅OLi. As shown in Table 3, **3a** could be used five times in the cyanation of **1a**.

Table 3: Recycled use of **3a** in the hydrocyanation of **1a**.^[a]

Run number ^[b]	Conversion [%] ^[c]	Yield [%] ^[d]	<i>ee</i> [%] ^[c]
1	> 99	97	93
2	> 99	93	94
3	> 99	99	93
4	> 99	99	93
5	> 99	99	92

[a] Reactions were conducted using **1a** (10 mmol) and HCN (15 mmol) in *tert*-C₄H₉OCH₃ (60 mL) at 0 °C for 5 h with a solid (S,S,S)-**3a** and C₆H₅OLi (50 mm in THF). **1a**/**3a**/C₆H₅OLi = 500:1:1 (initial). HCN was prepared in situ from (CH₃)₃SiCN and CH₃OH in a 1:1 ratio. [b] Number of times the catalyst was used. [c] Determined by GC on a chiral stationary phase. [d] Yield of isolated **2a**.

with an initial S/C of 500 at 0 °C with maintaining high enantioselectivity. The total turnover number was about 2500. The notable robustness and reusability of **3a** make it suitable for practical use.

In addition, different α,β-unsaturated ketones **1** were cyanated sequentially with this catalyst-reuse procedure. Table 4 lists the results. The catalyst efficiency and enantioselectivity for all runs were comparable to those of the regular single-run reactions shown in Table 2.

In summary, we have reported here the efficient enantioselective conjugate addition of HCN into the α,β-unsaturated ketones to afford the β-cyano ketones catalyzed by our

Table 4: Sequential hydrocyanation of different substrates.^[a]

Run number ^[b]	1	<i>t</i> [h]	Yield [%] ^[c]	<i>ee</i> [%] ^[d]
1	1n	3	96	93
2	1p	12	96	90
3	1j	5	98	96
4	1l	3	95	95
5	1k ^[e]	12	99	96

[a] Reactions were conducted using **1** (8.2 mmol) and HCN (11.8 mmol) in *tert*-C₄H₉OCH₃ (48 mL) at 0 °C with a solid (S,S,S)-**3a** and C₆H₅OLi (50 mm in THF). **1**/**3a**/C₆H₅OLi = 500:1:1 (initial). HCN was prepared in situ from (CH₃)₃SiCN and CH₃OH in a 1:1 ratio. [b] Number of times the catalyst was used. [c] Yield of isolated **2**. [d] Determined by GC or HPLC on a chiral stationary phase. [e] Reaction using **1k** (3.2 mmol) and HCN (4.8 mmol) in *tert*-C₄H₉OCH₃ (19 mL).

original [Ru(phgly)₂(binap)]/C₆H₅OLi system. The reaction was carried out with an S/C in the range of 200–1000 at –20 °C→0 °C. A series of aryl-, hetero-aryl-, and alkyl-substituted enones was converted into the 1,4-addition products in up to 98 % *ee* without formation of a detectable amount of the 1,2-adducts. The reaction of cyclohexenyl pentenyl ketone afforded the monocyanated product at the less-hindered site in high regio- and enantioselectivity. The robust [Ru(phgly)₂(binap)] complex can be reused with addition of fresh C₆H₅OLi without loss of the stereoselectivity. We hope these findings will contribute to the progress of synthetic organic chemistry.

Experimental Section

The typical procedure for the hydrocyanation of 1-phenyl-2-buten-1-one (**1a**): **Caution:** (CH₃)₃SiCN and HCN that is formed in situ must be used in a well-ventilated fume hood owing to their high toxicity. Ruthenium complex (S,S,S)-**3a** (6.2 mg, 6.1 μmol)^[7,8] was placed in a 50 mL Schlenk flask, and the air present in this apparatus was replaced by argon. Anhydrous CH₃OH (146 mg, 4.6 mmol) was added to this flask, and the mixture was cooled to 0 °C. Then (CH₃)₃SiCN (445 mg, 4.5 mmol) was added in a dropwise manner, and the mixture was stirred for 15 min. To the solution containing HCN, anhydrous *tert*-C₄H₉OCH₃ (18 mL) and C₆H₅OLi (60 mm in THF, 100 μL, 6.0 μmol) were added at 0 °C, and the mixture was stirred for 30 min. Then **1a** (447 mg, 3.1 mmol) was added to this solution in a dropwise manner over 5 min, and the reaction mixture was stirred for 5 h. After the solvent and the volatile compounds were evaporated under reduced pressure, the residue was purified by column chromatography on silica gel to give (S)-**2a** (colorless oil, 531 mg, 99 % yield, 94 % *ee*). [α]_D²⁴ = –6.7 deg cm³ g^{–1} dm^{–1} (*c* = 1.07 g cm^{–3}, CHCl₃); literature^[2a] [α]_D²⁵ = –6.2 deg cm³ g^{–1} dm^{–1} (*c* = 0.6 g cm^{–3}, CHCl₃), 88 % *ee* (absolute configuration was unreported); ¹H NMR (400 MHz, CDCl₃): δ = 1.43 (d, 3 H, *J* = 6.8 Hz, CH₃), 3.23 (dd, 1 H, *J* = 6.5, 17.0 Hz, CHH), 3.31–3.40 (m, 1 H, CHCN), 3.43 (dd, 1 H, *J* = 6.2, 17.0 Hz, CHH), 7.48–7.51 (m, 2 H, aromatic H), 7.60–7.63 (m, 1 H, aromatic H), 7.95–7.97 ppm (m, 2 H, aromatic H); ¹³C NMR (67.7 MHz, CDCl₃): δ = 17.8 (CH₃), 20.5 (CH), 42.2 (CH₂), 122.6 (C), 128.0 (CH), 128.8 (CH), 133.8 (CH), 135.8 (C), 195.1 ppm (C); HRMS (ESI): *m/z* calcd for C₁₁H₁₁ClNO: 208.05292 [*M*+Cl][–]; found: 208.05292. The *ee* value of **2a** was determined by GC on a chiral stationary phase using an InertCap CHIRAMIX column (0.25 mm × 30 m, depth of film = 0.25 μm, GL Science); carrier gas: helium (217 kPa); column temp: 170 °C heating to 179 °C at a rate of 0.5 °C min^{–1}; injection temp: 250 °C; retention time (*t*_R) of (R)-**2a**: 17.5 min (3.1 %), *t*_R of (S)-**2a**: 16.5 min (96.9 %). The *ee* value was not changed by purification with column chromatography. The absolute

configuration was determined after conversion to 2-methyl-4-oxo-4-phenylbutanoic acid in 55% *ee*. $[\alpha]_{\text{D}}^{26} = -18.4 \text{ deg cm}^3 \text{ g}^{-1} \text{ dm}^{-1}$ ($c = 0.592 \text{ g cm}^{-3}$, CHCl_3); literature^[13] $[\alpha]_{\text{D}}^{20} = -32.5 \text{ deg cm}^3 \text{ g}^{-1} \text{ dm}^{-1}$ ($c = 0.69 \text{ g cm}^{-3}$, CHCl_3) for the *S* enantiomer.

Received: February 7, 2011

Revised: March 29, 2011

Published online: May 3, 2011

Keywords: asymmetric catalysis · hydrocyanation · lithium · ruthenium · α,β -unsaturated ketones

- [1] HCN is a toxic compound, but it is utilized in industrial processes for the production of useful chemical compounds, such as α -hydroxy acids, α -amino acids, and methacrylates; for example, see: P. Poechlauer, W. Skranc, M. Wubbolts in *Asymmetric Catalysis on Industrial Scale: Challenge, Approaches and Solutions* (Eds.: H. U. Blaser, E. Schmidt), Wiley-VCH, Weinheim, **2004**, pp. 151–164.
- [2] a) Y. Tanaka, M. Kanai, M. Shibasaki, *J. Am. Chem. Soc.* **2008**, *130*, 6072–6073; b) Y. Tanaka, M. Kanai, M. Shiabasaki, *J. Am. Chem. Soc.* **2010**, *132*, 8862–8863.
- [3] For cyanation of α,β -unsaturated imides, see: a) G. M. Sammis, E. N. Jacobsen, *J. Am. Chem. Soc.* **2003**, *125*, 4442–4443; b) G. M. Sammis, H. Danjo, E. N. Jacobsen, *J. Am. Chem. Soc.* **2004**, *126*, 9928–9929; c) C. Mazet, E. N. Jacobsen, *Angew. Chem.* **2008**, *120*, 1786–1789; *Angew. Chem. Int. Ed.* **2008**, *47*, 1762–1765; d) N. Madhavan, M. Weck, *Adv. Synth. Catal.* **2008**, *350*, 419–425.
- [4] For cyanation of α,β -unsaturated *N*-acylpyrroles, see: a) T. Mita, K. Sasaki, M. Kanai, M. Shibasaki, *J. Am. Chem. Soc.* **2005**, *127*, 514–515; b) I. Fujimori, T. Mita, K. Maki, M. Shiro, A. Sato, S. Furusho, M. Kanai, M. Shibasaki, *Tetrahedron* **2007**, *63*, 5820–5831.
- [5] For cyanation of other activated alkenes using acetone cyanohydrin or ethyl cyanofornate as a cyanide source, see: a) L. Bernardi, F. Fini, M. Fochi, A. Ricci, *Synlett* **2008**, 1857–1861; b) J. Wang, W. Li, Y. Liu, Y. Chu, L. Lin, X. Liu, X. Feng, *Org. Lett.* **2010**, *12*, 1280–1283.
- [6] Use of pure HCN as a cyanide source gave low yield and enantioselectivity in the reaction with the Gd catalyst. The Sr catalyst is expected to be labile in the presence of a large excess of HCN; for details see reference [2].
- [7] a) N. Kurono, K. Arai, M. Uemura, T. Ohkuma, *Angew. Chem.* **2008**, *120*, 6745–6748; *Angew. Chem. Int. Ed.* **2008**, *47*, 6643–6646; b) N. Kurono, M. Uemura, T. Ohkuma, *Eur. J. Org. Chem.* **2010**, 1455–1459.
- [8] N. Kurono, T. Yoshikawa, M. Yamasaki, T. Ohkuma, *Org. Lett.* **2011**, *13*, 1254–1257.
- [9] O. Glemser in *Handbook of Preparative Inorganic Chemistry*, 2nd ed. (Eds.: G. Brauer, R. F. Riley), Academic Press, New York, **1963**, pp. 658–660.
- [10] Tol-binap = 2,2'-bis(di-4-tolylphosphanyl)-1,1'-binaphthyl.
- [11] The β -amino ketones **2a** in 88% *ee* and **2s** in 92% *ee* were obtained by Shibasaki's Gd system (see reference [2a]).
- [12] The conjugate addition of $(\text{CH}_3)_3\text{SiCN}$ to **1h** catalyzed by a chiral sodium phosphate with an S/C of 5 gave **2h** in 71% *ee*. See: J. Yang, S. Wu, F.-X. Chen, *Synlett* **2010**, 2725–2728.
- [13] R. V. Hoffman, H.-O. Kim, *J. Org. Chem.* **1995**, *60*, 5107–5113.