

Berichtigung

Fused Indolines by Palladium-Catalyzed Asymmetric C–C Coupling Involving an Unactivated Methylene Group

M. Nakanishi, D. Katayev, C. Besnard,
E. P. Kündig* ————— **7576–7579**

Angew. Chem. **2011**, 123

DOI 10.1002/ange.201102639

Die Fußnote [a] von Tabelle 3 in dieser Zuschrift muss wie folgt lauten: „[a] Condition: **1** (0.2 mmol), $[\{\text{Pd}(\eta^3\text{-cinnamyl})\text{Cl}\}_2]$ (0.005 mmol), NHC-HI (0.01 mmol), cesium pivalate (0.2 mmol), and cesium carbonate (0.3 mmol), mesitylene.“ Weiterhin ist das korrekte Literaturzitat [3] hier gezeigt.

[3] a) An enantioselective (37% *ee*) coupling between *n*-butylboronic acid and a CH₃ group in 2-propylpyridine was reported: B.-F. Shi, N. Mangel, Y.-H. Zhang, J.-Q. Yu, *Angew. Chem.* **2008**, 120, 4960–4964; *Angew. Chem. Int. Ed.* **2008**, 47, 4882–4886; b) for a Pd⁰-catalyzed asymmetric β-arylation of enantiotopic CH₃ groups in β-position of carboxylic esters (55–70% yield, up to 54% *ee*), see: A. Renaudat, L. Jean-Gerard, R. Jazsar, C. E. Kefalidis, E. Clot, O. Baudoin, *Angew. Chem.* **2010**, 122, 7419–7423; *Angew. Chem. Int. Ed.* **2010**, 49, 7261–7265.