


Multicomponent Reactions

Rhodium-Catalyzed Three-Component Cross-Addition of Silylacetylenes, Alkynyl Esters, and Electron-Deficient Alkenes or Isocyanates**


Yuki Hoshino, Yu Shibata, and Ken Tanaka*

The transition-metal-catalyzed hydroalkynylation of unsaturated compounds with terminal alkynes has been extensively studied for the atom-economical synthesis of functionalized alkynes.^[1] Although a number of the hydroalkynylation reactions of alkynes,^[2] allenes,^[3] and alkenes^[4] with terminal alkynes has been reported, the cross-addition reactions of terminal alkynes and two unsaturated compounds through the formation of two C–C bonds are rare.^[5] In particular, the reactions of terminal alkynes and two different unsaturated compounds are difficult as a result of the challenge in controlling the chemoselectivity.^[6,7] Ogata, Fukuzawa, and co-workers realized such challenging transformations by using bulky triisopropylsilylacetylene as a terminal alkyne and nickel(0)/phosphine complexes as catalysts.^[6,7] In 2009, they reported the nickel-catalyzed three-component cross-addition of triisopropylsilylacetylene and two different internal alkynes ($R^1, R^2, R^4 \neq H, R^3 = CH_2OR^5$) by controlling the chemoselective addition between two different internal alkynes (Scheme 1a).^[6a] In 2011, they realized the more difficult cross-addition of two different terminal alkynes [triisopropylsilylacetylene and terminal alkynes ($R^1 = \text{alkyl}, R^2 = H$)] and internal alkynes ($R^3, R^4 = Ar$) by controlling the chemoselective C–H bond activation between two different terminal alkynes as well as the chemoselective addition between the terminal and internal alkynes (Scheme 1a).^[6b] In 2010, they also reported that norbornene is able to participate in this three-component cross-addition to give the corresponding 1,5-enynes (Scheme 1a).^[7] However, the available alkenes were unfortunately strictly limited to norbornene derivatives. Herein, we report the unprecedented transition-metal-catalyzed chemo-, regio-, and stereoselective three-component cross-addition of silylacetylenes, alkynes, and alkenes, thus leading to substituted 1,3-dienes (Scheme 1b) by using cationic rhodium(I) complexes as catalysts.^[8] The analogous unprecedented three-component cross-addition involving isocyanates in place of alkenes is also disclosed.^[9]



Scheme 1. Transition-metal-catalyzed three-component cross-addition of silylacetylenes, alkynes, and alkenes.

Our research group previously reported that a cationic rhodium(I)/(cod)₂ (cod = 1,5-cyclooctadiene) complex is a highly effective catalyst for the [2+2+1] cross-cyclotrimerization of silylacetylenes and two alkynyl esters, thus leading to substituted silylfulvenes.^[10,11] For example, triisopropylsilylacetylene (**1a**) reacted with two ethyl 2-butynoates (**2a**) to give the corresponding silylfulvene **5aa** in high yield (Scheme 2).^[10] Interestingly, the use of the sterically demand-



Scheme 2. Rhodium-catalyzed reactions of triisopropylsilylacetylene (**1a**) and alkynyl esters **2**.

ing methyl 4,4-dimethyl-2-butynoate (**2b**) in place of **2a** did not afford the cross-cyclotrimerization product **5aa**, but instead delivered the cross-dimerization product **3ab**^[2g] as a major product along with the linear cross-trimerization product **4ab**,^[5k] which was generated from **2b** and two units of **1a**, as a minor product (Scheme 2).


We anticipated that when adding an excess of an electron-deficient alkene, this alkene would be incorporated into the product as a third component in place of **1a**. Pleasingly, the

[*] Y. Hoshino, Y. Shibata, Prof. Dr. K. Tanaka

Department of Applied Chemistry, Graduate School of Engineering
Tokyo University of Agriculture and Technology
Koganei, Tokyo 184-8588 (Japan)
E-mail: tanaka-k@cc.tuat.ac.jp
Homepage: <http://www.tuat.ac.jp/~tanaka-k/>

[**] This work was supported partly by Grants-in-Aid for Scientific Research (Nos. 23105512, 20675002, and 21-906) from the MEXT (Japan). We thank Umicore for generous support in supplying rhodium complexes.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/ange.201204646>.



Scheme 3. Rhodium-catalyzed reactions of triisopropylsilylacetylene (**1a**), the alkynyl ester **2b**, and acrylate **6a**.

reaction of **1a**, **2b**, and butyl acrylate (**6a**, 2 equiv; Scheme 3) at 60 °C afforded the expected three-component cross-addition product **7aba** along with the corresponding dehydrogenated product **8aba**, however, the major product was the cross-dimerization product **3ab**.

To improve the yield of **7aba**, the effect of the solvents was first examined (entries 1–6, Table 1). Ether solvents were

Table 1: Effect of catalysts and solvents on reaction of **1a**, **2b**, and **6a**.^[a]

Entry	Catalyst	Solvent	Yield [%] ^[b]		
			7aba	8aba	3ab
1	$[Rh(cod)_2]BF_4$	1,4-dioxane	20	11	43
2	$[Rh(cod)_2]BF_4$	THF	32	14	26
3	$[Rh(cod)_2]BF_4$	DME	16	6	49
4	$[Rh(cod)_2]BF_4$	$(CH_2Cl)_2$	0	0	48
5	$[Rh(cod)_2]BF_4$	toluene	0	3	39
6	$[Rh(cod)_2]BF_4$	CH_3CN	0	0	0
7	$\{[RhCl(cod)]_2\}/2AgBF_4$	THF	30	7	24
8	$\{[IrCl(cod)]_2\}/2AgBF_4$	THF	0	0	49
9	$[Rh(nbd)]_2BF_4$	THF	0	6	58
10	$\{[RhOAc(cod)]_2\}$	THF	7	8	41
11	$[Rh(cod)_2]BF_4/rac\text{-}binap$	THF	0	0	0
12	$\{[RhCl(cod)]_2\}$	THF	0	0	0
13	$[Rh(cod)_2]OTf$	THF	31	0	14
14	$[Rh(cod)_2]SbF_6$	THF	46	0	31
15	$\{[RhCl(cod)]_2\}/2NaBAR_F^{[c]}$	THF	0	2	0
16 ^[d]	$[Rh(cod)_2]SbF_6$	THF	62	0	10
17 ^[d,e]	$[Rh(cod)_2]SbF_6$	THF	74	0	16

[a] Reactions were conducted using the catalyst (0.010 mmol of metal), **1a** (0.11 mmol), **2b** (0.10 mmol), **6a** (0.20 mmol), and solvent (2.0 mL) at 60 °C for 16 h. [b] Determined by 1H NMR spectroscopy.

[c] $NaBAR_F$ = sodium tetrakis[3,5-bis(trifluoromethyl)phenyl]borate.

[d] **6a**: 5 equiv. [e] At 80 °C. DME = 1,2-dimethoxyethane, THF = tetrahydrofuran.



effective for this transformation (entries 1–3) and the use of THF furnished **7aba** in the highest yield (entry 2). However, less-coordinating solvents (entries 4 and 5) and a highly coordinating solvent (entry 6) did not furnish **7aba** at all. Next, the effect of catalysts was examined (entries 7–14). An in situ generated cationic rhodium(I)/cod complex catalyzed the formation of **7aba** as well as a significant amount of the oligomerization products from **1a** (entry 7). The use of a cationic iridium(I)/cod complex, $[Rh(nbd)]_2BF_4$, and $\{[Rh(cod)OAc]_2\}$ predominantly furnished the cross-dimerization product **3ab** (entries 8–10). The use of a cationic rhodium(I)/*rac*-binap complex [binap = 2,2'-bis(diphenylphosphino)-1,1'-

binaphthyl] and $\{[RhCl(cod)]_2\}$ failed to catalyze the reaction (entries 11 and 12). Interestingly, changing the counter anion from BF_4 to OTf (entry 13) or SbF_6 (entry 14) completely suppressed the formation of **8aba**, and the use of $[Rh(cod)_2]SbF_6$ furnished **7aba** in the highest yield (entry 14). However, the use of BAR_F as the counter anion significantly suppressed the substrate conversion (entry 15). Increasing the amount of **6a** improved the selectivity of **7aba** over **3ab**, but the reaction rate decreased (entry 16). Elevating the reaction temperature to 80 °C increased the reaction rate and further improved the yield of **7aba** (entry 17).

The scope of this rhodium(I)-catalyzed three-component cross-addition was examined as shown in Scheme 4.^[12] With respect to electron-deficient alkenes, various acrylates (**6a–c**) and methyl vinyl ketone (**6d**) were suitable substrates for this process, while the use of *N,N*-dimethylacrylamide (**6e**) lowered the yield of the corresponding three-component cross-addition product. With respect to the silylacetylenes, not only triisopropylsilylacetylene (**1a**) but also *tert*-butyldiphenyl- and *tert*-butyldimethylsilylacetylenes (**1b** and **1c**, respectively) could be employed, although excess **6d** (20 equiv) was required. With respect to the alkynyl esters, tertiary-alkyl- (**2b**), secondary-alkyl- (**2c,d**), primary-alkyl- (**2e–g**), and aryl-substituted (**2h,i**) alkynyl esters could react with **1a** and excess **6d** (20 equiv) to give the corresponding three-component cross-addition products in moderate to good yields. Importantly, these reactions proceeded with complete regio- and stereoselectivity. Interestingly, the reactions of propargyl acetates **2j,k** proceeded at room temperature to give the corresponding three-component cross-addition products in good yields, although that of benzoate **2l** required an elevated temperature. The use of the less Lewis-acidic $[Rh(cod)_2]BF_4$ in place of $[Rh(cod)_2]SbF_6$ was necessary to avoid the undesired elimination of carboxylic acids. In these reactions, stereoselectivities were opposite to those using **2b–i**.

As the third addition component, isocyanates could be employed in place of electron-deficient alkenes by using the less Lewis-acidic $[Rh(cod)_2]BF_4$ as a catalyst (Scheme 5).^[13] The primary and secondary alkyl isocyanates **8a,b** reacted with **1a** and **2b** to give the corresponding enyne amides **9**, and the use of the arylisocyanates **8c–e** improved the product yields. Although the product yield decreased, *tert*-butyldiphenylsilylacetylene (**1b**) could also be employed.



A mechanistic proposal for this rhodium(I)-catalyzed three-component cross-addition reaction is shown in Scheme 6. Carborhodation of the alkynyl ester **2** with the rhodium acetylide **A**, generated by the reaction of the silylacetylene **1**^[14] and the cationic rhodium(I) complex, affords the (alkenyl)rhodium **B**, which reacts with electron-deficient alkene **6** to generate the intermediate **C**. Protonation of **C** affords the 1,3-enyne **7** and regenerates the rhodium(I) catalyst. The isocyanate **8** is also able to react with the (alkenyl)rhodium **B** to give the intermediate **D**, the protonation of which affords the enyne amide **9**. In contrast, the reaction of the rhodium acetylide **A** with the propargyl ester **2** [$R^1 = CR^5R^6(OCOR^7)$] affords the (alkenyl)rhodium **E**, in which the carbonyl oxygen atom coordinates to rhodium, through *E/Z* isomerization of **B**.^[15] This intermedi-




Scheme 4. Three-component cross-addition of silylacetylenes **1a–c**, alkynyl esters **2b–l**, and electron-deficient alkenes **6a–e**. Reaction conditions: $[\text{Rh}(\text{cod})_2]\text{SbF}_6$ (0.010 mmol), **1a–c** (0.22 mmol), **2b–l** (0.20 mmol), **6a–e** (1.0–4.0 mmol), and THF (2.0 mL) were used. Cited yields are of isolated products. $[\text{a}] [\text{Rh}(\text{cod})_2]\text{BF}_4$ (0.010 mmol) was used. Bz = benzoyl.

ate **E** reacts with **6** to give the intermediate **F**, subsequent protonation of which affords the 1,3-enyne **7'**.

Transformations of the present three-component cross-addition products were briefly examined as shown in Scheme 7. Treatment of the 1,3-enynes **7abd** and **7aed** with TBAF (tetra-*n*-butylammonium fluoride) furnished the terminal alkynes **10a** and **10b**, respectively, in good to high yields. The palladium-catalyzed Sonogashira cross-coupling of **10a** with iodobenzene furnished the phenylacetylene




Scheme 5. Three-component cross-addition of silylacetylenes **1a,b**, alkynyl ester **2b**, and isocyanates **8a–e**. Reaction conditions: [Rh-(cod)₂]BF₄ (0.010 mmol), **1a,b** (0.22 mmol), **2b** (0.20 mmol), **8a–e** (1.0 mmol), and THF (2.0 mL) were used. Reported yields are of those of the isolated products.



Scheme 6. Proposed reaction mechanism.

derivative **11** in high yield. Alternatively, the rhodium(I)/H₈-binap [2,2'-bis(diphenylphosphino)-5,5',6,6',7,7',8,8'-octahydro-1,1'-binaphthyl] complex catalyzed [2+2+2] cycloaddition of **10b** with the 1,6-diyne **12** to furnish the densely substituted benzene **13** in good yield. In contrast, treatment of



Scheme 7. Transformations of three-component cross-addition products. Ts = 4-toluenesulfonyl.

the enyne amide **9abc** with TBAF furnished the desilylated cyclization product **14** in high yield.

In conclusion, we have developed an unprecedented rhodium-catalyzed chemo-, regio-, and stereoselective three-component cross-addition of a silylacetylene, alkynyl ester, and electron-deficient alkene, thereby leading to substituted 1,3-dienes. In addition, the analogous and unprecedented three-component cross-addition involving isocyanates in place of electron-deficient alkenes has also been developed. Future work will focus on further exploitation of the catalytic three-component cross-addition reactions by using the cationic rhodium(I) complexes as catalysts.

Received: June 14, 2012

Published online: August 21, 2012

Keywords: alkenes · alkynes · multicomponent reactions · rhodium · synthetic methods

- [1] For reviews, see: a) B. M. Trost, A. H. Weiss, *Adv. Synth. Catal.* **2009**, *351*, 963; b) S. Fujimori, T. F. Knöpfel, P. Zarotti, T. Ichikawa, D. Boyall, E. M. Carreira, *Bull. Chem. Soc. Jpn.* **2007**, *80*, 1635; c) L. Zani, C. Bolm, *Chem. Commun.* **2006**, 4263; d) P. G. Cozzi, R. Hilgraf, N. Zimmermann, *Eur. J. Org. Chem.* **2004**, 4095; e) C. Wei, Z. Li, C.-J. Li, *Synlett* **2004**, 1472; f) L. Pu, *Tetrahedron* **2003**, *59*, 9873; g) D. E. Frantz, R. Fässler, C. S. Tomooka, E. M. Carreira, *Acc. Chem. Res.* **2000**, *33*, 373.
- [2] a) B. M. Trost, M. C. McIntosh, *J. Am. Chem. Soc.* **1995**, *117*, 7255; b) B. M. Trost, A. J. Frontier, *J. Am. Chem. Soc.* **2000**, *122*, 11727; c) L. Chen, C. Li, *Tetrahedron Lett.* **2004**, *45*, 2771; d) T. Hirabayashi, S. Sakaguchi, Y. Ishii, *Adv. Synth. Catal.* **2005**, *347*, 872; e) H. Katayama, H. Yari, M. Tanaka, F. Ozawa, *Chem. Commun.* **2005**, 4336; f) J.-i. Ito, M. Kitase, H. Nishiyama, *Organometallics* **2007**, *26*, 6417; g) T. Nishimura, X.-X. Guo, K. Ohnishi, T. Hayashi, *Adv. Synth. Catal.* **2007**, *349*, 2669; h) T. Katagiri, H. Tsurugi, A. Funayama, T. Satoh, M. Miura, *Chem. Lett.* **2007**, *36*, 830; i) N. Tsukada, S. Ninomiya, Y. Aoyama, Y. Inoue, *Org. Lett.* **2007**, *9*, 2919; j) T. Katagiri, H. Tsurugi, T. Satoh, M. Miura, *Chem. Commun.* **2008**, 3405; k) K. Ogata, O. Oka, A. Toyota, N. Suzuki, S.-i. Fukuzawa, *Synlett* **2008**, 2663;

- l) N. Matsuyama, K. Hirano, T. Satoh, M. Miura, *J. Org. Chem.* **2009**, *74*, 3576.
- [3] a) B. M. Trost, G. Kottirsch, *J. Am. Chem. Soc.* **1990**, *112*, 2816; b) M. Yamaguchi, K. Omata, M. Hiramata, *Tetrahedron Lett.* **1994**, *35*, 5689; c) M. Yamaguchi, Y. Kido, K. Omata, M. Hiramata, *Synlett* **1995**, 1181; d) D. Bruyere, R. Grigg, J. Hinsley, R. K. Hussain, S. Korn, C. O. De La Cierva, V. Sridharan, J. Wang, *Tetrahedron Lett.* **2003**, *44*, 8669; e) M. Rubin, J. Markov, S. Chuprakov, D. J. Wink, V. Gevorgyan, *J. Org. Chem.* **2003**, *68*, 6251; f) T. Nishimura, X.-X. Guo, T. Hayashi, *Chem. Asian J.* **2008**, *3*, 1505; g) X.-X. Guo, T. Sawano, T. Nishimura, T. Hayashi, *Tetrahedron: Asymmetry* **2010**, *21*, 1730.
- [4] a) G. I. Nikishin, I. P. Kovalev, *Tetrahedron Lett.* **1990**, *31*, 7063; b) R. V. Lerum, J. D. Chisholm, *Tetrahedron Lett.* **2008**, *49*, 1505; c) A. Tenaglia, L. Giordano, G. Buono, *Org. Lett.* **2006**, *8*, 4315; d) J. Yin, J. D. Chisholm, *Chem. Commun.* **2006**, 632; e) T. Nishimura, X.-X. Guo, N. Uchiyama, T. Katoh, T. Hayashi, *J. Am. Chem. Soc.* **2008**, *130*, 1576; f) M. Shirakura, M. Suginoe, *J. Am. Chem. Soc.* **2008**, *130*, 5410; g) M. Shirakura, M. Suginoe, *Org. Lett.* **2009**, *11*, 523; h) K. Kohno, K. Nakagawa, T. Yahagi, J.-C. Choi, H. Yasuda, T. Sakakura, *J. Am. Chem. Soc.* **2009**, *131*, 2784; i) M. Shirakura, M. Suginoe, *J. Am. Chem. Soc.* **2009**, *131*, 5060; j) E. Fillion, A. K. Zoritto, *J. Am. Chem. Soc.* **2009**, *131*, 14608; k) T. Nishimura, T. Sawano, T. Hayashi, *Angew. Chem.* **2009**, *121*, 8201; *Angew. Chem. Int. Ed.* **2009**, *48*, 8057; l) T. Nishimura, T. Sawano, S. Tokui, T. Hayashi, *Chem. Commun.* **2010**, 46, 6837; m) T. Nishimura, T. Sawano, K. Ou, T. Hayashi, *Chem. Commun.* **2011**, 47, 10142.
- [5] a) M. Ishikawa, J. Ohshita, Y. Ito, A. Minato, *J. Chem. Soc. Chem. Commun.* **1988**, 804; b) H.-F. Klein, M. Mager, *Organometallics* **1992**, *11*, 3174; c) H. Matsuzaka, Y. Takagi, Y. Ishii, M. Nishio, M. Hidai, *Organometallics* **1995**, *14*, 2153; d) B. M. Trost, M. T. Sorum, C. Chan, A. E. Harms, G. Rühter, *J. Am. Chem. Soc.* **1997**, *119*, 698; e) A. Haskel, T. Straub, A. K. Dash, M. S. Eisen, *J. Am. Chem. Soc.* **1999**, *121*, 3014; f) A. Haskel, J. Q. Wang, T. Straub, T. G. Neyron, M. S. Eisen, *J. Am. Chem. Soc.* **1999**, *121*, 3025; g) A. K. Dash, I. Gourevich, J. Q. Wang, J. Wang, M. Kapon, M. S. Eisen, *Organometallics* **2001**, *20*, 5084; h) S.-i. Yoshimitsu, S. Hikichi, M. Akita, *Organometallics* **2002**, *21*, 3762; i) J. Q. Wang, M. S. Eisen, *J. Organomet. Chem.* **2003**, *670*, 97; j) M. V. Jiménez, E. Sola, F. J. Lahoz, L. A. Oro, *Organometallics* **2005**, *24*, 2722; k) N. Matsuyama, H. Tsurugi, T. Satoh, M. Miura, *Adv. Synth. Catal.* **2008**, *350*, 2274; l) K. Ogata, H. Murayama, J. Sugawara, N. Suzuki, S.-i. Fukuzawa, *J. Am. Chem. Soc.* **2009**, *131*, 3176.
- [6] a) K. Ogata, J. Sugawara, S.-i. Fukuzawa, *Angew. Chem.* **2009**, *121*, 6194; *Angew. Chem. Int. Ed.* **2009**, *48*, 6078; b) K. Ogata, Y. Atsuumi, S.-i. Fukuzawa, *Org. Lett.* **2011**, *13*, 122.
- [7] K. Ogata, J. Sugawara, Y. Atsuumi, S.-i. Fukuzawa, *Org. Lett.* **2010**, *12*, 148.
- [8] For precedented examples of the transition-metal-catalyzed intermolecular reactions involving successive insertion of internal alkynes and alkenes, see: a) K. Shibata, T. Satoh, M. Miura, *Org. Lett.* **2005**, *7*, 1781; b) T. Kurahashi, H. Shinokubo, A. Osuka, *Angew. Chem.* **2006**, *118*, 6484; *Angew. Chem. Int. Ed.* **2006**, *45*, 6336; c) H. Horiguchi, K. Hirano, T. Satoh, M. Miura, *Adv. Synth. Catal.* **2009**, *351*, 1431.
- [9] For precedented examples of the transition-metal-catalyzed intermolecular reactions involving the arylmetalation of isocyanates, see: a) T. Koike, M. Takahashi, N. Arai, A. Mori, *Chem. Lett.* **2004**, *33*, 1364; b) T. Miura, Y. Takahashi, M. Murakami, *Chem. Commun.* **2007**, 3577; c) Y. Kuninobu, K. Kikuchi, Y. Tokunaga, Y. Nishina, K. Takai, *Tetrahedron* **2008**, *64*, 5974; d) K. D. Hesp, R. G. Bergman, J. A. Ellman, *J. Am. Chem. Soc.* **2011**, *133*, 11430.
- [10] Y. Shibata, K. Tanaka, *Angew. Chem.* **2011**, *123*, 11109; *Angew. Chem. Int. Ed.* **2011**, *50*, 10917.

- [11] Our research group reported the rhodium(I)-catalyzed cross-trimerization of propargyl esters and two arylacetylenes leading to substituted dihydropentalenes, a reaction initiated by the hydroalkynylation of propargyl esters with arylacetylenes. See: Y. Shibata, K. Noguchi, K. Tanaka, *Org. Lett.* **2010**, *12*, 5596.
- [12] When the yields of the desired products **7** were low, the following cross-dimerization products **3** or cross-trimerization products **5** were generated as major by-products: **7abe** (**3ab**: 71 %), **7cbd** (**3cb**: 33 %), **7add** (**5ad**: 35 %), **7afd** (**5af**: 31 %), **7agd** (**5ag**: 46 %), **7aid** (**5ai**: 14 %), **7ald** (**5al**: 21 %).
- [13] The following cross-dimerization products **3** and cross-trimerization products **4** were generated as major by-products: **9aba** (**3ab**: 14 % and **4ab**: 11 %), **9abb** (**3ab**: 28 % and **4ab**: 40 %), **9abc** (**3ab**: 35 % and **4ab**: 9 %), **9abd** (**3ab**: 22 % and **4ab**: 12 %), **9abe** (**3ab**: 44 % and **4ab**: 7 %), **9bbc** (**3bb**: 24 % and **4bb**: 40 %).
- [14] For examples of the formation of rhodium acetylides from silylacetylenes, see: Ref. [2g].
- [15] We believed that this isomerization step from the intermediate **B** to intermediate **E** might be irreversible.