


An Organosilicon Cluster with an Octasilacuneane Core: A Missing Silicon Cage Motif**

Shintaro Ishida,* Kyohei Otsuka, Yuki Toma, and Soichiro Kyushin*

Organosilicon clusters (caged oligosilanes) are a fascinating class of molecules because 1) they can be regarded as partial structures of bulk crystalline or amorphous silicon and 2) the electronic properties of silicon clusters depend on silicon numbers and structural modification.^[1] Therefore, synthetic studies on organosilicon clusters have been extensively developed. Persilapolyhedranes,^[2–4] decasilaadamantane,^[5] pentasila[1.1.1]propellane,^[6] persila[*n*]staffane (catenated silicon cages),^[7] and related siliconoids^[8,9] have all been shown to possess unique electronic characteristics owing to developed σ conjugation among the caged Si–Si bonds. Octasilacuneane, a silicon analog of cuneane^[10] (pentacyclo[3.3.0.0^{2,4}.0^{3,7}.0^{6,8}]octane), is a missing saturated silicon cluster motif, but potentially important because octasilacuneane is an isomeric silicon cage of octasilacubane,^[4] one of the most well-known silicon polyhedra. Moreover, through theoretical studies at the MP2/6-31G(d) + ZPE level, Versteeg and Koch have demonstrated that parent octasilacuneane (Si₈H₈) is only 3.3 kcal mol^{−1} less stable than the isomeric octasilacubane, which is in sharp contrast to the energetic relation between cubane and cuneane: cubane (C₈H₈) is much more unstable (by 43.7 kcal mol^{−1}) than cuneane in their calculations.^[11] Cubanes readily isomerize to the corresponding cuneanes by metal-ion catalyzed skeletal rearrangements.^[10a]

Herein, we report the synthesis, structure, and properties of cyclotetrasilane-fused persilacuneane (**1**), which consists of sixteen silicon atoms. Compound **1** was found in the course of our study on ladder oligosilanes.^[12] As shown in Equation (1), the reductive tetramerization (formation of eight Si–Si bonds during the reaction) of tetrachlorocyclotetrasilane **2**^[13] with sodium gave silicon cluster **1** as air-sensitive, thermally stable orange crystals in 15 % yield. No structural isomers of **1** were found in the reaction mixture. The ¹H NMR spectrum of the reaction mixture showed a set of eight singlet signals from the *tert*-butyl protons of **1** together with a very broad signal



around 1–2 ppm, owing to an intractable polymeric mixture. Compound **1** was sparingly soluble in common organic solvents.

The construction of the silicon cage could be ascribed to the formation of an energetically reasonable structure by successive bond formation during the reaction. To elucidate this idea, we estimated the relative energies of two fused persilacubanes and three fused persilacuneanes (**3**-*D*_{2d}, **3**-*D*_{4d}, **4**-*C*₂, **4**-*C*_{2v}, and **4**-*C*_s) by theoretical calculations at the B3LYP/6-31G(d) level, as shown in Figure 1.^[14] Whereas




Figure 1. Relative energies (kcal mol^{−1}) of compounds **3** and **4**.

parent persilacuneane is less stable than persilacubane by 3.3 kcal mol^{−1} at the MP2/6-31G(d) + ZPE level^[11] (8.3 kcal mol^{−1} at the B3LYP/6-31G(d) level),^[14] **4**-*C*₂, a model of **1**, is the most stable among the five isomers. The introduction of cyclotetrasilane units into the silicon cluster can change the relative energies of different silicon cages.

The molecular structure of **1** was determined by X-ray crystallography, as shown in Figure 2.^[15] Compound **1** has crystallographic *C*₂ symmetry with the axis through the midpoints of the Si1–Si1* and Si5–Si5* bonds, and a well-developed network structure consisting of sixteen silicon

[*] Dr. S. Ishida,^[14] Dr. K. Otsuka, Y. Toma, Prof. Dr. S. Kyushin
Department of Chemistry and Chemical Biology, Graduate School of
Engineering, Gunma University, Kiryu, Gunma 376-8515 (Japan)
E-mail: kyushin@gunma-u.ac.jp

[†] Present address: Department of Chemistry, Graduate School of
Science, Tohoku University, Aoba-ku, Sendai 980-8578 (Japan)
E-mail: sishida@m.tohoku.ac.jp

[**] This work was supported in part by Grants-in-Aid for Scientific
Research from the Ministry of Education, Culture, Sports, Science
and Technology (Japan) and the Japan Society for the Promotion of
Science. We also thank Prof. Dr. Takeshi Yamanobe, Gunma
University (Japan) for measurement of the CP-MAS NMR spectrum.

Supporting information for this article is available on the WWW
under <http://dx.doi.org/10.1002/anie.201208506>.




Figure 2. Molecular structure of **1**. Thermal ellipsoids are drawn at the 30% probability level. Selected bond lengths [pm]: Si1–Si2 238.2(1), Si1–Si4 238.4(1), Si1–Si1* 238.0(2), Si1–Si8* 233.2(1), Si2–Si3 252.2(1), Si3–Si4 241.3(1), Si4–Si5 240.4(1), Si4–Si8* 235.4(1), Si5–Si6 241.8(1), Si5–Si8 236.1(1), Si5–Si5* 241.0(2), Si6–Si7 251.0(1), Si7–Si8 242.0(1).

atoms. The Si–Si bond lengths of peripheral cyclotetrasilane moieties (238.2(1)–252.2(1) pm) are slightly elongated because of steric hindrance from the *tert*-butyl groups.

Although the recently reported compound, pentasil-[1.1.1]propellane, has inverted tetrahedral silicon atoms with a significantly long bridgehead Si–Si bond (263.6(1) pm; its bonding character has been controversial)^[6] the bridgehead Si–Si bond lengths of **1** (Si1–Si4 238.4(1) pm, Si5–Si8 236.1(1) pm) are within standard values.^[16,17] However, the bridgehead silicon atoms (Si1, Si4, Si5, and Si8) have highly deformed trigonal monopyramidal structures, as illustrated in Figure 3. Each silicon atom has a planar structure, in which




Figure 3. Geometries around the deformed silicon atoms of **1**.

the sum of three Si–Si–Si bond angles is almost 360°. The remaining Si–Si bond is almost perpendicular to the plane in the cases of Si5 and Si8, whereas the remaining Si–Si bond is tilted to form a three-membered ring in the cases of Si1 and Si4. Similar trigonal monopyramidal silicon atoms have been found in the silicon clusters, tris(triisopropylsilyl)silane,^[18] and unsolvated tris(di-*tert*-butylmethylsilyl)silyllithium.^[19]

Figure 4 shows the solid state ²⁹Si CP-MAS NMR spectrum of **1**. Eight signals were observed, which is





Figure 4. ²⁹Si cross polarization magic angle spinning (CP-MAS) NMR spectrum of **1** at room temperature with signals calculated by the GIAO method at the B3LYP/6-311 + G(2d,p) level. Atom numbers correspond to those in Figure 2.

consistent with the molecular structure in the solid state. These signals were essentially reproduced by GIAO calculation at the B3LYP/6-311 + G(2d,p) level using the optimized structure, and can be assigned as follows: Three characteristic signals shifted upfield at –71.8, –62.4, and –56.6 ppm are due to the ²⁹Si nuclei of the cyclotrisilane rings (Si1, Si4, and Si8).^[20,21] The signals at 25.0, 29.1, 35.4, and 39.0 ppm can be assigned to any of ²⁹Si nuclei of Si2, Si3, Si5, and Si6.^[22,23]

Notably, the ²⁹Si signal of Si7 at 66.4 ppm is shifted rather downfield. A possible explanation may be given by the large paramagnetic term (σ_p) of Si7. This term has a predominant effect on the chemical shift of ²⁹Si NMR. According to a theoretical equation,^[24] σ_p is roughly inversely proportional to the averaged transition energy. The downfield shift of the ²⁹Si signal of Si7 is probably due to its contribution to the HOMO and the LUMO of **1**.

Theoretical calculations were carried out to obtain insight into the electronic characterization of **1**. Frontier orbitals are located on the silicon skeleton, indicating well-developed σ conjugation (see the Supporting Information). The major contribution of the HOMO is an out-of-phase interaction among the $\sigma_{Si1-Si4}$, $\sigma_{Si1-Si8^*}$, and $\sigma_{Si7-Si8^*}$ orbitals and among the $\sigma_{Si1^*-Si4^*}$, σ_{Si1^*-Si8} , and $\sigma_{Si7-Si8}$ orbitals. The major contribution of the LUMO consists of an out-of-phase interaction between the pseudo- π^* orbitals of two peripheral cyclotetrasilane rings (Si5–Si6–Si7–Si8 and Si5*–Si6*–Si7*–Si8* rings).

The UV/Vis spectra of **1** and our heptacyclic ladder oligosilane, *i*Pr₂₀Si₁₆ (**5**),^[25] both of which consist of sixteen



silicon atoms, are compared in Figure 5. Compound **5** has well-developed σ conjugation along the long axis of the molecule and shows the lowest energy absorption as a distinctive band at 464 nm.^[25] In contrast, the absorption of **1** does not show clear bands above 250 nm and tails to ca. 570 nm, except for some shoulders at 345 and 380 nm. The absorption tailing is characteristic of silicon polyhedral




Figure 5. UV/Vis spectra of **1** in methylcyclohexane and **5** in hexane at room temperature. **1** (—), **5** (----).

compounds: UV/Vis absorption of hexasilaprismane (DipSi)₆ (Dip = 2,6-diisopropylphenyl) tails to ca. 500 nm,^[3] and those of octasilacubanes (DepSi)₈ (Dep = 2,6-diethylphenyl) and [(*t*BuMe₂Si)Si]₈ tail to ca. 470 nm.^[1c,4d] The absorption edge of **1** is remarkably red-shifted compared with those of these hexasilaprismane and octasilacubanes.

The characteristic tailing of **1** is probably due to close energy levels of molecular orbitals with a small band gap. The HOMO and the LUMO have energy levels close to other molecular orbitals, and therefore, absorption bands overlap to give a closely overlapped absorption curve without peaks. From these results, organosilicon cluster **1** may be regarded as an intermediate model between small organosilicon molecules and silicon semiconductors that have the band structures of orbitals.^[26]

In summary, we have synthesized a cyclotetrasilane-fused octasilacubane **1**, which consists of sixteen silicon atoms, by reductive tetramerization of tetrachlorocyclotetrasilane. X-ray crystallography, theoretical studies, and analysis by NMR and UV/Vis spectroscopy revealed the unique structural and electronic features of **1**, which are due to ring strain and developed σ conjugation. Investigations of the reactivity of **1** and the expansion of σ conjugation by introducing functional groups are now ongoing and will appear in the near future.

Received: October 22, 2012

Published online: January 30, 2013

Keywords: cluster compounds · cuneanes · organosilicon · propellanes · structure elucidation

- [1] For reviews on cyclic oligosilanes and silicon cage compounds, see: a) E. Hengge, R. Janoschek, *Chem. Rev.* **1995**, 95, 1495–1526; b) A. Sekiguchi, H. Sakurai, *Adv. Organomet. Chem.* **1995**, 37, 1–38; c) A. Sekiguchi, S. Nagase in *The Chemistry of Organic Silicon Compounds*, Vol. 2 (Eds.: Z. Rappoport, Y. Apeloig), Wiley, Chichester, **1998**, pp. 119–152; d) E. Hengge, H. Stüger in *The Chemistry of Organic Silicon Compounds*,

Vol. 2 (Eds.: Z. Rappoport, Y. Apeloig), Wiley, Chichester, **1998**, pp. 2177–2216.

- [2] For tetrasilatetrahedranes, see: a) N. Wiberg, C. M. M. Finger, K. Polborn, *Angew. Chem.* **1993**, 105, 1140–1142; *Angew. Chem. Int. Ed. Engl.* **1993**, 32, 1054–1056; b) M. Ichinohe, M. Toyoshima, R. Kinjo, A. Sekiguchi, *J. Am. Chem. Soc.* **2003**, 125, 13328–13329.
- [3] For hexasilaprismane, see: A. Sekiguchi, T. Yatabe, C. Kabuto, H. Sakurai, *J. Am. Chem. Soc.* **1993**, 115, 5853–5854.
- [4] For octasilacubanes, see: a) H. Matsumoto, K. Higuchi, Y. Hoshino, H. Koike, Y. Naoi, Y. Nagai, *J. Chem. Soc. Chem. Commun.* **1988**, 1083–1084; b) K. Furukawa, M. Fujino, N. Matsumoto, *Appl. Phys. Lett.* **1992**, 60, 2744–2745; c) A. Sekiguchi, T. Yatabe, H. Kamatani, C. Kabuto, H. Sakurai, *J. Am. Chem. Soc.* **1992**, 114, 6260–6262; d) H. Matsumoto, K. Higuchi, S. Kyushin, M. Goto, *Angew. Chem.* **1992**, 104, 1410–1412; *Angew. Chem. Int. Ed. Engl.* **1992**, 31, 1354–1356; e) K. Furukawa, M. Fujino, N. Matsumoto, *J. Organomet. Chem.* **1996**, 515, 37–41; f) M. Unno, T. Matsumoto, K. Mochizuki, K. Higuchi, M. Goto, H. Matsumoto, *J. Organomet. Chem.* **2003**, 685, 156–161.
- [5] J. Fischer, J. Baumgartner, C. Marschner, *Science* **2005**, 310, 825.
- [6] D. Nied, R. Köppe, W. Kloppe, H. Schnöckel, F. Breher, *J. Am. Chem. Soc.* **2010**, 132, 10264–10265, and references therein.
- [7] T. Iwamoto, D. Tsushima, E. Kwon, S. Ishida, H. Isobe, *Angew. Chem.* **2012**, 124, 2390–2394; *Angew. Chem. Int. Ed.* **2012**, 51, 2340–2344.
- [8] For a Si₈(SiR₃)₆ cluster, see: G. Fischer, V. Huch, P. Mayer, S. K. Vasisht, M. Veith, N. Wiberg, *Angew. Chem.* **2005**, 117, 8096–8099; *Angew. Chem. Int. Ed.* **2005**, 44, 7884–7887.
- [9] Scheschke and co-workers have proposed the term “siliconoids” for silicon clusters with unsubstituted silicon vertexes; see: a) D. Scheschke, *Angew. Chem.* **2005**, 117, 3014–3016; *Angew. Chem. Int. Ed.* **2005**, 44, 2954–2956; b) K. Abersfelder, A. J. P. White, H. S. Rzepa, D. Scheschke, *Science* **2010**, 327, 564–566; c) K. Abersfelder, A. J. P. White, R. J. F. Berger, H. S. Rzepa, D. Scheschke, *Angew. Chem.* **2011**, 123, 8082–8086; *Angew. Chem. Int. Ed.* **2011**, 50, 7936–7939; d) K. Abersfelder, A. Russell, H. S. Rzepa, A. J. P. White, P. R. Haycock, D. Scheschke, *J. Am. Chem. Soc.* **2012**, 134, 16008–16016.
- [10] For cuneanes, see: a) L. Cassar, P. E. Eaton, J. Halpern, *J. Am. Chem. Soc.* **1970**, 92, 6366–6368; b) R. Gleiter, S. Brand, *Tetrahedron Lett.* **1994**, 35, 4969–4972; c) H. Irngartinger, S. Strack, R. Gleiter, S. Brand, *Acta Crystallogr. Sect. C* **1997**, 53, 1145–1148; d) R. Gleiter, S. Brand, *Chem. Eur. J.* **1998**, 4, 2532–2538; e) M. V. Roux, J. Z. Dávalos, P. Jiménez, R. Notario, O. Castaño, J. S. Chickos, W. Hanshaw, H. Zhao, N. Rath, J. F. Liebman, B. S. Farivar, A. Bashir-Hashemi, *J. Org. Chem.* **2005**, 70, 5461–5470.
- [11] U. Versteeg, W. Koch, *J. Comput. Chem.* **1994**, 15, 1151–1162.
- [12] For reviews on ladder oligosilanes, see: a) H. Matsumoto, S. Kyushin, M. Unno, R. Tanaka, *J. Organomet. Chem.* **2000**, 611, 52–63; b) S. Kyushin, H. Matsumoto, *Adv. Organomet. Chem.* **2003**, 49, 133–166.
- [13] Tetrachlorocyclotetrasilane **2** was prepared by the head-to-head reductive dimerization of 1,2-dichlorodisilane **S3** and subsequent dephenylchlorination. For details of the synthesis and X-ray crystal analyses of **1** and **2**, see the Supporting Information.
- [14] We also calculated the energies of the parent cubane, cuneane, octasilacubane, and octasilacubane at the B3LYP/6-31G(d) level. Our results were comparable to the previous report by Versteeg and Koch.^[11] See the Supporting Information for details.
- [15] CCDC 906219 (**S4**), 906220 (**2**), and 906221 (**1**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

- [16] The silicon skeleton of **1** contains hitherto unknown persila-[3.2.1]- and [3.2.2]propellanes and spiro[3.3]- and [4.3]oligosilanes as substructures. Compound **1** also has a spiro-[3.2]hexasilane moiety. Some spiro[3.2]hexasilane derivatives have been reported.^[8,17d]
- [17] For spirooligosilanes, see: a) P. Boudjouk, R. Sooriyakumaran, *J. Chem. Soc. Chem. Commun.* **1984**, 777–778; b) T. Iwamoto, M. Tamura, C. Kabuto, M. Kira, *Science* **2000**, 290, 504–506; c) J. Hlina, C. Mechtler, H. Wagner, J. Baumgartner, C. Marschner, *Organometallics* **2009**, 28, 4065–4071; d) T. Iwamoto, Y. Furiya, H. Kobayashi, H. Isobe, M. Kira, *Organometallics* **2010**, 29, 1869–1872.
- [18] a) S. Kyushin, H. Sakurai, H. Matsumoto, *Chem. Lett.* **1998**, 107–108; b) P. P. Gaspar, A. M. Beatty, T. Chen, T. Haile, D. Lei, W. R. Winchester, J. Braddock-Wilking, N. P. Rath, W. T. Klooster, T. F. Koetzle, S. A. Mason, A. Albinati, *Organometallics* **1999**, 18, 3921–3932.
- [19] M. Nakamoto, T. Fukawa, V. Ya. Lee, A. Sekiguchi, *J. Am. Chem. Soc.* **2002**, 124, 15160–15161.
- [20] The ²⁹Si signals of silyl-substituted cyclotrisilanes appear in upfield regions: (Et₃Si)₃Si₃ = –174.4 ppm,^[21a] (tBuMe₂Si)₃Si₃ = –156.5 ppm,^[21b] and (iPr₂MeSi)₃Si₃ = –148.6 ppm.^[21b]
- [21] a) H. Matsumoto, A. Sakamoto, Y. Nagai, *J. Chem. Soc. Chem. Commun.* **1986**, 1768–1769; b) M. Kira, T. Iwamoto, T. Maruyama, T. Kuzuguchi, D. Yin, C. Kabuto, H. Sakurai, *J. Chem. Soc. Dalton Trans.* **2002**, 1539–1544.
- [22] The ²⁹Si signals of Si₂, Si₃, and Si₆ were observed within the range of reported *tert*-butyl-substituted cyclotetrasilanes: tBu₇Si₄H = 31.7 and 41.8 ppm,^[23a] *trans*-tBu₆Si₄H₂ = 32.0 ppm,^[23a] tBu₂(tBu₂MeSi)₃Si₄Br₃ = 37.3 ppm,^[23b] and *trans*-tBu₆Si₄Br₂ = 24.7 ppm.^[23c]
- [23] a) S. Kyushin, H. Sakurai, H. Matsumoto, *J. Organomet. Chem.* **1995**, 499, 235–240; b) A. Sekiguchi, T. Matsuno, M. Ichinohe, *J. Am. Chem. Soc.* **2001**, 123, 12436–12437; c) S. Kyushin, H. Kawai, H. Matsumoto, *Organometallics* **2004**, 23, 311–313.
- [24] a) N. F. Ramsey, *Phys. Rev.* **1950**, 78, 699–703; b) A. Saika, C. P. Slichter, *J. Chem. Phys.* **1954**, 22, 26–28.
- [25] S. Kyushin, Y. Ueta, R. Tanaka, H. Matsumoto, *Chem. Lett.* **2006**, 35, 182–183.
- [26] For reviews on group 13 and group 14 clusters, and their intermediate role between small molecules and the bulk state, see: a) H. Schnöckel, *Dalton Trans.* **2005**, 3131–3136; b) A. Schnepf, *Chem. Soc. Rev.* **2007**, 36, 745–758; c) A. Schnepf, *New J. Chem.* **2010**, 34, 2079–2092. See also Reference [9d].