


Catalytic Asymmetric [8+2] Cycloaddition: Synthesis of Cycloheptatriene-Fused Pyrrole Derivatives**

Mingsheng Xie, Xiaohua Liu, Xiaoxia Wu, Yunfei Cai, Lili Lin, and Xiaoming Feng*

Dedicated to Professor Guo-Qiang Lin on the occasion of his 70th birthday

The higher-order cycloaddition represents a highly valuable strategy for the preparation of medium-sized ring systems.^[1] For instance, fulvenes have been applied as 6 π components in enantioselective [6+2]^[2a] and [6+3]^[2b,c] cycloadditions. For the construction of seven-membered rings,^[3–5] asymmetric [5+2]^[4a,b] and [4+3]^[5] reactions were developed. However, as a direct method to construct 7,5-fused heterocyclic compounds, the asymmetric [8+2] cycloaddition was less studied.^[1d,e] The heteroanalogues of heptafulvenes are the most-studied substrates for the [8+2] cycloadditions.^[6–15] In general, the multiple reactivity profiles of the polyene analogues resulted in modest chemical efficiency in such higher-order cycloadditions. For example, tropones have been reported to serve as 4 π components in asymmetric [4+2] cycloadditions,^[10a] while as 6 π components in asymmetric [6+3]^[11a,b] and [6+4] cycloadditions.^[12a] Tropothione received scant concern owing to its inaccessibility and dimerization properties.^[14] Comparatively, azaheptafulvene has attracted much attention for it could undergo efficient [8+2] cycloadditions with electron-deficient π systems, as well as for its accessibility and stability (Scheme 1).^[15] However, to our knowledge no examples of catalytic asymmetric [8+2] cycloadditions have been reported to date. The cycloheptatriene unit in the initially formed [8+2] adducts could easily undergo isomerization, which led to the occurrence of epimerization (Scheme 1).^[16] Moreover, it is too difficult to obtain optically pure bicyclo[5.3.0] ring products by separating these isomers. In the context of ongoing projects in developing asymmetric catalysis promoted by an *N,N'*-dioxide–metal complex,^[17] we became interested in the enantioselective [8+2] cycloaddition. The azaheptafulvene was chosen as excellent 8 π component in the catalytic asymmetric [8+2] cycloaddition,



Scheme 1. The [8+2] cycloaddition of azaheptafulvenes.

affording functionalized cycloheptatriene-fused pyrrole derivatives,^[18] which represent a framework found in a number of biologically active molecules.^[19]

Initially, we investigated the reaction of azaheptafulvene **1a** with diethyl 2-benzylidenemalonate **2a** promoted by *N,N'*-dioxide **L1** with various metals (Table 1, entries 1–4). With Ni(BF₄)₂·6H₂O as the central metal salt, the direct [8+2] cycloaddition product **3a** was obtained with the highest *ee* value (75 % *ee*). However, the allylic rearrangement product **4a** was detected as the major form, along with the 1,5- and 1,7-H shift products in less than 5 % yield (Table 1, entries 1–4). Both the bulky amide subunits and the amino acid backbone of the ligands were crucial for the improvement of the outcome of the reaction. Decrease of the steric hindrance of the amide substituents deteriorated both yields and enantioselectivities (Table 1, entries 4–6). Compared with L-proline-derived ligand **L1** and L-pipecolic acid derived **L4**, L-ramipril-derived *N,N'*-dioxide **L5** afforded a higher enantioselectivity (Table 1, entry 8 versus entries 4 and 7). Complete conversion with 93 % *ee* was obtained by using catalyst **L5**–Ni(BF₄)₂·6H₂O. The ratio of the product **3a** and the allylic rearrangement product **4a** was nearly unchanged even when varying the counteranion of the nickel salts, reaction solvents, temperature, and additives (see the Supporting Information for details); therefore we wondered whether the rearrangement reaction was promoted by the Lewis acid catalyst or other factors. The issue was resolved by column chromatography experiments. When the reaction mixture was purified by chromatography using silica gel (basification by 3 % Et₃N), the formation of the allylic rearrangement product **4a** was almost suppressed, albeit a trace of Et₃N was eluted as well (Table 1, entry 9). Accordingly, another ground matrix was chosen as the stationary phase, and the desired product **3a** was obtained in 99 % yield and 93 % *ee* after being purified by flash chromatography on basic Al₂O₃ (Table 1, entry 10).

[*] M. S. Xie, Prof. Dr. X. H. Liu, X. X. Wu, Y. F. Cai, Dr. L. L. Lin, Prof. Dr. X. M. Feng
Key Laboratory of Green Chemistry & Technology, Ministry of Education, College of Chemistry, Sichuan University
Chengdu 610064 (China)
E-mail: xmfeng@scu.edu.cn

Prof. Dr. X. M. Feng
State Key Laboratory of Applied Organic Chemistry
Lanzhou University, Lanzhou 730000 (China)

[**] We thank the National Basic Research Program of China (973 Program: No. 2011CB808600), the National Natural Science Foundation of China (Nos. 21021001, 21290182, and 21172151) and the Ministry of Education (No. 20110181130014) for financial support.


Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/ange.201209601>.

Table 1: Optimization of the reaction conditions.^[a]

		1a Ar = 4-MeC ₆ H ₄	2a: R = CO ₂ Et 2b: R = CN	metal-ligand (x mol%) CH ₂ Cl ₂ , 0 °C	3a: R = CO ₂ Et 3ab: R = CN	4a: R = CO ₂ Et 4ab: R = CN	
Entry	Metal	Ligand	x	t [h]	Yield [%] ^[b]	3a/4a ^[c]	ee (3a) [%] ^[c]
1	Sc(OTf) ₃	L1	10	40	31	22:78	52
2	Mg(ClO ₄) ₂	L1	10	40	99	25:75	58
3	Co(BF ₄) ₂ ·6 H ₂ O	L1	10	40	75	21:79	63
4	Ni(BF ₄) ₂ ·6 H ₂ O	L1	10	40	94	21:79	75
5	Ni(BF ₄) ₂ ·6 H ₂ O	L2	10	20	88	21:79	35
6	Ni(BF ₄) ₂ ·6 H ₂ O	L3	10	20	73	19:81	— ⁷
7	Ni(BF ₄) ₂ ·6 H ₂ O	L4	10	20	92	25:75	39
8	Ni(BF ₄) ₂ ·6 H ₂ O	L5	10	20	99	26:74	93
9 ^[d]	Ni(BF ₄) ₂ ·6 H ₂ O	L5	10	20	99	97:3	93
10 ^[e]	Ni(BF ₄) ₂ ·6 H ₂ O	L5	10	20	99	99:1	93
11 ^[e]	Ni(BF ₄) ₂ ·6 H ₂ O	L5	0.5	15	99	99:1	93
12 ^[e,f]	Ni(BF ₄) ₂ ·6 H ₂ O	L5	0.5	2	99	99:1	92
13 ^[e,f]	Ni(BF ₄) ₂ ·6 H ₂ O	L5	0.1	17	99	99:1	88
14 ^[e,f,g]	Ni(BF ₄) ₂ ·6 H ₂ O	L5	0.1	2	99	99:1	0
15 ^[f]	—	—	—	24	17	99:1	0
16 ^[f]	—	L5	10	24	15	99:1	0

[a] Reaction conditions: metal/L (1:1), **1a** (0.11 mmol), and **2a** (0.1 mmol) in CH₂Cl₂ (0.5 mL) at 0 °C, and the products were purified by silica gel column chromatography. [b] Yields of isolated products.

[c] Determined by HPLC on a chiral stationary phase and ¹H NMR spectroscopy, and the d.r. > 95:5 was determined by ¹H NMR spectroscopy. [d] The product was purified by silica gel (basification by 3% Et₃N) column chromatography. [e] The product was purified by flash chromatography on basic Al₂O₃. [f] At 35 °C. [g] **2b** was used to react with **1a** to give **3ab/4ab**.



Remarkably, the catalytic efficiency was also satisfactory when the catalyst loading was lowered to 0.5 mol % (Table 1, entry 11 versus entry 10). The reaction time could be shortened to 2 h with similar results when the reaction was carried out at 35 °C (Table 1, entry 12 versus entry 11). Further lowering the catalyst loading to 0.1 mol % afforded the product in quantitative yield with slightly decreased enantioselectivity (Table 1, entry 13). Meanwhile, when ethyl 2-cyano-3-phenylacrylate **2b** was used as the 2π component, the corresponding racemic adduct was obtained within 2 h (Table 1, entry 14). This result showed that the bidentate dicarbonyl functional groups were indispensable for the chiral control. Besides, blank experiments without metal salt were performed, and lower yields and racemic adducts with high diastereoselectivities (> 95:5 d.r.) were obtained, thereby indicating that the central metal salt was essential for the chiral induction (Table 1, entries 15–16).

This simple catalyst system was quite versatile in the [8+2] cycloaddition with other azaheptafulvenes and alkylidene malonates (Table 2). The 8π components bearing electron-

Table 2: Substrate scope of the asymmetric [8+2] cycloaddition.^[a]

1	2			3			
Entry	R ¹	R ² , R ³	3	<i>x</i>	<i>t</i> [h]	Yield [%] ^[b]	<i>ee</i> [%] ^[c]
1	Me	Et, Ph	3 a	0.5	15	99	93
2 ^[d]	MeO	Et, Ph	3 b	1	24	98	92
3	Cl	Et, Ph	3 c	0.5	9	99	96
4	Br	Et, Ph	3 d	0.5	9	95	95
5	Br	Et, Ph	3 d	0.2	15	95	94
6	Me	Me, Ph	3 e	0.5	15	99	93
7	Me	Me, 2-FC ₆ H ₄	3 f	1	15	99	91
8	Me	Et, 3,4-Cl ₂ C ₆ H ₃	3 g	1	11	99	91
9	Me	Et, 4-BrC ₆ H ₄	3 h	1	21	93	95
10	Me	Et, 4-CF ₃ C ₆ H ₄	3 i	1	15	95	94
11	Me	Et, 3-NO ₂ C ₆ H ₄	3 j	1	16	99	92
12	Me	Me, 4-NO ₂ C ₆ H ₄	3 k	1	15	99	94
13	Me	Me, 3-MeC ₆ H ₄	3 l	1	17	99	95
14	Me	Et, 4-MeC ₆ H ₄	3 m	1	15	99	94
15	Me	Et, 4-MeOC ₆ H ₄	3 n	0.5	9	99	96
16	Me	Et, 2-thienyl	3 o	1	16	99	97
17	Me	Et, cyclohexyl	3 p	1	8	90	95
18 ^[d,e]	Me	Et, Et	3 q	2	35	38	93
19 ^[d,f]	Me	Et, <i>i</i> Pr	3 r	2	22	92	96

[a] Reaction conditions: Ni(BF₄)₂·6H₂O/L5 (1:1), **1** (0.11 mmol), and **2** (0.1 mmol) in CH₂Cl₂ (0.5 mL) at 0 °C, and the products were purified by flash chromatography on basic Al₂O₃. [b] Yields of isolated products.

[c] Determined by HPLC on a chiral stationary phase, and the d.r. > 95:5 was determined by ¹H NMR spectroscopy. The total amount of isomerization products was less than 5%. [d] The azaheptafulvene was generated in situ from the tetrafluoroborate salt with Et₃N. [e] **3q/4q** = 3:7. [f] **3r/4r** = 23:76. **4** is the corresponding allylic rearrangement product (see Scheme 1).

donating or electron-withdrawing substituents on the *N*-aryl group underwent efficient cyclization in excellent enantioselectivities (92–96% *ee*), high diastereoselectivities (> 95:5 d.r.), and high yields (Table 2, entries 1–5). Dimethyl 2-benzylidenemalonate gave comparable enantioselectivity and yield (Table 2, entry 6). The reactions were remarkably tolerant of functional groups regardless of the electronic properties and steric hindrance of the substituents on the 2π components (Table 2, entries 7–15). Among the substrates **2** investigated, *o*-fluoro-substituted and disubstituted ones afforded slightly lower enantioselectivities (Table 2, entries 7, 8 versus entries 9–15). Additionally, olefins containing heteroaromatic and aliphatic groups also underwent the [8+2] cycloaddition, affording the desired adducts with high enantioselectivities (93–97% *ee*; Table 2, entries 16–19). Allylic rearrangement products could not be excluded for substrates **2q** and **2r** derived from aliphatic aldehydes (Table 2, entries 18–19). In all cases, the cyclization reactions exhibited excellent diastereoselectivity, and only *trans*-isomers of **3** were found. The absolute configuration of the cycloheptatriene-fused pyrrole derivative **3d** was determined to be (2*S*, 3*aS*) by single-crystal X-ray diffraction analysis.^[20]

To further evaluate the synthetic potential of the catalytic system, a gram-scale synthesis of chiral cycloheptatriene-

- Williams, J.-Y. Yoon, *J. Am. Chem. Soc.* **2006**, *128*, 6302; b) R. Shintani, H. Nakatsu, K. Takatsu, T. Hayashi, *Chem. Eur. J.* **2009**, *15*, 8692; For examples of non-asymmetric [5+2] cycloadditions, see: c) P. A. Wender, H. Takahashi, B. Witulski, *J. Am. Chem. Soc.* **1995**, *117*, 4720; d) B. M. Trost, F. D. Toste, H. Shen, *J. Am. Chem. Soc.* **2000**, *122*, 2379; e) P. A. Wender, C. M. Barzilay, A. J. Dyckman, *J. Am. Chem. Soc.* **2001**, *123*, 179; f) G. Zuo, J. Louie, *J. Am. Chem. Soc.* **2005**, *127*, 5798; g) A. Fürstner, K. Majima, R. Martín, H. Krause, E. Kattinig, R. Goddard, C. W. Lehmann, *J. Am. Chem. Soc.* **2008**, *130*, 1992; h) P. A. Wender, A. B. Lesser, L. E. Sirois, *Org. Synth.* **2011**, *88*, 109; i) X. Xu, P. Liu, A. Lesser, L. E. Sirois, P. A. Wender, K. N. Houk, *J. Am. Chem. Soc.* **2012**, *134*, 11012.
- [5] For examples of catalytic asymmetric [4+3] cycloadditions, see: a) M. Harmata, S. K. Ghosh, X. Hong, S. Wacharasindhu, P. Kirchhoefer, *J. Am. Chem. Soc.* **2003**, *125*, 2058; b) J. Huang, R. P. Hsung, *J. Am. Chem. Soc.* **2005**, *127*, 50; c) X. Dai, H. M. L. Davies, *Adv. Synth. Catal.* **2006**, *348*, 2449; d) R. P. Reddy, H. M. L. Davies, *J. Am. Chem. Soc.* **2007**, *129*, 10312; e) M. Gullías, J. Durán, F. López, L. Castedo, J. L. Mascareñas, *J. Am. Chem. Soc.* **2007**, *129*, 11026; f) R. Shintani, M. Murakami, T. Hayashi, *J. Am. Chem. Soc.* **2007**, *129*, 12356; g) B. D. Schwartz, J. R. Denton, Y. Lian, H. M. L. Davies, C. M. Williams, *J. Am. Chem. Soc.* **2009**, *131*, 8329; h) R. Shintani, M. Murakami, T. Tsuji, H. Tanno, T. Hayashi, *Org. Lett.* **2009**, *11*, 5642; i) I. Alonso, H. Faustino, F. López, J. L. Mascareñas, *Angew. Chem.* **2011**, *123*, 11698; *Angew. Chem. Int. Ed.* **2011**, *50*, 11496.
- [6] For examples of [8+2] cycloadditions with dienyisobenzofurans as 8 π components, see: a) Y. Luo, J. W. Herndon, F. Cervantes-Lee, *J. Am. Chem. Soc.* **2003**, *125*, 12720; b) L. Zhang, Y. Wang, C. Buckingham, J. W. Herndon, *Org. Lett.* **2005**, *7*, 1665; c) Y. Chen, S. Ye, L. Jiao, Y. Liang, D. K. Sinha-Mahapatra, J. W. Herndon, Z.-X. Yu, *J. Am. Chem. Soc.* **2007**, *129*, 10773; d) P. Roy, B. K. Ghorai, *Tetrahedron Lett.* **2011**, *52*, 5668.
- [7] For examples of [8+2] cycloadditions with indolizines as 8 π components, see: a) J. C. Godfrey, *J. Org. Chem.* **1959**, *24*, 581; b) A. Galbraith, T. Small, R. A. Barnes, V. Boekelheide, *J. Am. Chem. Soc.* **1961**, *83*, 453; c) M. Aginagalde, Y. Vara, A. Arrieta, R. Zangi, V. L. Cebolla, A. Delgado-Camón, F. P. Cossío, *J. Org. Chem.* **2010**, *75*, 2776.
- [8] For examples of [8+2] cycloadditions with benzothietes as 8 π components, see: a) K. Kanakarajan, H. Meier, *J. Org. Chem.* **1983**, *48*, 881; b) H. Meier, A. Mayer, *Angew. Chem.* **1994**, *106*, 493; *Angew. Chem. Int. Ed. Engl.* **1994**, *33*, 465; c) H. Meier, B. Rose, D. Schollmeyer, *Liebigs Ann.* **1997**, 1173.
- [9] For examples of [8+2] cycloadditions with heptafulvenes as 8 π components, see: a) W. von E. Doering, D. W. Wiley, *Tetrahedron* **1960**, *11*, 183; b) C.-Y. Liu, J. Mareda, K. N. Houk, F. R. Fronczek, *J. Am. Chem. Soc.* **1983**, *105*, 6714; c) C.-Y. Liu, H.-Y. Shie, S.-Y. Chen, C.-Y. You, W.-C. Wang, L.-N. Hua, H.-J. Yang, C.-M. Tseng, *Tetrahedron* **1997**, *53*, 17275.
- [10] For an example of a catalytic asymmetric [4+2] cycloaddition with tropones as 4 π components, see: a) P. Li, H. Yamamoto, *J. Am. Chem. Soc.* **2009**, *131*, 16628; For examples of non-asymmetric [4+2] cycloadditions with tropones as 4 π components, see: b) J. H. Rigby, J.-M. Sage, J. Raggon, *J. Org. Chem.* **1982**, *47*, 4815; c) R. L. Funk, G. L. Bolton, *J. Am. Chem. Soc.* **1986**, *108*, 4655; d) T. Asao, S. Itô, I. Murata, *Eur. J. Org. Chem.* **2004**, 899.
- [11] For examples of catalytic asymmetric [6+3] cycloadditions with tropones as 6 π components, see: a) B. M. Trost, P. J. McDougall, O. Hartmann, P. T. Wathen, *J. Am. Chem. Soc.* **2008**, *130*, 14960; b) B. M. Trost, P. J. McDougall, *Org. Lett.* **2009**, *11*, 3782; For an example of a non-asymmetric [6+3] cycloaddition with tropones as 6 π components, see: c) Y. Du, J. Feng, X. Lu, *Org. Lett.* **2005**, *7*, 1987.
- [12] For an example of a catalytic asymmetric [6+4] cycloaddition with tropones as 6 π components, see: a) J. H. Rigby, M. Fleming, *Tetrahedron Lett.* **2002**, *43*, 8643; for examples of non-asymmetric [6+4] cycloadditions with tropones as 6 π components, see: b) K. N. Houk, C. R. Watts, *Tetrahedron Lett.* **1970**, *11*, 4025; c) K. N. Houk, L. J. Luskus, N. S. Bhacca, *J. Am. Chem. Soc.* **1970**, *92*, 6392; d) D. Mukherjee, C. R. Watts, K. N. Houk, *J. Org. Chem.* **1978**, *43*, 817; e) J. H. Rigby, H. S. Ateeq, N. R. Charles, S. V. Cuisiat, M. D. Ferguson, J. A. Henshilwood, A. C. Krueger, C. O. Ogbu, K. M. Short, M. J. Heeg, *J. Am. Chem. Soc.* **1993**, *115*, 1382; f) L. Isakovic, J. A. Ashenhurst, J. L. Gleason, *Org. Lett.* **2001**, *3*, 4189; g) J. H. Rigby, G. Chourraqui, *Synlett* **2005**, 2501; h) J. A. Ashenhurst, L. Isakovic, J. L. Gleason, *Tetrahedron* **2010**, *66*, 368.
- [13] For examples of tropone as the 8 π component see: a) K. Kumar, A. Kapur, M. P. S. Ishar, *Org. Lett.* **2000**, *2*, 787; b) J. Okamoto, S. Yamabe, T. Minato, T. Hasegawa, T. Machiguchi, *Helv. Chim. Acta* **2005**, *88*, 1519; c) V. Nair, M. Poonoth, S. Vellalath, E. Suresh, R. Thirumalai, *J. Org. Chem.* **2006**, *71*, 8964.
- [14] For examples of tropothione as the 8 π component see: a) T. Machiguchi, T. Hasegawa, H. Otani, Y. Ishii, *J. Chem. Soc. Chem. Commun.* **1987**, 1375; b) T. Machiguchi, *Tetrahedron* **1995**, *51*, 1133; c) V. Nair, K. G. Abhilash, E. Suresh, *Tetrahedron Lett.* **2006**, *47*, 9329; d) A. R. Rivero, I. Fernández, M. A. Sierra, *J. Org. Chem.* **2012**, *77*, 6648.
- [15] For examples of azaheptafulvenes as 8 π components see: a) K. Yamamoto, S. Kajigaeshi, S. Kanemasa, *Chem. Lett.* **1977**, 85; b) K. Yamamoto, S. Kajigaeshi, S. Kanemasa, *Chem. Lett.* **1977**, 91; c) W. E. Truce, J. P. Shepherd, *J. Am. Chem. Soc.* **1977**, *99*, 6453; d) V. Nair, K. G. Abhilash, *Tetrahedron Lett.* **2006**, *47*, 8707; e) M. L. Lage, I. Fernández, M. A. Sierra, M. R. Torres, *Org. Lett.* **2011**, *13*, 2892.
- [16] a) K. Sanechika, S. Kajigaeshi, S. Kanemasa, *Chem. Lett.* **1977**, 861; b) T. Ishizu, K. Harano, M. Yasuda, K. Kanematsu, *J. Org. Chem.* **1981**, *46*, 3630; c) K. Hayakawa, H. Nishiyama, K. Kanematsu, *J. Org. Chem.* **1985**, *50*, 512; d) J. Barluenga, J. García-Rodríguez, S. Martínez, Á. L. Suárez-Sobrinho, M. Tomás, *Chem. Asian J.* **2008**, *3*, 767; e) J. Barluenga, J. García-Rodríguez, Á. L. Suárez-Sobrinho, M. Tomás, *Chem. Eur. J.* **2009**, *15*, 8800.
- [17] For selected examples using chiral *N,N'*-dioxide ligands, see: a) X. H. Liu, L. L. Lin, X. M. Feng, *Acc. Chem. Res.* **2011**, *44*, 574; b) A. Hassner, I. Namboothiri in *Organic Syntheses Based on Name Reactions*, 3th ed., Elsevier, Oxford, **2011**, p. 408; c) W. Li, X. H. Liu, X. Y. Hao, Y. F. Cai, L. L. Lin, X. M. Feng, *Angew. Chem.* **2012**, *124*, 8772; *Angew. Chem. Int. Ed.* **2012**, *51*, 8644; d) L. Zhou, X. H. Liu, J. Ji, Y. H. Zhang, X. L. Hu, L. L. Lin, X. M. Feng, *J. Am. Chem. Soc.* **2012**, *134*, 17023.
- [18] For other methods to construct the cyclohepta-fused pyrrole derivatives, see: a) L. E. Overman, E. J. Jacobsen, R. J. Doedens, *J. Org. Chem.* **1983**, *48*, 3393; b) X. Zhao, E. Zhang, Y.-Q. Tu, Y.-Q. Zhang, D.-Y. Yuan, K. Cao, C.-A. Fan, F.-M. Zhang, *Org. Lett.* **2009**, *11*, 4002.
- [19] a) X. Zhou, T. Xiao, Y. Iwama, Y. Qin, *Angew. Chem.* **2012**, *124*, 4993; *Angew. Chem. Int. Ed.* **2012**, *51*, 4909; b) N. Abe, T. Gunji, *Heterocycles* **2010**, *82*, 201; c) M. Nakatani, H. Fujii, T. Murafuji, T. Gunji, R. Ikeda, T. Konakahara, N. Abe, *Heterocycles* **2012**, *84*, 461.
- [20] CCDC 892341 (**3d**) and 892839 (**(±)-4h**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [21] The mechanism of the allylic rearrangement reaction was still unknown, for more information, see the Supporting Information.