

Heterocycles

Palladium-Catalyzed Hydrobenzylation of *ortho*-Tolyl Alkynyl Ethers by Benzylic C–H Activation: Remarkable Alkynoxy-Directing Effect**


Yasunori Minami,* Kotomi Yamada, and Tamejiro Hiyama*

Straightforward C(sp³)–H bond functionalization by transition-metal complexes allows novel synthetic manipulations of substrates without the requirement for prefunctionalization and thus contributes to environmentally benign transformations through efficient use of natural resources and energy.^[1] Notably, a process that involves C(sp³)–H bond activation by oxidative insertion and subsequent carbometalation of unsaturated substrates represents an ideal new carbon–carbon bond formation with 100 % atom economy.^[2]

As we have reported recently, alkynyl aryl ethers react with internal alkynes through selective palladium(0)-catalyzed *ortho* C–H activation, thus providing efficient access to 2-methylidene-2*H*-chromene derivatives.^[3] The facile and straightforward transformation can be attributed to the coordination of the alkynoxy directing group with palladium to effect the *ortho* C–H activation.^[4] On the basis of this preliminary observation, we were intrigued by the possibility of benzylic C–H activation. Although cross-coupling reactions involving benzylic C–H bond cleavage have been briefly explored to some extent,^[5–7] insertion of unsaturated partners is less studied. In this respect, the work by Yamamoto and co-workers^[8] and Chatani and co-workers^[9] are remarkable examples of M^{II}-catalyzed (M = Pt, Ru) benzylic C–H cleavage of 1-alkyl-2-ethynylbenzenes by vinylidene complexes. Herein, we report that a hydrobenzylation of *ortho*-tolyl alkynyl ethers takes place through palladium-catalyzed benzylic C(sp³)–H bond insertion and subsequent *syn*-1,2-addition across the alkyne. The adducts, 2-methylidene-2,3-dihydrobenzofurans, easily react with acetic acid, azo and carbonyl compounds, and molecular oxygen to give variously functionalized benzofurans.^[10,11]


The reaction of triisopropylsilyl-2,6-xylyl ether (**1a**) was conducted in the presence of Pd(OAc)₂ (1 mol %), tricyclohexylphosphine (2 mol %), and Zn metal (1 mol %, in situ Pd^{II} reduction) in toluene at 90 °C for 30 minutes to

provide 97 % of the *Z*-exocyclic alkylidene product **2a** along with 3 % of the benzofuran **3a** as determined by NMR spectroscopy [Eq. (1); TIPS = triisopropylsilyl]. The stereochemistry of **2a** was unambiguously determined by NOE and



NOESY experiments. Attempted isolation of **2a** by silica gel or alumina column chromatography resulted in its isomerization into **3a**, to some extent. Thus, treatment of the crude reaction mixture with acetic acid prior to work-up gave **3a** in 98 % yield upon isolation.^[12] Since Pd(OAc)₂, PdCl₂, and PtCl₂ failed to effect the desired reaction, it is apparent that the cyclization is not promoted by either the Lewis acid, PCy₃, or Zn(OAc)₂ alone.^[13] Of note, the reaction took place in the absence of zinc, albeit at a slower rate, thus indicating that the combination of Pd(OAc)₂, PCy₃, and zinc promotes the cyclization. As the carbon analogue, 4-MeOC₆H₄CH(OMe)C≡CTIPS, was inert to the standard reaction conditions, the alkynoxy moiety is definitely essential for the intramolecular hydrobenzylation reaction.

Other silyl groups instead of TIPS were used in the cyclization as depicted in Equation (2). Substrates containing



tert-butyldiphenylsilyl (TBDPS), *tert*-butyldimethylsilyl (TBDMS), and triethylsilyl (TES) groups on the terminal ethynyl carbon atom underwent the hydrobenzylation to produce **2b**, **2c**, and **2d**, respectively, in quantitative yields as estimated by NMR assays. Isomerization by acetic acid led to the isolation of the corresponding **3b**, **3c**, and **3d** in essentially quantitative yields. In contrast, the dimethylphenylsilyl-substituted ethynyl ether **1e** was gave **2e** in 60 % yield only by

[*] Y. Minami, T. Hiyama


Research and Development Initiative, Chuo University, JST, ACT-C Kasuga, Bunkyo-ku, Tokyo 112-8551 (Japan)
E-mail: yminami@kc.chuo-u.ac.jp
thiyama@kc.chuo-u.ac.jp

K. Yamada
Graduate School of Science and Engineering, Chuo University Kasuga, Bunkyo-ku, Tokyo 112-8551 (Japan)


[**] This work has been supported financially by a Grant-in-Aid for Scientific Research (S) (21225005 to T.H.) and Young Scientists (B) (25870747 to Y.M.) from JSPS.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/ange.201304893>.

functionality. Protodesilylation of **3a** with TBAF gave 2,7-dimethyl benzofuran in a quantitative yield, and thus the silyl moiety behaves as a latent nucleophile as is shown by the TBAF-mediated 1,2-addition of **3e** to benzaldehyde, thereby leading to the corresponding alcohol **9** being isolated in 62 % yield [Eq. (3); TBAF = tetra-*n*-butylammonium fluoride].





The Pd/Zn-catalyzed intramolecular hydrobenzylation reaction is applicable to the aryl ethynyl ethers **10** bearing a carbonaceous substituent instead of a silyl group [Eq. (4)]. The cyclization smoothly took place to give the corresponding



products **11**, which upon treatment with acetic acid gave the 2-alkenyl-substituted benzofurans **12** by elimination of methanol.^[15] For example, cyclization of the 2,6-xylyl 2-(1-methoxycyclohexyl)ethynyl ethers **10a** ($R = R' = -(CH_2)_5-$), **10b** ($R = R' = Ph$), and **10c** ($R = Ph$, $R' = H$) took place in the presence of 5 mol % of palladium catalyst to form the corresponding products (**11**), which readily converted into the 2-alkenyl benzofurans **12** in moderate yields over two steps. In the case of **10c**, the β -styryl-substituted benzofuran **12c** was obtained as a single isomer. However, the reaction of a less bulky substrate, the 3-methoxypropynyl ether **10d**, turned out futile. These results clearly demonstrate that the presence of a bulky substituent on the propargylic position is essential for the success of the hydrobenzylation reaction.

To gain insight into the mechanism of the present hydrobenzylation, we conducted deuterium-labeling experiments. The reaction of $[D_6]$ -**1a** in toluene resulted in the formation of $[D_6]$ -**2a** with 97 % deuterium incorporation at the vinyl carbon atom as was evaluated by 1H NMR analysis [Eq. (5)]. This result demonstrates that the deuterium located originally on the benzylic carbon atom is transposed cleanly to the 2-methylidene carbon atom.

The kinetic isotope effect (KIE) was subsequently investigated (Scheme 2). The relative rate, k_H/k_D , of the reaction of




Scheme 2. Kinetic isotope effect experiments.

1a versus $[D_6]$ -**1a** was observed under the catalysis with 5 mol % of $Pd(OAc)_2/2PCy_3/Zn$ in C_6D_6 for 2 hours at $50^\circ C$, and the ratio was determined to be 12.7 (Scheme 2a), thus indicating that the C–H bond is probably cleaved by a concerted metalation/deprotonation (CMD) transition state (TS).^[16] The substrate $[D_3]$ -**1a**, containing both CH_3 and CD_3 groups, was exposed to the hydrobenzylation conditions ($50^\circ C$, 14 h in C_6D_6) and the resultant KIE value was found to be 6.9, although an H/D exchange occurred to some extent at the vinyl carbon atom α to the silyl group (Scheme 2b). Accordingly, the most probable rate-determining step may be attributed to the C–H bond cleavage.

The effect of $Zn(OAc)_2$, generated in situ from the reaction of $Pd(OAc)_2$ with Zn, is obvious. The $[Pd(dba)_2]/PCy_3$ -catalyzed reaction of **1a** in the presence or absence of $Zn(OAc)_2$ was monitored at $60^\circ C$. The hydrobenzylation took place in the presence of $Zn(OAc)_2$; without $Zn(OAc)_2$ **1a** was recovered almost unchanged. To clarify the role of $Zn(OAc)_2$, we examined other additives such as $ZnCl_2$ (20 mol %), $Zn(OTf)_2$ (20 mol %), NBu_4OAc (2 mol %), or $AcOH$ (2 mol %) for the cyclization of **1a** in the presence of $[Pd(dba)_2]$ (1 mol %) and PCy_3 (2 mol %), and found that only $AcOH$ only promoted the reaction and gave **2a** quantitatively. In contrast, the less bulky TES-substituted **1d** cyclized to form the isomerized product **3d** in 45 % in the presence of NBu_4OAc (2 mol %), thus indicating that the generated $Zn(OAc)_2$ is not merely for the acetate ion donor.^[17] These observations combined with the previous results indicate that in situ generated $Zn(OAc)_2$ acts as a Lewis acid and acetate ion donor to promote the cleavage of benzylic C–H bond.

Although more information is necessary to discuss the reaction mechanism in detail, we propose a mechanism for the $Pd(OAc)_2/PCy_3/Zn$ -catalyzed hydrobenzylation as summarized in Scheme 3 with **1i** as a representative substrate. First, the alkyne-coordinated palladium(0) complex **13** is formed, wherein the intramolecular nucleophilic attack of electron-rich palladium(0) on the α -carbon atom of the alkynoxy group presumably occurs to give the complex **14**. The alkynoxy group is reasonably assumed to have a ketene-like polarized resonance structure, thus indicating that the α -carbon atom bound to the oxygen center is electrophilic and the β -carbon atom nucleophilic.^[18] Thus, the β -carbanionic center in **14** is neutralized by $Zn(OAc)_2$ to give the vinyl



Scheme 3. Proposed reaction mechanism.

palladium acetate **15** by the migration of the generated acetate ion to the palladium center. An alternative pathway from **13** to **15** may be also possible: ligation of the β -carbon atom of **13** to $\text{Zn}(\text{OAc})_2$ could give **16**, whose palladium ligated by acetoxy group on Zn attacks the α -carbon atom to give **15**. The benzylic C–H bond is cleaved by CMD to give the acetic-acid-coordinated palladacycle intermediate **17**,^[19,20] with subsequent rapid protonation of the vinyl zinc moiety by acetic acid and reductive elimination to the *cis*-adduct **2i** and regeneration of the palladium(0) complex and $\text{Zn}(\text{OAc})_2$.^[21] It is evident that the substituents on the ethynyl carbon atom are crucial for the success of the cyclization. Bulky substituents on the terminal ethyne carbon atom apparently push the η^2 -palladium(0) complex to the α -carbon atom owing to steric repulsion. Moreover, the silyl group (e.g., TIPS) stabilizes the negative charge at the silicon-bearing carbon atom (α effect) to promote the cyclization.

In conclusion, the present study demonstrates that the palladium-catalyzed hydrobenzylation of *ortho*-tolyl alkynyl ethers provides efficient access to reactive 2-methylidene-2,3-dihydrobenzofurans by selective benzylic $\text{C}(\text{sp}^3)\text{--H}$ activation. The alkynoxy moiety containing bulky substituents as a directing group plays a key role in the present transformation. It is worthy of note that $\text{Zn}(\text{OAc})_2$ promotes the cyclization. The proper choice of the substituents bound to the terminal ethynyl carbon atom allows additional transformations. Efforts are currently directed toward further understanding of the reaction mechanism and synthesis of various heterocycles useful for organic light-emitting materials.

Received: June 7, 2013

Revised: July 5, 2013

Published online: August 16, 2013

Keywords: alkynes · arenes · heterocycles · palladium · synthetic methods

- [1] For recent reviews, see: a) F. Kakiuchi, T. Kochi, *Synthesis* **2008**, 3013; b) O. Daugulis, H. Q. Do, D. Shabashov, *Acc. Chem. Res.* **2009**, *42*, 1074; c) X. Chen, K. M. Engle, D.-H. Wang, J.-Q. Yu, *Angew. Chem.* **2009**, *121*, 5196; *Angew. Chem. Int. Ed.* **2009**, *48*, 5094; d) R. Giri, B.-F. Shi, K. M. Engle, N. Maugel, J.-Q. Yu, *Chem. Soc. Rev.* **2009**, *38*, 3242; e) T. W. Lyons, M. S. Sanford, *Chem. Rev.* **2009**, *109*, 1147; f) G. P. Chiusoli, M. Catellani, M.

- Costa, E. Motti, N. Della Ca', G. Maestri, *Coord. Chem. Rev.* **2010**, *254*, 456; g) T. W. Lyons, M. S. Sanford, *Chem. Rev.* **2010**, *110*, 1147; h) R. Jazzar, J. Hitce, A. Renaudat, J. Sofack-Kreutzer, O. Baudoin, *Chem. Eur. J.* **2010**, *16*, 2654; i) J.-Q. Yu, M. Shi, *C-H Activation*, Springer, Heidelberg, **2010**; j) O. Baudoin, *Chem. Soc. Rev.* **2011**, *40*, 4902; k) C. L. Sun, B. J. Li, Z. J. Shi, *Chem. Rev.* **2011**, *111*, 1293.

- [2] B. M. Trost, *Science* **1991**, *254*, 1471.
[3] Y. Minami, Y. Shiraishi, K. Yamada, T. Hiyama, *J. Am. Chem. Soc.* **2012**, *134*, 6124.
[4] For examples of C–C multiple bonds as a directing group, see: a) R. Takeuchi, H. Yasue, *J. Org. Chem.* **1993**, *58*, 5386; b) K. Tanaka, G. C. Fu, *Org. Lett.* **2002**, *4*, 933; c) M. Murakami, T. Itahashi, Y. Ito, *J. Am. Chem. Soc.* **2002**, *124*, 13976; d) K. Takeishi, K. Sugishima, K. Sasaki, K. Tanaka, *Chem. Eur. J.* **2004**, *10*, 5681; e) N. Chernyak, V. Gevorgyan, *J. Am. Chem. Soc.* **2008**, *130*, 5636; f) M. Tobisu, I. Hyodo, M. Onoe, N. Chatani, *Chem. Commun.* **2008**, 6013; g) N. Chernyak, V. Gevorgyan, *Adv. Synth. Catal.* **2009**, *351*, 1101; h) N. Chernyak, S. I. Gorelsky, V. Gevorgyan, *Angew. Chem.* **2011**, *123*, 2390; *Angew. Chem. Int. Ed.* **2011**, *50*, 2342; i) P. Gandeepan, C.-H. Chen, *J. Am. Chem. Soc.* **2012**, *134*, 5738; j) T. Xu, G. Dong, *Angew. Chem.* **2012**, *124*, 7685; *Angew. Chem. Int. Ed.* **2012**, *51*, 7567.
[5] For directing-group-assisted benzylic C–H activation, see: a) C.-H. Jun, D.-C. Hwang, S.-J. Na, *Chem. Commun.* **1998**, 1405; b) F. Kakiuchi, K. Tsuchiya, M. Matusmoto, E. Mezushima, N. Chatani, *J. Am. Chem. Soc.* **2004**, *126*, 12792.
[6] For recent examples of benzyl C–H functionalization, see: a) T. Xiong, Y. Li, X. Bi, Y. Lv, Q. Zhang, *Angew. Chem.* **2011**, *123*, 7278; *Angew. Chem. Int. Ed.* **2011**, *50*, 7140; b) M. Ichinose, H. Suematsu, Y. Yasutomi, Y. Nishioka, T. Uchida, T. Katsuki, *Angew. Chem.* **2011**, *123*, 10058; *Angew. Chem. Int. Ed.* **2011**, *50*, 9884; c) H. J. Kim, J. Kim, S. H. Cho, S. Chang, *J. Am. Chem. Soc.* **2011**, *133*, 16382; d) J. Zhang, C. Stanciu, B. Wang, M. M. Hussain, C.-S. Da, P. J. Carroll, S. D. Dreher, P. J. Walsh, *J. Am. Chem. Soc.* **2011**, *133*, 20552; e) Z. Ni, Q. Zhang, T. Xiong, Y. Zheng, Y. Li, H. Zhang, J. Zhang, Q. Liu, *Angew. Chem.* **2012**, *124*, 1270; *Angew. Chem. Int. Ed.* **2012**, *51*, 1244; f) J.-J. Chen, S. Onogi, Y.-C. Hsieh, C.-C. Hsiao, S. Higashibayashi, H. Sakurai, Y.-T. Wu, *Adv. Synth. Catal.* **2012**, *354*, 1551; g) P. Xie, Y. Xie, B. Qian, H. Zhou, C. Xia, H. Huang, *J. Am. Chem. Soc.* **2012**, *134*, 9902; h) J. Zhang, A. Bellomo, A. D. Creamer, S. D. Dreher, P. J. Walsh, *J. Am. Chem. Soc.* **2012**, *134*, 13765.
[7] For recent examples of aza-aryl methyl C–H functionalization, see: a) H. Tsurugi, K. Yamamoto, K. Mashima, *J. Am. Chem. Soc.* **2011**, *133*, 732; b) P. M. Burton, J. A. Morris, *Org. Lett.* **2010**, *12*, 5359; c) H. Komai, T. Yoshino, S. Matsunaga, M. Kanai, *Org. Lett.* **2011**, *13*, 1706; d) G. Song, Y. Su, X. Gong, K. Han, X. Li, *Org. Lett.* **2011**, *13*, 1968; e) H. Komai, T. Yoshino, S. Matsunaga, M. Kanai, *Synthesis* **2012**, 2185; f) S. Duez, A. K. Steib, S. M. Manolikakes, P. Knochel, *Angew. Chem.* **2011**, *123*, 7828; *Angew. Chem. Int. Ed.* **2011**, *50*, 7686.
[8] G. B. Bajracharya, N. K. Pahadi, I. D. Gridnev, Y. Yamamoto, *J. Org. Chem.* **2006**, *71*, 6204.
[9] M. Tobisu, H. Nakai, N. Chatani, *J. Org. Chem.* **2009**, *74*, 5471.
[10] Tanaka and co-workers reported the synthesis of dibenzofurans by rhodium-catalyzed [2+2+2] cycloaddition of *ortho*-ethynyl alkynyl aryl ethers with alkynes or nitriles. Y. Komine, A. Kamisawa, K. Tanaka, *Org. Lett.* **2009**, *11*, 2361.
[11] Benzofurans represent an important structural motif. For selected reviews on bioactive natural products and medicinal chemistry, see: a) G. W. Gribble in *Comprehensive Heterocyclic Chemistry II*, Vol. 2 (Eds.: A. R. Katritzky, C. W. Rees, E. F. V. Scriven), Pergamon, Oxford, **1996**, pp. 207–257; b) X.-L. Hou, Z. Yang, H. N. C. Wong in *Progress in Heterocyclic Chemistry*, Vol. 14 (Eds.: G. W. Gribble, T. L. Gilchrist), Pergamon, Oxford, **2002**, pp. 139–179; c) T. Kawasaki, K. Higuchi, *Nat. Prod. Rep.*

- 2007, 24, 843. For selected examples in material chemistry, see: d) H. Tsuji, C. Mitsui, L. Ilies, Y. Sato, E. Nakamura, *J. Am. Chem. Soc.* **2007**, 129, 11902; e) H. Tsuji, C. Mitsui, Y. Sato, E. Nakamura, *Adv. Mater.* **2009**, 21, 3776; f) A. Muranaka, S. Yasuike, C.-Y. Liu, J. Kurita, N. Kakusawa, T. Tsuchiya, M. Okuda, N. Kobayashi, Y. Matsumoto, K. Yoshida, D. Hashizume, M. Uchiyama, *J. Phys. Chem. A* **2009**, 113, 464; g) C. Mitsui, J. Soeda, K. Miwa, H. Tsuji, J. Takeya, E. Nakamura, *J. Am. Chem. Soc.* **2012**, 134, 5448.
- [12] a) J. D. Brewer, J. A. Elix, *Aust. J. Chem.* **1975**, 28, 1059. For 1,3-migration of hydroxy group, see: b) B. Gabriele, R. Mancuso, G. Salerno, *J. Org. Chem.* **2008**, 73, 7336.
- [13] For reactions of alkynyl ethers with Lewis acids, see: a) L. Zhang, S. A. Kozmin, *J. Am. Chem. Soc.* **2004**, 126, 11806; b) A. S. K. Hashmi, M. Rudolph, J. W. Bats, W. Frey, F. Rominger, T. Oeser, *Chem. Eur. J.* **2008**, 14, 6672; c) A. S. K. Hashmi, M. Rudolph, J. Huck, W. Frey, J. W. Bats, M. Hamzić, *Angew. Chem.* **2009**, 121, 5962; *Angew. Chem. Int. Ed.* **2009**, 48, 5848.
- [14] The reaction in 1.0M concentration gave a mixture of **3i** and a dimeric product, by *ortho* C–H activation, in 31 and 55% yield, respectively.
- [15] L. Yu Ukhin, L. V. Belousova, Z. I. Orlova, M. S. Korobov, G. S. Borodkin, *Chem. Heterocycl. Compd.* **2002**, 38, 1174.
- [16] N. Guimond, S. I. Gorelsky, K. Fagnou, *J. Am. Chem. Soc.* **2011**, 133, 6449.
- [17] It is assumed that the steric repulsion between the TIPS group and NBU₄OAc interferes with the interaction of a palladium center with the acetate ion to prevent the cyclization.
- [18] a) R. Tanaka, S. I. Miller, *Tetrahedron Lett.* **1971**, 12, 1753; b) P. J. Stang, K. A. Roberts, *J. Am. Chem. Soc.* **1986**, 108, 7125; c) M. Shindo, *Tetrahedron* **2007**, 63, 10.
- [19] a) H.-Y. Aun, S. I. Gorelsky, D. R. Stuart, L.-C. Campeau, K. Fagnou, *J. Org. Chem.* **2010**, 75, 8180; b) S. Rousseaux, S. I. Gorelsky, B. K. W. Chung, K. Fagnou, *J. Am. Chem. Soc.* **2010**, 132, 10692.
- [20] Selective benzylic C–H cleavage rather than *ortho* C–H bond by intramolecular imino-coordinated bromoplatinum(II) complex was reported. See: M. Crespo, C. M. Anderson, N. Kfoury, M. Font-Bardia, T. Calvet, *Organometallics* **2012**, 31, 4401.
- [21] The cyclization proceeded in the absence of Zn under the standard reaction conditions. In this case, it is assumed that the [Pd(PCy₃)(OAc)][−] complex is the active catalyst which attacks the α -carbon atom of the alkynoxy group to give a vinyl palladium acetate. See: a) C. Amatore, A. Jutand, M. A. M'Barkl, *Organometallics* **1992**, 11, 3009; b) C. Amatore, E. Carré, A. Jutand, M. A. M'Barkl, *Organometallics* **1995**, 14, 1818.