


Harnessing Reversible Oxidative Addition: Application of Diiodinated Aromatic Compounds in the Carboiodination Process**

David A. Petrone, Matthias Lischka, and Mark Lautens*

Transition-metal-mediated couplings of halogenated aromatics have been extensively studied since the 1960s.^[1] Notwithstanding the success of recent aromatic functionalization strategies such as directed *ortho* metalation^[2] and C–H functionalization,^[3–5] Pd-catalyzed aromatic carbon–halogen bond functionalization remains a central strategy. In this regard, the application of polyhalogenated aromatic compounds in site-selective transformations is one of the remaining challenges in this field,^[6] due to overcoupling^[6h] and chemoselectivity issues.^[7] Methods that take advantage of the intrinsic steric and electronic differences between different carbon–halogen bonds have been developed.^[6a,i] However, most of these strategies remain limited, requiring difficult substrate prefunctionalization to enforce the desired selectivity. Under traditional catalytic conditions, irreversible oxidative addition to a carbon–halogen bond of **A** occurs to give **B** (Scheme 1a), which lacks a productive reaction pathway. The presence of intermolecular Heck acceptors, or the addition of nucleophiles, is a strategy used to promote a catalytic cycle, as it allows catalytic dead ends to be avoided while increasing product complexity.^[6b,f]

A more general and attractive solution would be to use catalysts capable of undergoing reversible oxidative addition.^[8] Building on the stoichiometric experiments on reductive elimination from $\text{ArPd}^{\text{II}}\text{X}$ complexes conducted by Hartwig et al.,^[9] and the contributions of Buchwald et al., which afford aromatic C–F^[10a] and C–Br bonds;^[10b] our group has developed Pd⁰-catalyzed transformations exhibiting reversible oxidative addition as a key to catalysis.^[8,11] To this end, the application of our carboiodination method to diiodinated substrates would highlight the unique capabilities of the Pd/QPhos combination: the ability to oxidatively add reversibly to carbon–halogen bonds, and to promote sp^3 carbon–iodine reductive elimination. Herein, we report both



Scheme 1. a) Catalyst deactivation by irreversible oxidative addition; b) Strategy to overcome catalyst deactivation (when R ≠ H). QPhos = 1,2,3,4,5-pentaphenyl-1'-[(di-*tert*-butylphosphino)ferrocene].

the carboiodination, and the sequential intramolecular carboiodination/intermolecular Heck reaction of diiodinated aromatic substrates (Scheme 1b).

We began by optimizing the intramolecular carboiodination of **1a** (Table 1). Although 5 mol % of [Pd(PrBu₃)₂] in toluene at 100 °C led to full conversion, the desired product (**2a**) was obtained in only 30 % isolated yield after 18 h.^[12] The yield of **2a** depended on the Pd⁰ precatalyst used, as well as the presence of both additional QPhos and base.^[13] Based on previous synthetic reports^[8,11] and computational evidence,^[14a] we believe that the steric bulk of QPhos make it ideal for promoting carbon–iodine reductive elimination.^[14b] The optimized conditions were found to be [Pd(QPhos)₂] (**Pd-1**; 5 mol %), additional QPhos (10 mol %), and 1,2,2,6,6-pentamethylpiperidine (PMP; 2 equiv) in toluene at 110 °C.^[15] Under these conditions, **2a** was isolated in 74 % yield. With the optimized reaction conditions in hand, we examined a series of diiodinated compounds, **1b–1f**.


Varying the location of the halogen substituent did not affect the transformation as **1b** could be efficiently cyclized to **2b** in 76 % yield (Table 1, entry 2). Six-membered rings could be accessed, as **1c** cyclized to afford chroman **2c** in 74 % yield (entry 3). Dihydrobenzofuran **2d** and oxindole **2f** were isolated in 60 % and 68 % yield when the reaction times

[*] D. A. Petrone, M. Lischka, Prof. Dr. M. Lautens
Department of Chemistry, University of Toronto
80 St. George St. Toronto, Ontario (Canada)
E-mail: mlautens@chem.utoronto.ca
Homepage: <http://www.chem.utoronto.ca/staff/ML/index.php>

[**] We thank the Natural Sciences and Engineering Research Council of Canada (NSERC) and the University of Toronto for financial support. NSERC/Merck-Frosst Canada is thanked for an Industrial Research Chair. D.A.P. thanks NSERC for a postgraduate scholarship (CGS-D). The authors wish to acknowledge CFI project number 19119 for funding the CSICOMP NMR Facility. Dr. Timothy Burrow is sincerely thanked for his dedicated assistance with NMR. We thank Johnson Matthey for kind gifts of QPhos, [Pd(QPhos)₂], and [Pd(crotlyl)QPhosCl].

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/ange.201304923>.

Table 1: Scope of the intramolecular carboiodination of diiodoarenes.^[a]




Entry	Substrate 1	Product 2	t [h]	Yield [%] ^[b]
1			5	74 (67) ^[c]
2			7	76
3			8	74
4			20	60
5			11	58
6			30	68

[a] Reactions conditions: diiodoarene (0.1–0.2 mmol), [Pd(QPhos)₂], (5 mol %), QPhos (10 mol %), PMP (2 equiv), PhMe, 110 °C. [b] Yield of isolated product. [c] Reaction run on 2.5 mmol scale.

were increased to 20 h and 30 h, respectively (entries 4 and 6). Substrate **1e** contains iodides on different aryl rings, but gave **2e** in 58 % yield after 11 h (entry 5).

We next sought to apply diiodinated compounds in an orthogonal intramolecular carboiodination/intermolecular Heck reaction sequence. Among the challenges are (Scheme 2): 1) starting material homo-dimerization; 2) undesired Heck reactions occurring either prior to or after the desired intermolecular Heck reaction (path a); 3) reactive



Scheme 2. Challenges of the reaction sequence (R = CO₂tBu).

Table 2: Optimization of the orthogonal intramolecular carboiodination/intermolecular Heck reaction sequence.^[a]

Reaction scheme showing the synthesis of compounds **2a**, **3a**, and **4a** from substrate **1a** and **tBuO₂C-CH=CH₂** using **Pd-1** or **Pd-2** (5 mol%), **QPhos** (*x* mol%), and **NR₃** (*y* equiv) in **PhMe** (0.1 M) at 100 °C for 24 h.

Entry	Pd Source	QPhos (<i>x</i> mol %)	NR ₃ (<i>y</i> equiv)	Yield [%] ^[b]	2a	3a	4a
1	Pd-1	–	NEt ₃ (2)	16	16	40	
2	Pd-1	–	NEt ₃ (4)	18	18	21	
3 ^[c]	Pd-1	–	NEt ₃ (4)	8	5	55	
4	Pd-1	10	NEt ₃ (2)	10	7	57	
5 ^[c]	Pd-1	10	NEt ₃ (2)	14	8	57	
6 ^[c]	Pd-1	10	PMP (2)	4	–	86	
7 ^[c]	Pd-1	5	PMP (2)	5	16	62	
8 ^[c,d]	Pd-2	10	PMP (2)	–	–	86	
9 ^[c–f]	Pd-2	8	PMP (2)	–	–	88 ^[g] (80) ^[h]	


[a] 0.2 mmol scale, 0.1 M in toluene. [b] Yields determined by ¹H NMR analysis of the crude reaction mixture using 1,3,5-trimethoxybenzene as an internal standard. [c] TBA (1.5 equiv). [d] KOtBu (20 mol %) used. [e] **Pd-2** (4 mol %). [f] **1a** (0.125 M). [g] Yield of isolated product. [h] Reaction run on a 2.5 mmol scale.

neopentyl Pd^{II} intermediate^[16] **D** could undergo an intermolecular Heck reaction (path b) instead of reductive elimination to form **4a** (path c).

We began by optimizing the reaction between **1a** and *tert*-butyl acrylate (TBA; Table 2). With **Pd-1** (5 mol %) and NEt₃ (2 equiv) in toluene at 100 °C for 24 h, **4a** was obtained as the major product in 40 % yield along with 16 % of both **2a** and **3a** (entry 1). The reaction was completely chemoselective, as no undesired Heck products were observed. Increasing the amount of NEt₃ to 4 equiv led to a decrease in the yield of **4a** to 21 % (entry 2). However, increasing the amount of olefin to 1.5 equiv, or adding an additional 10 mol % of QPhos increased the yield of **4a** to 55 % and 57 %, respectively, while simultaneously decreasing the amount of **2a** and **3a** (entries 3 and 4). When PMP was used, **4a** was furnished in 86 % yield, with only 4 % of **2a** and no trace of **3a** (entry 6). The complex [Pd(crotyl)QPhosCl] (**Pd-2**) was an equally effective Pd⁰ precursor,^[17] and employing 5 mol %, while maintaining all established conditions, gave rise to **4a** in 86 % yield (entry 8). Using this catalyst is ideal, as it is air stable and allows the ligand loading to be decreased. The final optimized conditions were TBA (1.5 equiv), **Pd-2** (4 mol %), QPhos (8 mol %), and PMP (2 equiv) in toluene at 100 °C for 24 h, where **4a** was isolated in 88 % yield with no observed **2a** or **3a** (entry 9).


Control experiments were then conducted to determine if all proposed intermediate products could ultimately convert into the desired adducts (Scheme 3). We found that both **2a** and **3a** could be converted into **4a** in 83 % and 91 % yield, respectively. These experiments suggest that either intermediate could proceed towards the desired product, and that both pathways could be occurring simultaneously.

We tested this proposal using ¹H NMR spectroscopy to monitor the reaction between **1a** and TBA in situ



Scheme 3. Control experiments concerning reaction intermediates. Conditions: **Pd-2** (4 mol%), QPhos (8 mol%), KOtBu (20 mol%), PMP (2 equiv), toluene, 100 °C, 12 h. R = CO₂tBu.

(Scheme 4).^[18] We observed similar increases in the concentration of both products **2a** and **3a**, from the intramolecular carboiodination and the intermolecular Heck reaction, respectively. In contrast to Table 1 where selective *ortho*




Scheme 4. In situ monitoring of the reaction of **1a** and TBA. Conditions: **Pd-2** (0.04 equiv), QPhos (0.08 equiv), KOtBu (0.2 equiv), **1a** (0.087 M), TBA (0.122 M), PMP (0.216 M), [D₈]toluene, 373 K, 700 MHz.

coupling occurs, this experiment strongly suggests that the Pd⁰ catalyst can oxidatively add reversibly into both carbon–iodine bonds of **1a**, allowing both reactions to proceed simultaneously. Additionally, there are similar decreases in the concentrations of **2a** and **3a**, thus highlighting that both intermediates can converge to **4a**.

Finally, the optimized conditions were applied to a series of Heck acceptors and diiodoarenes (Table 3). When the reaction time was doubled to 48 h, the loadings of **Pd-2** and QPhos could be decreased to 2 mol% and 4 mol%, respectively, when the reaction was run on a 2.5 mmol scale. This combination allowed the isolation of **4a** in 79% yield (entry 1). Methyl acrylate and *N,N*-dimethyl acrylamide were well tolerated as acceptors, and their adducts, **4b** and **4c**, were isolated in 83% and 73% yield, respectively (entries 2 and 3). Subjecting styrene and acrylonitrile to the standard conditions delivered the desired products **4d** and **4e** in 82% and 90% yield, respectively (entries 4 and 5). Methyl methacrylate reacted selectively to form **4f** in 77% yield (entry 6) in the presence of 5 equiv of olefin. *N*-tosyl indolines could also be obtained as **4g** and **4h** were afforded in 86% and 84% (entries 8 and 9). Electron-deficient olefins were tolerated in the intramolecular carboiodination component,

Table 3: Reaction scope.^[a]

			
Entry	Product	4	Yield [%] ^[b]
1	R = CO ₂ tBu	4a	88 (79) ^[c]
2	R = CO ₂ Me	4b	83
3 ^[c]	R = CONMe ₂	4c	73
4	R = Ph	4d	82
5	R = CN	4e	90 ^[d]
		(4:1 <i>E/Z</i>)	
6		4f (R = Me)	77
7		4f' (R = Ph)	25
8 ^[e]	R = CO ₂ tBu	4g	86
9	R = 2-py	4h	84
10 ^[e]		4i	87
11		4j ^[f] (d.r.=87:13)	86
12	R = CO ₂ tBu	4k	91
13	R = CONMe ₂	4l	88
14	R = 2-py	4m	74
15 ^[e]		4n	70

[a] 0.2 mmol scale. [b] Yield of isolated product. [c] **1a** (2.5 mmol, 0.25 M), **Pd-2** (2 mol%), QPhos (4 mol%), KOtBu (10 mol%), toluene, 48 h. [d] Combined yield of both isomers. [e] Reaction heated for 48 h. [f] Stereochemistry assigned based on analogy to a previous report, see Ref. [11c]. 2-py = 2-pyridyl.

as *N*-methyl oxindole **4i** could be synthesized in 87% yield (entry 10). Switching to a *para*-iodo substitution pattern had no effect (entries 12–14). A nitrogen-containing polycycle **4n** could also be obtained in 70% yield (entry 15).

In conclusion, we have shown the utility of diiodinated aromatic compounds in a Pd⁰-catalyzed intramolecular carboiodination. These two complementary reactions allow access to a diverse range of cross-coupling substrates and olefin containing heterocycles. Commonly encountered problems associated with this important class of compounds are effectively avoided by carefully selecting a ligand that allows reductive elimination from unproductive catalytic intermediates.

Experimental Section

KOrBu (4.5 mg, 0.04 mmol, 20 mol %), [Pd(crotyl)QPhosCl] (7.3 mg, 0.008 mmol, 4 mol %), and QPhos (11.4 mg, 0.016 mmol, 8 mol %) were added to a dry screw cap vial equipped with a stir bar, which was purged with dry argon. The contents were taken up in toluene (0.8 mL) and stirred at room temperature for 15 min. A toluene solution (0.8 mL) of **1a** (79.9 mg, 0.2 mmol, 1 equiv), *tert*-butyl acrylate (44 μ L, 0.3 mmol, 1.5 equiv), and PMP (72 μ L, 0.4 mmol, 2 equiv) was then added to the pre-mixed catalyst solution. The vial was sealed and placed into a preheated oil bath (100°C). After 24 h, the reaction was filtered through silica, and concentrated. The crude product was purified by column chromatography using hexanes/EtOAc (100:0–100:3 v/v) to give a clear, colourless oil (70.4 mg, 0.176 mmol, 88 %).

Received: June 7, 2013

Revised: July 11, 2013

Published online: August 12, 2013

Keywords: carboiodination · Heck reaction · heterocycles · oxidative addition · palladium

- [1] a) E. Negishi in *Handbook of Organopalladium Chemistry for Organic Synthesis*, Vol. 2, Wiley, New York, **2002**; b) E. Negishi, *Acc. Chem. Res.* **1982**, *15*, 340; c) N. Miyaura, A. Suzuki, *Chem. Rev.* **1995**, *95*, 2457; d) J. Hassan, M. Sévignon, C. Gozzi, E. Schulz, M. Lemaire, *Chem. Rev.* **2002**, *102*, 1359; e) A. F. Littke, G. C. Fu, *Angew. Chem.* **2002**, *114*, 4350; *Angew. Chem. Int. Ed.* **2002**, *41*, 4176; f) L. F. Tietze, H. Ila, H. P. Bell, *Chem. Rev.* **2004**, *104*, 3453; g) J. Dupont, C. S. Consorti, J. Spencer, *Chem. Rev.* **2005**, *105*, 2527; h) G. Zeni, R. C. Larock, *Chem. Rev.* **2006**, *106*, 4644.
- [2] M. C. Whisler, S. MacNeil, V. Snieckus, P. Beak, *Angew. Chem.* **2004**, *116*, 2256; *Angew. Chem. Int. Ed.* **2004**, *43*, 2206.
- [3] a) T. W. Lyons, M. S. Sanford, *Chem. Rev.* **2010**, *110*, 1147; b) G. Song, F. Wang, X. Li, *Chem. Soc. Rev.* **2012**, *41*, 3651; c) D. A. Colby, R. G. Bergman, J. A. Ellman, *Chem. Rev.* **2010**, *110*, 624; d) P. B. Arockian, C. Bruneau, P. H. Dixneuf, *Chem. Rev.* **2012**, *112*, 5879.
- [4] D. Alberico, M. E. Scott, M. Lautens, *Chem. Rev.* **2007**, *107*, 174.
- [5] S. Rousseaux, B. Liégault, K. Fagnou in *C–H Functionalization: A New Strategy for the Synthesis of Biologically Active Natural Products* (Eds.: J. Cossy, S. Arseniyadis), Wiley, Hoboken, **2012**, pp. 1–32.
- [6] a) For a review, see: S. Schröter, C. Stock, T. Bach, *Tetrahedron* **2005**, *61*, 2245, and references therein; b) M. Lautens, Y.-Q. Fang, *Org. Lett.* **2003**, *5*, 3679; c) K. T. J. Loones, B. U. W. Maes, C. Meyers, J. Deruytter, *J. Org. Chem.* **2006**, *71*, 260; d) D. Conreux, E. Bossharth, N. Monteiro, P. Desbordes, J.-P. Vors, G. Balme, *Org. Lett.* **2007**, *9*, 271; e) A. K. Yadav, S. Verbeeck, S. Hostyn, P. Franck, S. Sergeyev, B. U. W. Maes, *Org. Lett.* **2013**, *15*, 1060; f) T. Saget, D. Perez, N. Cramer, *Org. Lett.* **2013**, *15*, 1354; g) A. M. Hamdy, N. Eleya, H. H. Mohammed, T. Patonay, A. Spinnenberg, P. Langer, *Tetrahedron* **2013**, *69*, 2081; h) S. Tanaka, G. Tatsuta, A. Sugie, A. Mori, *Tetrahedron Lett.* **2013**, *54*, 1976; i) Z. Hassan, T. Patonay, P. Langer, *Synlett* **2013**, 412.
- [7] J.-R. Wang, K. Manabe, *Synthesis* **2009**, *9*, 1405.
- [8] S. G. Newman, M. Lautens, *J. Am. Chem. Soc.* **2010**, *132*, 11416.
- [9] a) A. H. Roy, J. F. Hartwig, *J. Am. Chem. Soc.* **2001**, *123*, 1232; b) A. H. Roy, J. F. Hartwig, *J. Am. Chem. Soc.* **2003**, *125*, 13944; c) A. H. Roy, J. F. Hartwig, *Organometallics* **2004**, *23*, 1533; d) F. Barrios-Landeros, B. P. Carrow, J. F. Hartwig, *J. Am. Chem. Soc.* **2009**, *131*, 8141.
- [10] a) D. A. Watson, M. Su, G. Teverovskiy, Y. Zhang, J. García-Fortanet, T. Kinzel, S. L. Buchwald, *Science* **2009**, *325*, 1661; b) X. Shen, A. M. Hyde, S. L. Buchwald, *J. Am. Chem. Soc.* **2010**, *132*, 14076.
- [11] a) S. G. Newman, M. Lautens, *J. Am. Chem. Soc.* **2011**, *133*, 1778; b) S. G. Newman, J. K. Howell, N. Nicolaus, M. Lautens, *J. Am. Chem. Soc.* **2011**, *133*, 14916; c) D. A. Petrone, H. A. Malik, A. Clemenceau, M. Lautens, *Org. Lett.* **2012**, *14*, 4806; d) X. Jia, D. A. Petrone, M. Lautens, *Angew. Chem.* **2012**, *124*, 10008; *Angew. Chem. Int. Ed.* **2012**, *51*, 9870.
- [12] Decomposition products made up the remainder of the mass balance.
- [13] E. M. Vogl, G. Gröger, M. Shibasaki, *Angew. Chem.* **1999**, *111*, 1672; *Angew. Chem. Int. Ed.* **1999**, *38*, 1570.
- [14] a) Y. Lan, P. Liu, S. G. Newman, M. Lautens, K. N. Houk, *Chem. Sci.* **2012**, *3*, 1987; b) J. F. Hartwig, *Inorg. Chem.* **2007**, *46*, 1936.
- [15] For a thorough discussion on the optimization, see the Supporting Information.
- [16] For examples, see: a) C. Copéret, E.-i. Negishi, *Org. Lett.* **1999**, *1*, 165; b) R. Grigg, V. Sridharan, *J. Organomet. Chem.* **1999**, *576*, 65.
- [17] a) C. C. C. Johansson Seechurn, S. L. Parisel, T. J. Colacot, *J. Org. Chem.* **2011**, *76*, 7918; b) H. Li, C. C. C. Johansson Seechurn, T. J. Colacot, *ACS Catal.* **2012**, *2*, 1147; c) X. Pu, H. Li, T. J. Colacot, *J. Org. Chem.* **2013**, *78*, 568.
- [18] The optimized **[1a]** was not suitable for this experiment, owing to severe peak broadening.