

Angewandte Berichtigung

Die Autoren dieser Zuschrift möchten einen zusätzlichen Beitrag zitieren. Zu diesem Zweck ist im zweiten Absatz der rechten Spalte auf S. 13851 ein Verweis auf Lit. [18] einzufügen: „Direct transformation of readily available alkanes into valuable complex alkyl ethers by transition-metal-catalyzed C(sp³)-H functionalization of unactivated methylenes is arguably a highly efficient and atom-economic method toward these compounds.^[18]“

[18] During the revision of our manuscript, an alkyl ether synthesis (alkoxylation of the β-C(sp³)-H bonds) through sp³ (methylene) C-H activation was reported by Shi's group with a new pyridine-based bidentate directing group and PhI(OAc)₂ as the oxidant. γ-Alkoxylation of C(sp²)-H bonds was also performed in this work: F. Chen, S. Zhao, F. Hu, K. Chen, Q. Zhang, S. Zhang, B. Shi, *Chem. Sci.* **2013**, *4*, 4187..

An Efficient Palladium-Catalyzed C-H Alkoxylation of Unactivated Methylene and Methyl Groups with Cyclic Hypervalent Iodine (I³⁺) Oxidants

G. Shan, X. Yang, Y. Zong,
Y. Rao* **13851–13855**

Angew. Chem. **2013**, 125

DOI: 10.1002/ange.201307090