



Auf diesen Seiten weisen wir auf wichtige aktuelle Beiträge in unseren Schwesterzeitschriften hin. Wenn Sie die Seiten online lesen, dann können Sie

die Artikel mit einem Klick direkt aufrufen, ansonsten sind sie durch Eingabe der DOIs über Wiley Online Library leicht online zugänglich.




Model Catalysis

D. J. Harding, A. Fielicke*

Platinum Group Metal Clusters: From Gas-Phase Structures and Reactivities towards Model Catalysts

Caught in the act: Transition-metal clusters have long been proposed as model systems to study heterogeneous catalysts. This Concept article shows how advanced spectroscopic techniques can be used to investigate gas-phase metal clusters and their complexes with small molecules, with the aim of developing an understanding of their reactivity (see figure).



Chem. Eur. J.
DOI: 10.1002/chem.201304586




Reaction Mechanisms


M. G. Apenova, V. A. Akhmetov, N. M. Belov, A. A. Goryunkov,*
I. N. Ioffe, N. S. Lukonina, V. Y. Markov, L. N. Sidorov

Alkali-Metal Trichloroacetates for Dichloromethylenation of Fullerenes: Nucleophilic Addition-Substitution Route

Looking for clues: The first experimental evidence that fullerenes react with alkali-metal trichloroacetates through a nucleophilic addition-substitution route to ultimately yield dichloromethylenefullerenes is reported (see picture). The structures of $C_{60}(CCl_3)H$ and four isomeric $C_{70}(CCl_3)H$ derivatives were unambiguously determined by means of NMR and UV/Vis spectroscopy.



Chem. Asian J.
DOI: 10.1002/asia.201301413




Tuberculosis


R. Dai, D. J. Wilson, T. W. Geders, C. C. Aldrich, B. C. Finzel*

Inhibition of *Mycobacterium tuberculosis* Transaminase BioA by Aryl Hydrazines and Hydrazides

TB or not TB: 7,8-Diaminopelargonic acid synthase (BioA) of *Mycobacterium tuberculosis* (*Mtb*) is a recently validated target in therapeutic interventions for tuberculosis (TB). In this paper, an aryl hydrazine *Mtb* BioA inhibitor is described that reversibly reacts with the PLP cofactor to form a stable *cis*-azo quinonoid species, which was structurally characterized by X-ray crystallography.



ChemBioChem
DOI: 10.1002/cbic.201300748





ChemPhysChem
DOI: 10.1002/cphc.201300958

Solar Cells

R. Narayanan, M. Deepa,* A. K. Srivastava, S. M. Shivaprasad

Efficient Plasmonic Dye-Sensitized Solar Cells with Fluorescent Au-Encapsulated C-Dots

Covering all possibilities: Au@C-dots function as plasmonic electron conduits and also undergo photo-induced charge separation in FRET-enabled high-performance ZnO based dye-sensitized solar cells.





ChemMedChem
DOI: 10.1002/cmdc.201300433

Epigenetics

A. S. Madsen, H. M. E. Kristensen, G. Lanz, C. A. Olsen*

The Effect of Various Zinc Binding Groups on Inhibition of Histone Deacetylases 1–11

Zinc about it: We synthesized a series of compounds containing diverse zinc binding motifs and profiled their inhibitory activity against the 11 human histone deacetylases (HDAC1–11). From the results, we discovered silanediol to be a novel functionality with potential as a suitable design element in the preparation of new HDAC inhibitors. Compounds of this type show promise as future drug leads.





ChemCatChem
DOI: 10.1002/cctc.201300811

Epoxidation

Y. Pang, X. Chen,* C. Xu, Y. Lei, K. Wei

High Catalytic Performance of MoO₃-Bi₂SiO₅/SiO₂ for the Gas-Phase Epoxidation of Propylene by Molecular Oxygen

A new sensation in epoxidation: We describe the probable synergistic effects of MoO₃ and Bi₂SiO₅ in propylene epoxidation. The reactive centers consist of nanoparticulate species of crystalline MoO₃ to activate the propylene and bismuth oxide cluster cations to activate O₂.




ChemPlusChem
DOI: 10.1002/cplu.201300277


Photocatalysis

S. Neubert, A. Ramakrishnan, J. Strunk, H. Shi, B. Mei, L. Wang, M. Bledowski, D. A. Guschin, M. Kauer, Y. Wang, M. Muhler, R. Beranek*

Surface-Modified TiO₂ Photocatalysts Prepared by a Photosynthetic Route: Mechanism, Enhancement, and Limits

Shedding light on things: Aromatic molecules like benzene or toluene can be activated by visible light ($\lambda > 455$ nm) at the surface of TiO₂, which leads to the formation of carbonaceous polymeric deposits on TiO₂ (see figure). The photosynthesized surface-modified TiO₂ shows enhanced photoactivity in the complete mineralization of 4-chlorophenol under visible-light irradiation.






Energy Materials

Y. Liu, S. Zhang,* T. Zhu*

Germanium-Based Electrode Materials for Lithium-Ion Batteries

In situ TEM electrochemistry is a powerful tool to study lithiation/delithiation and degradation mechanisms in battery electrodes in real time with high spatial resolution. Novel phenomena are uncovered in germanium-based electrode materials by using this technique.



ChemElectroChem
DOI: 10.1002/celc.201300195




Titanium Complexes


C. Adler, A. Bekurds, D. Haase, W. Saak, M. Schmidtman, R. Beckhaus*

Bulky Titanium Amides: C–H Bond Activation under Mild Conditions

The solid-state structures of tetrahedral and octahedral titanium amide complexes with bulky dicyclohexylamido ligands reveal a characteristic close contact between the titanium atom and a carbon atom in the β -position. The exchange of a chlorine atom by a bulky pentafulvene ligand increases this effect. The introduction of other sterically demanding amines to complexes initiates a spontaneous C–H activation.



Eur. J. Inorg. Chem.
DOI: 10.1002/ejic.201400022




Drug Design

S. Torijano-Gutiérrez, C. Vilanova, S. Díaz-Oltra, J. Murga,* E. Falomir, M. Carda, J. A. Marco*

Design and Synthesis of Pironetin Analogue/Combretastatin A-4 Hybrids and Evaluation of Their Cytotoxic Activity

The preparation of a series of hybrid molecules containing a combretastatin A-4 moiety and a pironetin analogue fragment linked by a spacer of variable length is described. The relationship between structure and cytotoxicity is discussed. Cytotoxicity values of some compounds were similar to those of the parent molecules, combretastatin A-4 and pironetin, and some were less toxic than the latter for normal cells.



Eur. J. Org. Chem.
DOI: 10.1002/ejoc.201301725




Heterocycles


M. Bielawski, J. Malmgren, L. M. Pardo, Y. Wikmark, B. Olofsson*

One-Pot Synthesis and Applications of N-Heteroaryl Iodonium Salts

N-Heteroarylation: An efficient one-pot synthesis of N-heteroaryl iodonium triflates from the corresponding N-heteroaryl iodide has been developed. The reactivity and chemoselectivity of these iodonium salts were demonstrated by selectively introducing a pyridyl moiety onto both oxygen and carbon nucleophiles in good yields.



ChemistryOpen
DOI: 10.1002/open.201300042




Asian J. Org. Chem.
DOI: 10.1002/ajoc.201400025

Organocatalysis

H. Qian, J. Sun*

Organocatalytic Enantio- and Diastereoselective Assembly of Thiazolidine Scaffolds by Formal [3+2] Annulation

Great balls of thia: An organocatalytic formal [3+2] annulation is described. With readily available achiral starting materials, a range of thiazolidines can be assembled in excellent yield and with moderate to good enantioselectivity and diastereoselectivity.



ChemViews magazine
DOI: 10.1002/chemv.201400002

Nanorods

J. Faiz

Copolymerization of Metal Nanoparticles

Professor Eugenia Kumacheva, Canada, talks to Dr. Jonathan Faiz, Senior Associate Editor for *Angewandte Chemie*, about her recent article. She discusses the significance of the molecular copolymerization approach described and how this strategy can have a broader impact by offering control over the structure and properties of nanoparticle assemblies.

