


Aerobic Oxidation in Nanomicelles of Aryl Alkynes, in Water at Room Temperature**

Sachin Handa, James C. Fennewald, and Bruce H. Lipshutz*

Abstract: On the basis of the far higher solubility of oxygen gas inside the hydrocarbon core of nanomicelles, metal and peroxide free aerobic oxidation of aryl alkynes to β -ketosulfones has been achieved in water at room temperature. Many examples are offered that illustrate broad functional group tolerance. The overall process is environmentally friendly, documented by the associated low *E* Factors.

Reactions in alternative media represent one approach to decreasing the huge amounts of organic waste generated by use of organic solvents in organic chemistry.^[1] While such options as ionic liquids, supercritical CO₂, and fluorinated media, among others, have made important inroads in this regard, the most likely and perhaps logical choice is water.^[2] Although we have investigated many processes enabled by designer surfactants where water serves as the gross reaction medium,^[3] synthetic advantage has yet to be taken of the well established far greater solubility of gases in organic media than in water.^[4] Since our reported cross-couplings and related reactions take place within the lipophilic cores of tailor-made micellar arrays, gases, as well as reactants and catalysts, should likewise co-exist in high concentrations and be available to participate in a given transformation. Surprisingly, there appears to be limited methodology^[5] of synthetic utility focused on the use of gases in micellar catalysis. Here we describe one such process involving dissolved oxygen serving as the stoichiometric oxidant, along with readily available aryl alkynes and sulfinic acids that leads to valuable β -ketosulfones under very mild, metal-free, and green conditions.


β -Ketosulfones can be derived from a radical reaction between an arylacetylene and a stoichiometric amount of a sulfinic acid. However, the short lifetime of the vinyl radical so generated^[6] (Scheme 1) makes trapping with oxygen difficult and prevents the desired reaction from having any generality in a purely aqueous or wet organic solvent. On the other hand, by virtue of the hydrophobic effect^[7] operating within the water-free lipophilic inner core of a nanomicelle, H-atom abstraction by a reactive vinyl radical by atom transfer radical addition (ATRA) from water or another



Scheme 1. Aerobic oxosulfonylation of arylacetylenes within nanomicelles in water.

molecule of alkyne should be minimized (path A). Rather, trapping of the vinyl radical by molecular oxygen is highly favored and ultimately leads to β -ketosulfone formation (path B). By contrast, related literature reports on this topic are far less environmentally friendly, as they take place in dry organic solvents at elevated temperatures with the use of metals and/or peroxide initiators,^[8] and offer no opportunities to recycle the reaction mixture. β -Ketosulfones are highly desirable materials, known to have fungicidal,^[9] antibacterial,^[10] as well as other biological properties.^[11] Moreover, numerous derivatives, such as olefins, disubstituted alkynes,^[12] allenes,^[13] and chiral vinyl sulfones^[14] and ketones^[13,14] have been prepared from such intermediates.

A model reaction between phenylacetylene **1** with *p*-toluenesulfinic acid sodium salt **2** was run at ambient temperature in an aqueous medium containing 2 wt. % TPGS-750-M (Scheme 2).^[15] The acid was generated in situ by the reaction of inexpensive **2** and HCl. Only traces of the desired product, however, were observed after 24 h. A large amount of unreacted phenylacetylene was recovered, along with byproducts, most likely due to rapid autoxidation of *p*-tolylsulfinic to the sulfonic acid and quenching of the vinyl radical by



Scheme 2. Aerobic difunctionalization of an arylacetylene.


[*] Dr. S. Handa, J. C. Fennewald, Prof. B. H. Lipshutz
Department of Chemistry & Biochemistry, University of California
Santa Barbara, CA 93106 (USA)
E-mail: lipshutz@chem.ucsb.edu

[**] Financial support for this work was provided by the U.S. National
Institutes of Health (GM 86485). We thank Shane Rainey for
technical assistance.

Supporting information for this article is available on the WWW
under <http://dx.doi.org/10.1002/ange.201310634>.

which is then trapped by oxygen to generate intermediate **33**. SET from another molecule of arylsulfinate to **33** generates arylperoxide anion **34**. The newly generated arylsulfonyl radical enters the next cycle, while **34** undergoes protonation either by water or a pyridinium cation to form arylhydroperoxide species **35**. Oxidation of arylsulfinate to arylsulfonate by **35** generates an enol that tautomerizes to final product **36**. The arylsulfonic acid generated as a byproduct remains in the aqueous phase, while the organic product can be isolated by extraction.

An E Factor^[20] of 5.3 was determined on the basis of organic solvent utilized for the model system (Scheme 8).^[21]



Scheme 8. E Factor and recycling studies.

This value compares quite favorably with those typical of the pharma and fine chemicals industries,^[22] as well as related literature.^[8,23] Moreover, recycling of the aqueous mixture led to good-to-excellent yields being obtained over three reaction cycles. The yield for the third cycle was noticeably lower, but this was due to practical considerations, as buildup of the sulfonic acid salt caused thickening and, therefore, problems with stirring on the scale at which the reaction was run.

In summary, an environmentally friendly aerobic oxidation has been developed for converting arylalkynes and arylsulfinate salts to β -ketosulfones. This process relies on the far greater solubility of oxygen in hydrocarbon media as found within micellar arrays than in the surrounding water. The process, enabled by a commercially available designer surfactant, is metal-free, takes place in water at room temperature using air as the stoichiometric oxidant, and is amenable to recycling of the aqueous reaction medium in which the amphiphile remains. Minimal amounts of organic solvent can be used to recover the desired product, which leads to a low E Factor. Experiments supporting a radical-based process are provided, along with data suggesting that the oxidation is taking place within the lipophilic core of the nanomicelles present in aqueous solution.

Received: December 7, 2013

Published online: February 24, 2014

Keywords: aerobic oxidation · E Factor · hydrophobic effect · micellar catalysis · TPGS-750-M

[1] a) D. J. Adams, P. J. Dyson, S. J. Tavener in *Chemistry in Alternative Reaction Media*, Wiley, Hoboken, **2005**, pp. 217–235; b) D. Reinhardt, F. Ilgen, D. Kralisch, B. Konig, G. Kreisel, *Green Chem.* **2008**, *10*, 1170–1181.

- [2] R. A. Sheldon, I. W. C. E. Arends, U. Hanefeld, *Green Chemistry and Catalysis*, Wiley-VCH, Weinheim, **2007**, pp. 1–47.
- [3] B. H. Lipshutz, S. Ghorai, *Aldrichimica Acta* **2012**, *45*, 3–16.
- [4] a) R. Battino, T. R. Rettich, T. Tominaga, *J. Phys. Chem. Ref. Data* **1983**, *12*, 163–178; b) I. B. Golovanov, S. M. Zhenodaro, *Russ. J. Gen. Chem.* **2005**, *75*, 1795–1797.
- [5] A. T. Straub, M. Otto, I. Usui, B. Breit, *Adv. Synth. Catal.* **2013**, *355*, 2071–2075.
- [6] a) P. J. Parsons, C. S. Penkett, A. J. Shell, *Chem. Rev.* **1996**, *96*, 195–206; b) U. Wille, *Chem. Rev.* **2013**, *113*, 813–853; c) K. Gilmore, I. V. Alabugin, *Chem. Rev.* **2011**, *111*, 6513–6556.
- [7] U. M. Lindström, F. Andersson, *Angew. Chem.* **2006**, *118*, 562–565; *Angew. Chem. Int. Ed.* **2006**, *45*, 548–551.
- [8] a) Q. Lu, J. Zhang, G. Zhao, Y. Qi, H. Wang, A. Lei, *J. Am. Chem. Soc.* **2013**, *135*, 11481–11484; b) Y. M. Markitanov, V. M. Timoshenko, Y. G. Shermolovich, *J. Sulfur Chem.* **2013**, 1–49; f) T. G. B. David, *Chemistry of Functional Groups*, Wiley, Hoboken, **2009**; c) W. Wei, C. Liu, D. Yang, J. Wen, J. You, Y. Suo, H. Wang, *Chem. Commun.* **2013**, *49*, 10239–10241; d) Q. Liu, R. Jackstell, M. Beller, *Angew. Chem. Int. Ed.* **2013**, *52*, 14115–14117; *Angew. Chem. Int. Ed.* **2013**, *52*, 13871–13873; e) Q. Lu, J. Zhang, F. Wei, Y. Qi, H. Wang, Z. Liu, A. Lei, *Angew. Chem.* **2013**, *125*, 7297–7300; *Angew. Chem. Int. Ed.* **2013**, *52*, 7156–7159.
- [9] W. M. Wolf, *J. Mol. Struct.* **1999**, *474*, 113–124.
- [10] C. Curti, M. Laget, A. O. Carle, A. Gellis, P. Vanelle, *Eur. J. Med. Chem.* **2007**, *42*, 880–884.
- [11] a) J. Xiang, M. Ipek, V. Suri, W. Massefski, N. Pan, Y. Ge, M. Tam, Y. Xing, J. F. Tobin, X. Xu, *Bioorg. Med. Chem. Lett.* **2005**, *15*, 2865–2869; b) J. Xiang, M. Ipek, V. Suri, M. Tam, Y. Xing, N. Huang, Y. Zhang, J. Tobin, T. S. Mansour, J. McKew, *Bioorg. Med. Chem.* **2007**, *15*, 4396–4405.
- [12] a) M. Ihara, S. Suzuki, T. Taniguchi, Y. Tokunaga, K. Fukumoto, *Tetrahedron* **1995**, *51*, 9873–9890; b) T. Mandai, T. Yanagi, K. Araki, Y. Morisaki, M. Kawada, J. Otera, *J. Am. Chem. Soc.* **1984**, *106*, 3670–3672.
- [13] J. E. Baldwin, R. M. Adlington, N. P. Crouch, R. L. Hill, T. G. Laffey, *Tetrahedron Lett.* **1995**, *36*, 7925–7928.
- [14] S. Sengupta, D. S. Sarma, S. Mondal, *Tetrahedron: Asymmetry* **1998**, *9*, 2311–2316.
- [15] B. H. Lipshutz, S. Ghorai, A. R. Abela, R. Moser, T. Nishikata, C. Duplais, A. Krasovskiy, R. D. Gaston, R. C. Gadwood, *J. Org. Chem.* **2011**, *76*, 4379–4391. Sigma–Aldrich catalog No. 733857.
- [16] For details about ATRA processes, see; a) C. P. Jasperse, D. P. Curran, T. L. Fevig, *Chem. Rev.* **1991**, *91*, 1237–1286; b) K. C. Majumdar, P. Debnath, *Tetrahedron* **2008**, *64*, 9799–9820; c) P. Balczewski, A. Szadowiak, T. Bialas, *Heteroat. Chem.* **2006**, *17*, 22–35.
- [17] For detailed optimization studies and procedures, see the Supporting Information.
- [18] Neither terminal alkylacetylenes, nor aryl alkyl alkynes reacted under these conditions.
- [19] N. A. Isley, F. Gallou, B. H. Lipshutz, *J. Am. Chem. Soc.* **2013**, *135*, 17707–17710.
- [20] a) R. A. Sheldon, *Green Chem.* **2007**, *9*, 1273–1283; b) S. Bonollo, D. Lanari, J. M. Longo, L. Vaccaro, *Green Chem.* **2012**, *14*, 164–169; c) M. M. Kirchhoff, *J. Chem. Educ.* **2013**, *90*, 683–684.
- [21] B. H. Lipshutz, N. A. Isley, J. C. Fennelwald, E. D. Slack, *Angew. Chem.* **2013**, *125*, 11156–11162; *Angew. Chem. Int. Ed.* **2013**, *52*, 10952–10958.
- [22] R. A. Sheldon, I. W. C. E. Arends, U. Hanefeld, *Green Chemistry and Catalysis*, Wiley-VCH, Weinheim, **2007**.
- [23] A. Weichert, H. M. R. Hoffmann, *J. Org. Chem.* **1991**, *56*, 4098–4112.