

Hexafluoroantimonic Acid Catalysis: Formal [3+2+2] Cycloaddition of Aziridines with Two Alkynes**


Ming-Bo Zhou, Ren-Jie Song, and Jin-Heng Li*

Dedicated to Professor Ming-Cai Chen on the occasion of his 60th birthday

Abstract: A practical method for the synthesis of azepine derivatives, a typical seven-membered heterocyclic ring system, was developed and involves the use of hexafluoroantimonic acid to catalyze a formal [3+2+2] cycloaddition of aziridines with two alkynes. This method was applicable to two of the same or different terminal alkynes for the [3+2+2] cycloaddition with unactivated aziridines, and furnished the corresponding azepine derivatives in good yields with good levels of chemo- and regioselectivity. The mechanism was also discussed according to the results of the *in situ* HRMS and ¹H NMR analysis.

The cycloaddition reaction has proven to be a powerful and straightforward synthetic tool for the atom-economical construction of cyclic compounds in modern organic chemistry.^[1–4] In the cycloaddition field, an important strategy involving the use of the ring-openings of small strained rings as a key step, fascinates numerous researchers because it can be used to meet the synthetic demand of making bioactive natural products containing heterocyclic rings.^[1–3] These cycloaddition processes allow the ring-opening of small strained rings and subsequent reaction with 2π components to construct various rings, specifically five- and six-membered rings, through [3+2] or [4+2] modalities. Particularly, cycloaddition reactions involving ring-opening reactions of strained aziridines have been widely applied in the construction of nitrogen-containing five-membered rings.^[3] However, methods for the selective construction of larger nitrogen-containing rings, including nitrogen-containing seven-membered rings, are lacking.^[4]

Generally, aziridines, a class of strained small heterocycles, are used as the precursors for both zwitterionic 1,3-



Scheme 1. The cycloaddition of aziridines. Ts = 4-toluenesulfonyl.

dipoles (**A**; in the presence of Lewis acids; Scheme 1) and azomethine ylides (under irradiation or thermolysis) for [3+2] cycloaddition with 2π components such as alkenes and alkynes.^[3] We reasoned that aziridines could undergo the [3+2+2] cycloaddition with two 2π components when the nucleophilicity of nitrogen anion in the intermediate **B** was reduced, thus enabling a subsequent electrophilic addition to another 2π component to form nitrogen-containing seven-membered rings. Herein, we report a new strategy to access the stable nitrogen anion in intermediate **B** using the superacid HSBF₆, thus triggering a new formal [3+2+2] cycloaddition of unactivated aziridines to two of the same or different terminal alkynes to construct azepine architectures (Scheme 1 b). Such a reaction would be particularly valuable for the synthesis of azepine derivatives,^[4,5] a typical seven-membered heterocyclic ring system, which are synthetically versatile compounds in synthesis and important skeletal units found in numerous natural products, potent pharmaceuticals, and peptidomimetics.^[6]

We first investigated the proposed [3+2+2] cycloaddition reaction between 2-phenyl-1-tosylaziridine (**1a**) with phenylacetylene to optimize the reaction conditions (Table 1). Examination of a range of reaction temperatures, Brønsted acids, and solvents (entries 1–11) revealed the combination of the HSBF₆ as the catalyst and CH₂Cl₂ as the solvent at 40 °C to be most effective: treatment of **1a** with phenylacetylene and 15 mol % HSBF₆ in CH₂Cl₂ at 40 °C for 24 hours regioselectively afforded the desired azepine **2** in 76 % yield (entry 1). The results demonstrated that the reaction temperature affected the reaction: the yield of **2** was reduced to 60 % when the reaction was carried out at room temperature (entry 2). Of the amounts of HSBF₆ examined, it turned out that 15 mol % of HSBF₆ was perfect for the reaction (entries 1, 3, and 4). Notably, the absence of HSBF₆ resulted in no detectable amounts of **2** (entry 5). Subsequently, several

[*] M.-B. Zhou, R.-J. Song, Prof. Dr. J.-H. Li
State Key Laboratory of Chemo/Biosensing and Chemometrics
College of Chemistry and Chemical Engineering
Hunan University, Changsha 410082 (China)
E-mail: jhli@hnu.edu.cn

Prof. Dr. J.-H. Li
State Key Laboratory of Applied Organic Chemistry
Lanzhou University, Lanzhou 730000 (China)

[**] We thank the Natural Science Foundation of China (No. 21172060), Specialized Research Fund for the Doctoral Program of Higher Education (No. 20120161110041), and Hunan Provincial Natural Science Foundation of China (No. 13JJ2018) for financial support. M.-B.Z. also thanks the Hunan Provincial Innovation Foundation for Postgraduate (CX2013B154).

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/ange.201310944>.

Table 1: Screening optimal reaction conditions.^[a]

Entry	Variation from the standard conditions	Yield [%] ^[b]
1	none	76
2	at room temperature	60
3	HSbF ₆ (10 mol %)	58
4	HSbF ₆ (30 mol %)	61
5	without HSBF ₆	0
6	HOTf instead of HSBF ₆	12
7	HOAc instead of HSBF ₆	trace
8	HBF ₄ instead of HSBF ₆	6
9	CH ₂ ClCH ₂ Cl instead of CH ₂ Cl ₂ for 48 h	52
10	toluene instead of CH ₂ Cl ₂ for 48 h	trace
11	MeNO ₂ instead of CH ₂ Cl ₂ for 48 h	14
12 ^[c]	none for 72 h	73

[a] Reaction conditions: **1a** (0.2 mmol), phenylacetylene (0.8 mmol), HSBF₆·6H₂O (15 mol %), and CH₂Cl₂ (2 mL) at 40 °C under an argon atmosphere for 24 h. [b] Yield of isolated product. [c] **1a** (6 mmol, 1.638 g).

other Brønsted acids, such as HOTf, HOAc, and HBF₄, were tested (entries 6–8). Both HOTf and HBF₄ could catalyze the reaction, albeit in low yields after 24 hours (entries 6 and 8). However, HOAc had no effect on the reaction (entry 7). Screening revealed that the effect of solvents had a fundamental influence on the reaction (entries 1 and 9–11). While CH₂ClCH₂Cl was still an efficient solvent for the reaction (entry 9), both toluene and MeNO₂ displayed lower activity (entries 10 and 11). It is noteworthy that the reaction of 1.638 g (6 mmol) **1a** proceeds in good yield (entry 12).

With the standard reaction conditions in hand, the scope of this HSBF₆-catalyzed [3+2+2] cycloaddition reaction, with respect to aziridines reacting with two of the same terminal alkynes, was first exploited (Table 2). The standard reaction conditions were found to be compatible with a wide range of terminal alkynes, including aryl, heteroaryl, and aliphatic alkynes (**3–11**). Furthermore, several substituents, such as Me, MeO, Cl, and Br, on the aryl ring of alkynes were well tolerated (**3–9**). Alkynes having a *para*- or *meta*-methyl-substituted aryl group underwent the reaction with **1a** and HSBF₆ smoothly, thus providing the desired products **3** and **7** in 73 and 66 % yield, respectively. Importantly, the halogens Cl and Br were tolerated under the reaction conditions, thereby facilitating additional modifications at the halogenated positions (**5**, **6**, and **8**). When using a dimethyl-substituted aryl alkyne, satisfactory yield was still achieved under the same reaction conditions (**9**). We were pleased to find that this [3+2+2] cycloaddition reaction was applicable to the preparation of the thiophen-3-yl-containing azepine **10** in 67 % yield. Ethynylcyclopropane was also a suitable substrate for the reaction (**11**).

Gratifyingly, this catalyzed [3+2+2] cycloaddition protocol was subject to a variety of 1-tosylaziridines (**1**; Table 2, **12–18**). 2-(3-Chlorophenyl)-1-tosylaziridine, for instance, was successfully reacted with phenylacetylene and HSBF₆ to afford the product **12** in 66 % yield. We were delighted to

Table 2: HSBF₆-catalyzed [3+2+2] cycloaddition of aziridines (**1**) with two of the same terminal alkynes.^[a]


3, 73% (R³ = 4-MeC₆H₄) 4, 62% (R³ = 4-MeOC₆H₄) 5, 65% (R³ = 4-ClC₆H₄) 6, 76% (R³ = 4-BrC₆H₄) 7, 66% 8, 72% 9, 63% 10, 67% 11, 41% 12, 66% 13, 71% (R¹ = 4-MeC₆H₄) 14, 67% (R¹ = 4-BrC₆H₄) 15, 65% (R¹ = 4-ClC₆H₄) 16, 57% (R¹ = 4-NO₂C₆H₄) 17, 63% 18, 67%		

[a] Reaction conditions: **1** (0.2 mmol), alkyne (0.8 mmol), HSBF₆·6H₂O (15 mol %), and CH₂Cl₂ (2 mL) at 40 °C under argon atmosphere for 24 h. Yields are those of the isolated products.

discover that a number of substituents, Me, Br, Cl, and NO₂, at the *para* position of the 2-aryl moiety were perfectly tolerated, thus resulting in the corresponding products **13–16** in moderate to good yields. Interestingly, the naphthalen-1-yl group could be readily introduced into the azepine structure (**17**). It was noted that 2-methyl-2-phenyl-1-tosylaziridine was also viable for the formation of the azepine **18** in 67 % yield.

In light of the results described above, we next decided to examine the possibility of synthesizing azepines having different substituents at the 5- and 7-positions by using two different terminal alkynes (Table 3). As expected, the reaction of **1a** with two different terminal alkynes was successfully performed, thus furnishing the desired azepines **19–27** in moderate to good yields. For example, when **1a** was treated with phenylacetylene (the first alkyne) and 5 mol % HSBF₆ in CH₂Cl₂ at 0 °C for 15 minutes, with subsequent addition of 4-methylphenylacetylene (the second alkyne) and 10 mol % HSBF₆ and an increase in the reaction temperature to 40 °C for about 24 hours, 3,5-diphenyl-7-*p*-tolyl-1-tosyl-2,3-dihydro-1*H*-azepine (**19**) was delivered in 78 % yield. It was noted that the same reaction conditions could be viable for the [3+2+2]


Table 3: [3+2+2] Cycloaddition of **1a** with two different terminal alkynes.^[a]



[a] Reaction conditions: a mixture of **1a** (0.2 mmol), the first alkyne (0.4 mmol), $\text{HSbF}_6 \cdot 6\text{H}_2\text{O}$ (5 mol%), and CH_2Cl_2 (2 mL) was stirred at 0 °C under an argon atmosphere. After 15 min, both the second alkyne (0.4 mmol) and $\text{HSbF}_6 \cdot 6\text{H}_2\text{O}$ (10 mol%) were added and the mixture was stirred at 40 °C for about 24 h. Yields are those of the isolated products. [b] The mixture of **1a** (0.2 mmol), the first alkyne (0.4 mmol), $\text{HSbF}_6 \cdot 6\text{H}_2\text{O}$ (5 mol%), and CH_2Cl_2 (2 mL) was first stirred at 15 °C under argon atmosphere for 30 min.

cycloaddition of **1a** with phenylacetylene and another alkyne, such as 2-methylphenylacetylene, 4-chlorophenylacetylene, 3-bromophenylacetylene, (2-thienyl)acetylene, or 2-chloro-4-methylphenylacetylene, thus leading to the corresponding azepines **20–24**, which have different substituents at the 7-position, in moderate to good yields. Interestingly, the substituent at the 5-position of the azepines could also be varied simply by the use of different alkynes the first step. Both phenylacetylene and 10 mol % HSbF_6 were added after 4-methoxyphenylacetylene reacted with **1a** and 5 mol % HSbF_6 in CH_2Cl_2 at 15 °C for 30 minutes, thus providing the 5-(4-methoxyphenyl)-substituted azepine **25** in 67% yield. When using 4-methylphenylacetylene or (2-thienyl)acetylene as the first alkyne, the corresponding 4-methylphenyl- and 4-(2-thienyl)-substituted azepines **26** and **27**, respectively, were also obtained in good yields.

However, phenyl(2-phenylaziridin-1-yl)methanone (**1b**) was unreactive for the [3+2+2] cycloaddition reaction [Eq. (1)]. To understand the mechanism, the reaction of the enyne **29** was carried out [Eq. (2)]. The results disclosed that **29** could not be converted into **2** under the standard reaction conditions, thus suggesting that the current reaction does not include an enyne intermediate.



Scheme 2. Possible mechanism.

Consequently, the working mechanism outlined in Scheme 2 was proposed on the basis of the present results and the literature reports.^[3,7,8] Initially, the zwitterionic 1,3-dipole intermediate **C** is formed from the reaction of **1a** with HSbF_6 ,^[3,8] with subsequent electrophilic addition to phenylacetylene to afford the intermediate **D**.^[7,8] In this step, HSbF_6 can also serve to stabilize the nitrogen anion. Subsequently, **D** undergoes the second electrophilic addition to a second molecule of phenylacetylene to give the intermediate **E**.^[8] Finally, dipolar cyclization of **E** gives the desired azepine **2**.

In summary, we have developed the first HSbF_6 -catalyzed formal [3+2+2] cycloaddition of 1-tosylaziridines with two alkynes. This novel method provides a mild and general access to the azepine architectures with both excellent functional-group tolerance and good levels of selectivity control, thus representing a new [3+2+2] cycloaddition transformation using 1-tosylaziridines as zwitterionic 1,3-dipoles. Studies on the mechanism and applications of this formal [3+2+2] cycloaddition method in organic synthesis are currently underway in our laboratory.

Received: December 18, 2013

Revised: January 16, 2014

Published online: March 11, 2014

Keywords: alkynes · aziridines · cycloaddition · heterocycles · superacidic systems

[1] For selected reviews, see: a) *Advances in Cycloaddition*, Vol. 1–6, JAI, Greenwich, CT, **1988–1999**; b) *Tetrahedron Organic Chemistry Series*: W. Carruthers, *Cycloaddition Reactions in Organic Synthesis*, Pergamon, Elmsford, NY, **1990**; c) M. Lautens, W. Klute, W. Tam, *Chem. Rev.* **1996**, 96, 49; d) C. P. Dell, *Contemp.*

- Org. Synth.* **1997**, *4*, 87; e) L. Yet, *Chem. Rev.* **2000**, *100*, 2963; f) *Cycloaddition Reactions in Organic Synthesis* (Eds.: S. Kobayashi, K. A. Jørgensen), Wiley-VCH, Weinheim, **2002**; g) F. López, J. L. Mascareñas, *Beilstein J. Org. Chem.* **2011**, *7*, 1075.
- [2] For special reviews on the cycloaddition reactions for the synthesis of seven-membered rings, specifically seven-membered carbocyclic rings, see: a) G. Dyker, *Angew. Chem.* **1995**, *107*, 2407; *Angew. Chem. Int. Ed. Engl.* **1995**, *34*, 2223; b) M. Harmata, *Acc. Chem. Res.* **2001**, *34*, 595; c) M. A. Battiste, P. M. Pelphrey, D. L. Wright, *Chem. Eur. J.* **2006**, *12*, 3438; d) H. Butenschön, *Angew. Chem.* **2008**, *120*, 5367; *Angew. Chem. Int. Ed.* **2008**, *47*, 5287; e) M. Harmata, *Chem. Commun.* **2010**, *46*, 8886; f) M. Harmata, *Chem. Commun.* **2010**, *46*, 8904; g) A. G. Lohse, R. P. Hsung, *Chem. Eur. J.* **2011**, *17*, 3812; h) H. Pellissier, *Adv. Synth. Catal.* **2011**, *353*, 189; i) T. V. Nguyen, J. M. Hartmann, D. Enders, *Synthesis* **2013**, 845; j) K. E. O. Ylijoki, J. M. Stryker, *Chem. Rev.* **2013**, *113*, 2244.
- [3] For special reviews on the [3+2] cycloaddition of aziridines, see: a) J. B. Sweeney, *Chem. Soc. Rev.* **2002**, *31*, 247; b) X. E. Hu, *Tetrahedron* **2004**, *60*, 2701; c) I. D. G. Watson, L. Yu, A. K. Yudin, *Acc. Chem. Res.* **2006**, *39*, 194; d) X. L. Hou, J. Wu, R. H. Fan, C. H. Ding, Z. B. Luo, L. X. Dai, *Synlett* **2006**, 181; e) P. Dauban, G. Malik, *Angew. Chem.* **2009**, *121*, 9188; *Angew. Chem. Int. Ed.* **2009**, *48*, 9026; f) P. Liu, *Tetrahedron* **2010**, *66*, 2549; g) S. H. Krake, S. C. Bergmeier, *Tetrahedron* **2010**, *66*, 7337; h) I. Coldham, R. Hufton, *Chem. Rev.* **2005**, *105*, 2765; i) G. S. Singh, M. D'hooghe, N. De Kimpe, *Chem. Rev.* **2007**, *107*, 2080; j) A. L. Cardoso, T. M. V. D. Pinho e Melo, *Eur. J. Org. Chem.* **2012**, 6479; for selected recent papers on the [3+2] cycloaddition reactions of aziridines with alkenes: k) I. Ungureanu, P. Klotz, A. Mann, *Angew. Chem.* **2000**, *112*, 4790; *Angew. Chem. Int. Ed.* **2000**, *39*, 4615; l) T. Ozawa, T. Kurahashi, S. Matsubara, *Synlett* **2013**, 24, 2673, and references therein; with alkynes: m) P. A. Wender, D. Strand, *J. Am. Chem. Soc.* **2009**, *131*, 7528; n) J. Fan, L. Gao, Z. Wang, *Chem. Commun.* **2009**, 5021.
- [4] For papers on the cycloaddition methods for azepine synthesis including the [4+3] cycloaddition, see: a) D. J. Anderson, A. Fiassner, *J. Am. Chem. Soc.* **1971**, *93*, 4339; b) Y. Bai, J. Fang, J. Ren, Z. Wang, *Chem. Eur. J.* **2009**, *15*, 8975; c) N. D. Shapiro, F. D. Toste, *J. Am. Chem. Soc.* **2008**, *130*, 9244; d) C. S. Jeffrey, K. L. Barnes, J. A. Eickhoff, C. R. Carson, *J. Am. Chem. Soc.* **2011**, *133*, 7688; e) H. Liu, X. Li, Z. Chen, W.-X. Hu, *J. Org. Chem.* **2012**, *77*, 5184; f) X.-Q. Hu, J.-R. Chen, S. Gao, B. Feng, L. Q. Lu, W.-J. Xiao, *Chem. Commun.* **2013**, *49*, 7905; g) S.-G. Huang, H.-F. Mao, S.-F. Zhou, J.-P. Zou, W. Zhang, *Tetrahedron Lett.* **2013**, *54*, 6178; h) S. Cui, Y. Zhang, D. Wang, Q. Wu, *Chem. Sci.* **2013**, *4*, 3912; the [5+2] cycloaddition: i) P. A. Wender, T. M. Pedersen, M. J. C. Scanio, *J. Am. Chem. Soc.* **2002**, *124*, 15154; j) X.-Z. Shu, S. Huang, D. Shu, I. A. Guzei, W. Tang, *Angew. Chem.* **2011**, *123*, 8303; *Angew. Chem. Int. Ed.* **2011**, *50*, 8153; k) X.-Z. Shu, X. Li, D. Shu, S. Huang, C. M. Schienebeck, X. Zhou, P. J. Robichaux, W. Tang, *J. Am. Chem. Soc.* **2012**, *134*, 5211; l) E. Kanno, K. Yamanoi, S. Koya, I. Azumaya, H. Masu, R. Yamasaki, S. Saito, *J. Org. Chem.* **2012**, *77*, 2142; m) N. Iqbal, A. Fiksdahl, *J. Org. Chem.* **2013**, *78*, 7885; n) M.-B. Zhou, R.-J. Song, C.-Y. Wang, J.-H. Li, *Angew. Chem.* **2013**, *125*, 11005; *Angew. Chem. Int. Ed.* **2013**, *52*, 10805.
- [5] For representative papers on the classical methods for azepine synthesis including radicals, see: a) L. Yet, *Tetrahedron* **1999**, *55*, 9349; b) C. E. Masse, A. J. Morgan, J. S. Panek, *Org. Lett.* **2000**, *2*, 2571; ring-expansion: c) T. J. V. Bergen, R. M. Kellogg, *J. Org. Chem.* **1971**, *36*, 978; d) E. J. Kantorowski, M. J. Kurth, *Tetrahedron* **2000**, *56*, 4317; e) E. A. Ilardi, J. T. Njardarson, *J. Org. Chem.* **2013**, *78*, 9533; ring-closing metathesis: f) M. E. Maier, *Angew. Chem.* **2000**, *112*, 2153; *Angew. Chem. Int. Ed.* **2000**, *39*, 2073; g) P. Jakubec, A. Hawkins, W. Felzmann, D. J. Dixon, *J. Am. Chem. Soc.* **2012**, *134*, 17482; nitrene insertion or diene-conjugated nitrile ylide cyclization: h) *Comprehensive Heterocyclic Chemistry*, Vol. 9 (Ed.: A. R. Katritzky II), Pergamon, New York, **1996**, p. 21; metal-catalyzed cyclization: i) L. F. Tietze, R. Schimpf, *Synthesis* **1993**, 876; j) V. Lyaskovskiy, K. Bergander, R. Fröhlich, E.-U. Würthwein, *Org. Lett.* **2007**, *9*, 1049; k) H. Ohno, H. Hamaguchi, M. Ohata, S. Kosaka, T. Tanaka, *J. Am. Chem. Soc.* **2004**, *126*, 8744; l) T. O. Vieira, H. Alper, *Org. Lett.* **2008**, *10*, 485; m) H. He, W.-B. Liu, L.-X. Dai, S.-L. You, *Angew. Chem.* **2010**, *122*, 1538; *Angew. Chem. Int. Ed.* **2010**, *49*, 1496; n) I. Nakamura, M. Okamoto, Y. Sato, M. Terada, *Angew. Chem.* **2012**, *124*, 10974; *Angew. Chem. Int. Ed.* **2012**, *51*, 10816; o) T. Zou, X.-G. Zhang, J.-H. Li, C.-L. Deng, R.-Y. Tang, *Adv. Synth. Catal.* **2012**, *354*, 889; p) Z. Shi, C. Grohmann, F. Glorius, *Angew. Chem.* **2013**, *125*, 5503; *Angew. Chem. Int. Ed.* **2013**, *52*, 5393; q) L. Wang, J. Huang, S. Peng, H. Liu, X. Jiang, J. Wang, *Angew. Chem.* **2013**, *125*, 1812; *Angew. Chem. Int. Ed.* **2013**, *52*, 1768; r) F. Grillet, K. M. Brummond, *J. Org. Chem.* **2013**, *78*, 3737; s) D. Solé, F. Mariani, *J. Org. Chem.* **2013**, *78*, 8136; t) Z. Dong, C.-H. Liu, Y. Wang, M. Lin, Z.-X. Yu, *Angew. Chem.* **2013**, *125*, 14407–14411; *Angew. Chem. Int. Ed.* **2013**, *52*, 14157–14161; Lewis acid mediated multicomponent reactions: u) G. Yin, Y. Zhu, P. Lu, Y. Wang, *J. Org. Chem.* **2011**, *76*, 8922.
- [6] For selected reviews, see: a) T. Kametani, K. Fukumoto, *Heterocycles* **1975**, *3*, 931; b) B. Renfroe, C. Harrington, G. R. Proctor, *Heterocyclic Compounds: Azepines*, Wiley & Interscience, New York, **1984**; c) C. R. Ganellin, D. J. Triggle, *Dictionary of Pharmacological Agents*, Chapman & Hall/CRC, London, **1996**; d) J. H. Ryan, C. Hyland, J. Just, A. G. Meyer, J. A. Smith, C. C. Williams, *Seven-Membered Rings in Progress in Heterocyclic Chemistry*, Vol. 25 (Eds.: G. W. Gribble, J. A. Joule), Elsevier, Amsterdam, **2013**, p. 455; for selected papers: e) S. Ohshima, M. Yanagisawa, A. Katoh, T. Fujii, T. Sano, S. Matsukuma, T. Furumai, M. Fujiu, K. Watanabe, K. Yokose, M. Arisawa, T. Okuda, *J. Antibiot.* **1994**, *47*, 639; f) R. M. Freidinger, D. E. Verber, D. Schwenk Perlow, J. R. Brooks, R. Saperstein, *Science* **1980**, *210*, 656; g) R. M. Freidinger, D. Schwenk Perlow, D. F. Verber, *J. Org. Chem.* **1982**, *47*, 104; h) K. Sakata, K. Aoki, C.-F. Chang, A. Sakurai, S. Tamura, S. Murakoshi, *Agric. Biol. Chem.* **1978**, *42*, 457; i) Y. Ye, G.-W. Qin, R.-S. Xu, *Phytochemistry* **1994**, *37*, 1205; j) S. Jiwajinda, N. Hirai, K. Watanabe, V. Santisopasri, N. Chuengsamarnyart, K. Koshimizu, H. Ohigashi, *Phytochemistry* **2001**, *56*, 693; k) J. Hu, M. J. Miller, *Tetrahedron Lett.* **1995**, *36*, 6379; l) Y.-F. Song, Y. Qu, X.-P. Cao, W. Zhang, *Mar. Biotechnol.* **2011**, *13*, 868; m) F. J. Villani, *J. Med. Chem.* **1967**, *10*, 497; n) H. J. M. Gijzen, D. Berthelot, M. Zaja, B. Bro ne, I. Geuens, M. Mercken, *J. Med. Chem.* **2010**, *53*, 7011; o) S. Tomasi, J. Renault, B. Martin, S. Duhieu, V. Cerec, M. L. Roch, P. Uriac, J.-G. Delcros, *J. Med. Chem.* **2010**, *53*, 7647; p) A. K. Ghose, T. Herbertz, H. Torsten, L. Robert, B. D. Dorsey, J. P. Mallamo, *ACS Chem. Neurosci.* **2012**, *3*, 50; q) R. N. Nicolas, R. Hernandez-Gallego, *Cardiovasc. Ther.* **2012**, *30*, 193; r) M. L. Granda, S. M. Carlin, C. K. Moseley, R. Neelamegam, J. B. Mandeville, J. M. Hooker, *ACS Chem. Neurosci.* **2013**, *4*, 261.
- [7] a) G. A. Olah, R. J. Spear, *J. Am. Chem. Soc.* **1975**, *97*, 1845; b) P. J. Stang, R. Summerville, *J. Am. Chem. Soc.* **1969**, *91*, 4600.
- [8] The formation of **A**, **B**, and **C** was supported by in situ HRMS analysis, and the detailed data summarized in Figure S1 in Supporting Information. In addition, four other intermediates were also observed by the in situ HRMS analysis:

