

Synthetic Methods

Direct Assembly of 3,4-Difunctionalized Benzofurans and Polycyclic Benzofurans by Phenol Dearomatization and Palladium-Catalyzed Domino Reaction**


Zhaomeng Han, Liang Zhang, Zhiming Li,* and Renhua Fan*

Dedicated to Professor Li-Xin Dai on the occasion of his 90th birthday

Abstract: A method to directly convert 2-alkynylphenols to 3,4-difunctionalized benzofurans and polycyclic benzofurans was developed. This protocol involves a hypervalent-iodine-mediated oxidative dearomatization to break the aromaticity of 2-alkynylphenols, and a palladium-catalyzed domino reaction to install two functional groups at the C3 and the C4 positions and restore the aromaticity of benzofurans.

The dearomatization of aromatic compounds provides numerous possibilities for the construction of complex molecules.^[1] The main advantage of the dearomatization is the possibility of converting an aromatic ring into a three-dimensional molecule. Moreover, since dearomatization offers unique strategic opportunities to circumvent the inherent *ortho/para* selectivity of electron-rich aromatic systems, the dearomatization strategy can also be used in the synthesis of multi-functionalized aromatic compounds that are difficult to prepare by electrophilic substitution reactions.


Among the family of benzofurans, 3,4-difunctionalized benzofurans are attractive synthetic targets because of their remarkable biological activities.^[2] The synthetic challenge is the selective functionalization of the C4 position of benzofurans, which is not a preferred site for electrophilic substitution reactions. 2-Alkynylphenols are the most commonly used precursors to prepare benzofurans. A number of cyclization or cascade cyclization/cross-coupling reactions of 2-alkynylphenols have been developed.^[3–12] However, these elegant methods only enable the synthesis of benzofurans with diverse substitutions on the five-membered ring. Herein, we report a method that directly converts 2-alkynylphenols to 3,4-difunctionalized benzofurans (Scheme 1). This protocol involves an oxidative dearomatization to break the aromaticity of 2-alkynylphenols, a palladium-catalyzed domino reaction to simultaneously install two functional groups at



Scheme 1. Preparation of 3,4-difunctionalized benzofurans from 2-alkynylphenols.

the C3 and the C4 positions, and an aromatization to restore the aromaticity.

(Diacetoxyiodo)benzene facilitated oxidative dearomatization of 4-methyl-2-(2-phenylethynyl)phenol **1** in methanol. The deleterious cyclization or oxidation of the sensitive alkynyl group was not observed. The crude dearomatization product was directly used to test the palladium-catalyzed domino reaction with *p*-toluidine and ethyl acrylate. When 0.1 equivalents of PdCl₂ were used together with 0.2 equivalents of Ph₃P, 4-amino-substituted 3-alkenylbenzofuran **2** was obtained in 7% yield. Its structure was confirmed by single-crystal diffraction analysis.^[13] A screening of solvents, temperatures, and bases did not improve the yield beyond 10%, thus indicating that the reaction with palladium was stoichiometric. Therefore, various oxidants were added to promote the regeneration of catalytic Pd^{II} from Pd⁰ formed in the Heck coupling. While the addition of 2 equivalents of benzoquinone (BQ) improved the yield to 31% [Eq. (1)],



compound **2** was not formed when Cu(OAc)₂, CuCl₂, AgOAc, *meta*-chloroperbenzoic acid (*m*-CPBA), or PhI(OAc)₂ were added. The product formation was further optimized by examining a variety of palladium salts and phosphine ligands. When PdBr₂ was used together with trifuran-2-ylphosphine, the yield of compound **2** increased to 76%.^[14]

After establishing the optimized reaction conditions, the scope of this transformation was investigated (Table 1). For a range of 2-alkynylphenols, the reactions proceeded smoothly, leading to the corresponding 3,4-difunctionalized benzofurans in moderate to good yields. In some cases, TsOH was added to promote the aromatization, and PdCl₂ was used

[*] Z. Han, L. Zhang, Prof. Dr. Z. Li, Prof. Dr. R. Fan
Department of Chemistry, Fudan University
220 Handan Road, Shanghai, 200433 (China)
E-mail: zml@fudan.edu.cn
rhfan@fudan.edu.cn

[**] We are grateful to the National Science Foundation of China (21332009, 21102019), and the Shanghai Science and Technology Committee (13431900103) for support of this research.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/ange.201404155>.

Table 1: Dearomatization and palladium-catalyzed domino reaction.

Entry	Product		Yield ^[a]
1		2: R ¹ = Ph	76
2 ^[b]		3: R ¹ = 4-MeC ₆ H ₄	61
3		4: R ¹ = 4-ClC ₆ H ₄	74
4 ^[c]		5: R ¹ = <i>n</i> Bu	65
5		6: R ¹ = <i>t</i> Bu	76
6		7: R ¹ = cyclopropyl	60
7 ^[b,c]		8: R ¹ = TMS	71
8 ^[b,c]		9: R ² = <i>n</i> Bu	55
9		10: R ² = Ph	0
10		11: R ² = OMe	65
11 ^[b]		12: R ³ = Me	78
12		13: R ⁴ = CN	63
13		14: R ⁴ = CON(Me) ₂	62
14		15: R ⁴ = Ph	0
15 ^[b]		16: R ⁵ = Ph	61
16 ^[b,c]		17: R ⁵ = 4- <i>i</i> PrC ₆ H ₄	72
17		18: R ⁵ = 4- <i>n</i> BuC ₆ H ₄	72
18		19: R ⁵ = 4-MeOC ₆ H ₄	82
19 ^[b,c]		20: R ⁵ = 3-MeOC ₆ H ₄	42
20		21: R ⁵ = 4-FC ₆ H ₄	61
21 ^[b,c]		22: R ⁵ = 4-ClC ₆ H ₄	65
22 ^[b,c]		23: R ⁵ = 4-BrC ₆ H ₄	71
23 ^[b,c]		24: R ⁵ = 2-IC ₆ H ₄	60
24		25: R ⁵ = Bn	0
25		26: R ⁵ = <i>n</i> Bu	0
26		27: R ⁶ = Me	64

[a] Reported yields are of the isolated products. [b] 4-Methylbenzenesulfonic acid (4 equiv) was added after 12 h. [c] PdCl₂ was used instead of PdBr₂. Bn = benzyl, TMS = trimethylsilyl.

instead of PdBr₂ to improve product yields. Electron-deficient alkenes, such as acrylonitrile and *N,N*-dimethylacrylamide, were also suitable reaction partners. When styrene was used, the reaction only afforded 4-amino-substituted benzofuran **28**. Various aromatic amines, including a secondary amine, could be used as nucleophile to be introduced at the C4 position of benzofuran. When benzylamine or butan-1-amine was used, no desired product was obtained, but a 4-methoxy-substituted 3-alkenylbenzofuran **29** was isolated. This compound was also the major product when diethyl malonate, phenol, or thiophenol were used as nucleophile. Interestingly, compound **29** was formed in a 63 % yield in the absence of an added nucleophile. When 1 equivalent of CD₃OD was added

as nucleophile, the incorporation of the OCD₃ group into compound **29** was not observed. This result indicated that compound **29** might be formed through a [1,2] transfer of the methoxy group.

To understand the formation of product **2**, 4-amino-substituted benzofuran **28** was treated with ethyl acrylate under the standard conditions. The formation of compound **2** was not observed. Meanwhile, the isolated non-aromatized product **30** was completely converted to compound **2** when it was treated with PdBr₂ in ClCH₂CH₂Cl at reflux.

To gain more insight into the reaction, B3LYP density functional theory (DFT) calculations were performed with the Gaussian09 software package.^[15] We initially evaluated the role of the palladium catalyst. Palladium(II) might work as a Lewis acid to coordinate with the carbonyl group of the dearomatization product by a monodentate coordination mode or with the carbonyl group and the triple bond by a bidentate mode. Palladium(II) might also work as a π acid to coordinate with the triple bond by a monodentate mode. This activation might induce a cyclization to generate a furan-like intermediate (see Scheme S1 in the Supporting Information).^[16] The computational results indicated that the furan-like intermediate **A** is the most stable intermediate (Figure 1).




Figure 1. Optimized structures of intermediate **A**. The numbers in parenthesis are the NBO charges on atoms and the numbers in square brackets are the condensed Fukui functions.

It is more than 5 kcal mol⁻¹ more stable than other intermediates. Therefore, palladium(II) acts as a π acid in the initial step. Moreover, although there are two possible reaction sites (C3 and C5) for the nucleophilic attack in the structure of intermediate **A**, the natural bond orbital (NBO) analysis showed that the C3 position is more positively charged (−0.02) compared to the C5 position (−0.11). The condensed Fukui function also predicted that the nucleophilic attack will occur at the C3 position. These results are in line with our experimental results.

We identified two possible reaction pathways from intermediate **A** to the non-aromatized product to understand the domino sequence. For clarity, only main stationary points are shown in Figure 2, and the relative free energies are based on the total free energy of intermediate **A**, *p*-toluidine, and methyl acrylate. In one pathway (in black), intermediate **A** is trapped by the electron-poor double bond of acrylate through






Figure 2. Free-energy reaction profile (kcal mol⁻¹) from intermediate **A** to the non-aromatized product, calculated at the PCM(DCE) B3LYP/6-31G(d,p) with LANL2DZ (for Pd and Br) level. TS = transition state.


syn insertion, thus affording intermediate **D**. Intermolecular nucleophilic attack of the nitrogen of toluidine on the C3 position forms intermediate **E**. Subsequent *syn* β -H elimination gives rise to the non-aromatized product and releases H₂CO₃ and HPdBr. In the alternative pathway (in red), intermolecular nucleophilic attack occurs before the insertion, thus leading to intermediate **J**. After releasing H₂CO₃, intermediate **K** reacts with acrylate through insertion, followed by β -H elimination to afford the non-aromatized product and generate HPdBr. The potential energy surfaces indicate that the rate-determining step for the first pathway is the β -H elimination step (TS EF), while it is only 7.9 kcal mol⁻¹ for the second one. This means that the domino reaction favors the second pathway to a large extent. On the basis of these results, a plausible mechanism is depicted in Scheme 2. Palladium might play



Scheme 2. Plausible mechanism.




the intramolecular insertion of organopalladium to the double bond of vinyl ketone or 2-methylenemalonate. The protonolysis of this intermediate might be more favorable than the β -hydride elimination to give a conjugate addition product.^[17] In these palladium(II)-catalyzed reactions, the addition of benzoquinone was not required. For most cases, polycyclic benzofurans were produced in moderate to good yields. The linkage between the alkynyl and the alkenyl group could be a benzene ring or an alkyl chain. The reaction of aromatic amines was found to tolerate a range of substituents, involving electron-withdrawing and electron-donating groups. When an unprotected aminophenol was employed,



Scheme 3. Construction of polycyclic benzofuran derivatives.

the reaction only afforded the *N*-arylated product. The electron-rich indoles could serve as suitable reaction partners in this process. With respect to other electron-rich aromatic compounds, such as *N,N*-dimethylaniline, 1,2-dimethoxybenzene, and thiophene, the reactions were complex and the desired products were not obtained.

With the aid of trifluoromethanesulfonic acid, compound **2** could be converted to compound **51** through an intramolecular Friedel–Crafts alkylation in a 94% yield [Eq. (3)]. Compound **51** has a dibenz[*b,f*]azepine core, which exists in



many natural products and medicinal compounds, such as carbazepine and trileptal.^[18]

In conclusion, we have developed a method to convert 2-alkynylphenols into 3,4-difunctionalized benzofurans and polycyclic benzofurans. This protocol involves a hypervalent-iodine-mediated oxidative dearomatization and a palladium-catalyzed domino reaction. The application of this strategy to the synthesis of natural products and investigations on a more detailed mechanism are currently underway in our laboratory.

Experimental Section

Representative procedure: $\text{PhI}(\text{OAc})_2$ (0.22 mmol) was added to a solution of 4-methyl-2-(2-phenylethynyl)phenol **1** (0.2 mmol) in MeOH (2.0 mL) at 25°C. After 5 min, the reaction mixture was concentrated in vacuo. The resulting crude product was mixed with ethyl acrylate (0.4 mmol), *p*-toluidine (0.3 mmol), PdCl_2 (0.02 mmol), trifuran-2-ylphosphine (0.04 mmol), benzoquinone (0.4 mmol), and K_2CO_3 (0.4 mmol) in $\text{ClCH}_2\text{CH}_2\text{Cl}$ (2 mL). The reaction was stirred under reflux. After the substrate was completely consumed (monitored by TLC analysis), the reaction mixture was passed through a short column of silica gel and then concentrated under reduced pressure. The residue was purified by flash column chromatography on silica gel (eluent: petroleum ether/ethyl acetate) to furnish the desired compound **2**.

Received: April 9, 2014

Published online: May 20, 2014

Keywords: benzofurans · dearomatization · domino reactions · multicomponent reactions · phenols

- [1] For recent reviews on dearomatization, see: a) A. R. Pape, K. P. Kaliappan, E. P. Kündig, *Chem. Rev.* **2000**, *100*, 2917–2940; b) D. Magdziak, S. J. Meek, T. R. R. Pettus, *Chem. Rev.* **2004**, *104*, 1383–1429; c) F. López Ortiz, M. J. Iglesias, I. Fernández, C. M. Andújar Sánchez, G. R. Gómez, *Chem. Rev.* **2007**, *107*, 1580–1691; d) S. Quideau, L. Pouységu, D. Deffieux, *Synlett* **2008**, 467–495; e) L. Pouységu, D. Deffieux, S. Quideau, *Tetrahedron* **2010**, *66*, 2235–2261; f) S. P. Roche, J. A. Porco, Jr., *Angew. Chem.* **2011**, *123*, 4154–4179; *Angew. Chem. Int. Ed.* **2011**, *50*, 4068–4093; g) C.-X. Zhuo, W. Zhang, S.-L. You,

Angew. Chem. **2012**, *124*, 12834–12858; *Angew. Chem. Int. Ed.* **2012**, *51*, 12662–12686; h) Q. Ding, Y. Ye, R. Fan, *Synthesis* **2013**, 1–13.

- [2] a) W. Friedrichsen in *Comprehensive Heterocyclic Chemistry II*, Vol. 2 (Eds.: C. W. Bird, A. R. Katritzky, C. W. Rees, E. F. V. Scriven), Pergamon, London, **1996**, pp. 368–378; b) *Comprehensive Heterocyclic Chemistry II*, Vol. 2 (Eds.: A. R. Katritzky, C. W. Rees, E. F. V. Scriven), Pergamon, Oxford, UK, **1996**, pp. 259–321; c) X. L. Hou, Z. Yang, K. S. Yeung, H. N. C. Wong in *Progress in Heterocyclic Chemistry*, Vol. 19 (Eds.: G. W. Gribble, J. A. Joule), Elsevier, Oxford, **2008**, pp. 176–207, and previous volumes in the series.
- [3] For recent reviews on the synthesis of benzofurans, see: a) S. Cacchi, G. Fabrizi, A. Goggioni, *Heterocycles* **2002**, *56*, 613–632; b) Y. Liao, Y. Hu, J. Wu, Q. Zhu, M. Donovan, R. Fathi, Z. Yang, *Curr. Med. Chem.* **2003**, *10*, 2285–2316; c) G. Zeni, R. C. Larock, *Chem. Rev.* **2004**, *104*, 2285–2309; d) I. Nakamura, Y. Yamamoto, *Chem. Rev.* **2004**, *104*, 2127–2198; e) F. Alonso, I. P. Beletskaya, M. Yus, *Chem. Rev.* **2004**, *104*, 3079–3160.
- [4] For carbonylative annulation of 2-alkynylphenols, see: a) Y. Nan, H. Miao, Z. Yang, *Org. Lett.* **2000**, *2*, 297–299; b) Y. Hu, Y. Zhang, *Org. Lett.* **2001**, *3*, 1387–1390; c) Y. Hu, Y. Zhang, Z. Yang, R. Fathi, *J. Org. Chem.* **2002**, *67*, 2365–2368; d) Y. Liao, J. Smith, R. Fathi, Z. Yang, *Org. Lett.* **2005**, *7*, 2707–2709.
- [5] For the synthesis of 2,3-functionalized benzofurans through 3-zincobenzofuran intermediates, see: a) M. Nakamura, L. Ilies, S. Otsubo, E. Nakamura, *Angew. Chem.* **2006**, *118*, 958–961; *Angew. Chem. Int. Ed.* **2006**, *45*, 944–947; b) M. Nakamura, L. Ilies, S. Otsubo, E. Nakamura, *Org. Lett.* **2006**, *8*, 2803–2805; c) H. Tsuji, C. Mitsui, L. Ilies, Y. Sato, E. Nakamura, *J. Am. Chem. Soc.* **2007**, *129*, 11902–11903.
- [6] For tandem cyclization/Heck coupling reactions, see: a) N. Isono, M. Lautens, *Org. Lett.* **2009**, *11*, 1329–1331; b) C. Martínez, R. Álvarez, J. M. Aurrecochea, *Org. Lett.* **2009**, *11*, 1083–1086; c) R. Álvarez, C. Martínez, Y. Madich, J. G. Denis, J. M. Aurrecochea, Á. R. de Lera, *Chem. Eur. J.* **2010**, *16*, 12746–12753; d) C. Martínez, J. M. Aurrecochea, Y. Madich, J. G. Denis, A. R. de Lera, R. Álvarez, *Eur. J. Org. Chem.* **2012**, 99–106.
- [7] For annulative electrophilic amination, see: a) N. Matsuda, K. Hirano, T. Satoh, M. Miura, *J. Org. Chem.* **2012**, *77*, 617–625; b) K. Hirano, T. Satoh, M. Miura, *Org. Lett.* **2011**, *13*, 2395–2397.
- [8] For palladium-catalyzed cyclization and coupling with triflates or halides, see: a) A. Arcadi, S. Cacchi, M. D. Rosario, G. Fabrizi, F. Marinelli, *J. Org. Chem.* **1996**, *61*, 9280–9288; b) Y. Hu, K. J. Nawoschik, Y. Liao, J. Ma, R. Fathi, Z. Yang, *J. Org. Chem.* **2004**, *69*, 2235–2239.
- [9] For palladium-catalyzed electrophilic annulation, see: a) Y. Liang, S. Tang, X.-D. Zhang, L.-Q. Mao, Y.-X. Xie, J.-H. Li, *Org. Lett.* **2006**, *8*, 3017–3020; b) H.-A. Du, X.-G. Zhang, R.-Y. Tang, J.-H. Li, *J. Org. Chem.* **2009**, *74*, 7844–7848.
- [10] For tandem cyclization/dehydrogenative coupling reactions, see: H. Hachiya, K. Hirano, T. Satoh, M. Miura, *Org. Lett.* **2011**, *13*, 3076–3079.
- [11] For carboalkoxylation of 1-alkynylphenyl ethers, see: a) A. Arcadi, S. Cacchi, S. D. Giuseppe, G. Fabrizi, F. Marinelli, *Org. Lett.* **2002**, *4*, 2409–2412; b) I. Nakamura, Y. Mizushima, Y. Yamamoto, *J. Am. Chem. Soc.* **2005**, *127*, 15022–15023; c) A. Fürstner, P. W. Davies, *J. Am. Chem. Soc.* **2005**, *127*, 15024–15025; d) Z. Liang, S. Ma, J. Yu, R. Xu, *J. Org. Chem.* **2007**, *72*, 9219–9224; e) A. Fürstner, E. K. Heilmann, P. W. Davies, *Angew. Chem.* **2007**, *119*, 4844–4847; *Angew. Chem. Int. Ed.* **2007**, *46*, 4760–4763.
- [12] For the cyclization with allyl carbonate, see: S. Cacchi, G. Fabrizi, L. Moro, *Synlett* **1998**, 741–745.

- [13] CCDC 915522 (**2**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [14] For the evaluation of conditions and catalysts, see Table S1 in the Supporting Information.
- [15] For computational details and references, see the Supporting Information.
- [16] a) T. Yao, X. Zhang, R. C. Larock, *J. Am. Chem. Soc.* **2004**, *126*, 11164–11165; b) T. Yao, X. Zhang, R. C. Larock, *J. Org. Chem.* **2005**, *70*, 7679–7685; c) Y. Xiao, J. Zhang, *Angew. Chem.* **2008**, *120*, 1929–1932; *Angew. Chem. Int. Ed.* **2008**, *47*, 1903–1906.
- [17] For competition between conjugate addition and Heck-coupling reaction, see: a) S. Cacchi, A. Arcadi, *J. Org. Chem.* **1983**, *48*, 4236–4240; b) T. Nishikata, Y. Yamamoto, N. Miyaara, *Angew. Chem.* **2003**, *115*, 2874–2876; *Angew. Chem. Int. Ed.* **2003**, *42*, 2768–2770; c) A. L. Gottumukkala, J. G. de Vries, A. J. Minnaard, *Chem. Eur. J.* **2011**, *17*, 3091–3095; d) Q. Peng, H. Yan, X. Zhang, Y.-D. Wu, *J. Org. Chem.* **2012**, *77*, 7487–7496.
- [18] a) R. G. Lande, *Int. J. Psychiatry Clin. Pract.* **2004**, *8*, 37–40; b) N. Della Cá, G. Maestri, M. Malacria, E. Derat, M. Catellani, *Angew. Chem.* **2011**, *123*, 12465–12469; *Angew. Chem. Int. Ed.* **2011**, *50*, 12257–12261.
-