

Characterization of Mononuclear Non-heme Iron(III)-Superoxo Complex with a Five-Azole Ligand Set**

Frédéric Oddon, Yosuke Chiba, Jun Nakazawa,* Takehiro Ohta, Takashi Ogura, and Shiro Hikichi*

Abstract: Reaction of O_2 with a high-spin mononuclear iron(II) complex supported by a five-azole donor set yields the corresponding mononuclear non-heme iron(III)-superoxo species, which was characterized by UV/Vis spectroscopy and resonance Raman spectroscopy. 1H NMR analysis reveals diamagnetic nature of the superoxo complex arising from antiferromagnetic coupling between the spins on the low-spin iron(III) and superoxide. This superoxo species reacts with H-atom donating reagents to give a low-spin iron(III)-hydroperoxo species showing characteristic UV/Vis, resonance Raman, and EPR spectra.

In biological systems, interaction between O_2 and iron centers in heme and non-heme proteins/enzymes occurs in aerobic respiration and metabolic processes. Oxidative addition of O_2 to the ferrous centers of the proteins/enzymes yields the corresponding ferric complexes of reduced O_2 species (O_2^- and O_2^{2-}).^[1] The O_2 activation mechanism of heme iron oxygenases has been explored extensively both on the enzymes and related model compounds.^[2] In line with the chemical analogy, a mononuclear iron center of non-heme oxygenases is inferred to activate O_2 by a similar mechanism.^[3] To date, a few mononuclear non-heme iron(III)-superoxo species have been characterized in some enzymes and synthetic model systems.^[4,5] However, information of the chemical properties of such non-heme iron(III)-superoxo species is still scarce. To clarify the correlation between the coordination environments and properties of mononuclear non-heme iron centers such as spin state and reactivity, interests in investigation of synthetic model compounds are

much growing.^[5–10] As reported herein, we have successfully characterized a mononuclear iron(III)-superoxo complex with azole-based ligands.

We have designed a single-site unsaturated iron(II) complex with five azole donors by assembling tridentate hydrotris(3,5-dimethylpyrazolyl)borate (Tp^{Me_2})^[11] and bidentate imidazolyl-based borate $[B(Im^{N-Me})_2MePh]^-$ (L^{Ph})^[12] ligands. A family of Tp^R (where R denotes substituent groups on the pyrazolyl rings) is one of extensively explored ligands for the synthetic models of non-heme O_2 -activating iron centers because facially capping N3-donating Tp^R ligands can reproduce the coordination structures of the active sites of some enzymes involving (imidazole)₂X triad motif where X is an imidazole residue of His or a carboxylate of Glu or Asp.^[3] To date, several O_2 -activating mononuclear iron(II)- Tp^R complexes bearing an additional bidentate ligand Y, $[Fe^{II}(Y)(Tp^R)]$ (Y = carboxylate, α -keto carboxylate, β -diketionate, cysteinate, etc.) have been reported. But putative mononuclear iron(III)-superoxo species have never been detected in these Tp^R systems because of the formation of dinuclear iron(III)- μ -peroxo compounds occurring through the change of coordination mode of Y from terminal chelating ligand to bridging one, or intramolecular oxygenation of the coordinating Y proceeds immediately as found in the enzymatic reactions.^[13] In this study, the imidazolyl-based ligand L^{Ph} has been employed because of its strong bidentate chelating nature as well as robustness toward oxidative degradation. A desired mixed ligand complex, $[Fe^{II}(L^{Ph})(Tp^{Me_2})]$ (**2**), was prepared by ligand exchange reaction between LiL^{Ph} and an iron(II) acetate complex, $[Fe^{II}(OAc)(Tp^{Me_2})]$ (**1**) (Scheme 1). The molecular structure of **2** was revealed by X-ray crystallography (Figure 1). The iron center is surrounded by five azole donors with square-pyramidal geometry. Equatorial Fe–N bonds are composed of two

[*] Dr. F. Oddon, Y. Chiba, Dr. J. Nakazawa, Prof. Dr. S. Hikichi
Department of Material and Life Chemistry
Faculty of Engineering, Kanagawa University
Kanagawa-ku, Yokohama, 222-8686 (Japan)
E-mail: jnaka@kanagawa-u.ac.jp
hikichi@kanagawa-u.ac.jp

Dr. T. Ohta, Prof. Dr. T. Ogura
Department of Life Science, University of Hyogo
Sayo-cho, Sayo-gun, Hyogo 679-5148 (Japan)

[**] This work was supported in part by a Cooperative Research Program of "Network Joint Research Center for Materials and Devices" (No. 2012348), Grant-in-Aid for Scientific Research (No. 26420788) and a Strategic Development of Research Infrastructure for Private Universities from the Ministry of Education, Culture, Sports, Science, and Technology (MEXT) (Japan). We acknowledge prof. Toshiro Takao in Tokyo Institute of Technology for EPR measurement.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201502367>.





Figure 1. ORTEP of $[Fe(L^{Ph})(Tp^{Me_2})]$ **2**. Hydrogen atoms are omitted for clarity; ellipsoids are set at 30% probability.^[23]



Scheme 1. Reactions of iron complexes in this study.

longer bonds derived from Tp^{Me_2} and two shorter bonds from L^{Ph} . This *trans* influence indicates that the imidazolyl groups on L^{Ph} are stronger donors compared to the pyrazolyl ligands of Tp^{Me_2} .

As we expected, the five-coordinate iron(II) complex **2** reacted with O_2 to give the corresponding O_2 adduct **3**. Bubbling of O_2 gas to a THF solution of **2** at -60°C led to change of the solution color from pale yellow to pale brown with increasing the absorption bands around 350 nm (Figure 2; red spectrum). Although the acetate complex **1** also reacted with O_2 , a resulting blue–green-colored species exhibited $\lambda_{\text{max}}=700\text{ nm}$, which would be attributed from




Figure 2. UV/Vis spectra of **2**, **3**, **4**, and **5** (0.5 mm) in THF at -60°C . Complexes **3** and **5** were prepared from **2** by in situ addition of O_2 gas (1 atm) and subsequent addition of AZADOL (50 equiv), respectively. Complex **4** was prepared from **2** by in situ addition of CO gas (1 atm).

a dinuclear iron(III)– μ -peroxo species analogous to the previously reported Tp^{Pr_2} derivatives.^[13a,b] The observed spectral pattern of the pale brown species **3** is similar to those of the synthetic complexes such as the recently characterized mononuclear non-heme iron(III)–superoxo species with a pentadentate ligand (330 nm)^[5a] and a terminal superoxide complex of iron(III) (325 nm)^[8] found in a oxygenation of the dinuclear iron(II) complex rather than that of the most recently reported mononuclear side-on iron (III)–superoxo species (490 nm).^[5b] Upon bubbling of Ar gas at low temperature (-60°C) or raising the temperature to -20°C , the pale brown color of **3** disappeared to recover the precursor **2**. Repetitious O_2 and Ar bubbling showed reversible change of the UV/Vis spectra corresponding to the addition and dissociation of O_2 to the iron center of **2**. The oxygenation process was also observed by low-temperature ^1H NMR spectroscopy in $[\text{D}_8]\text{THF}$ solution (Supporting Information, Figures S13, S14). Under an argon atmosphere, paramagnetically shifted signals of **2** were observed. On the basis of Evans method analysis, an iron(II) center of **2** in THF solution has $S=2$ spin configuration as expected from its pseudo-square pyramidal geometry determined by X-ray crystallography. Upon introduction of O_2 into this NMR sample, major signals were observed only in the 0–10 ppm region.^[14] The formed O_2 adduct **3** might be a diamagnetic species in which a low-spin iron(III) center ($S=1/2$) was antiferromagnetically coupled with superoxide anion radical ($S=1/2$). Such electronic configuration is found in the O_2 adduct of heme–iron compounds^[2,15] and Fe–Bleomycin.^[3,16] Furthermore, the high-spin ferrous center of **2** reacted with CO to give the corresponding low-spin iron(II)–carbonyl complex **4**, which exhibited characteristic features such as the $\nu(\text{C}=\text{O})$ band at 1967 cm^{-1} in the IR and diamagnetic signals in the ^1H NMR spectra (Figures S4, S15).^[17] Such behavior of **2** is also consistent with that of the iron(II) centers in the heme compounds and Fe–Bleomycin.

A resonance Raman spectrum of **3** generated in situ (derived by reaction of a THF solution of **2** with $^{16}\text{O}_2$) recorded at -60°C with 413 nm excitation exhibited an isotope sensitive Fermi doublet band at 1168 cm^{-1} (Table 1; Figure S6). This band was shifted to 1090 cm^{-1} on the $^{18}\text{O}_2$ derivative. These values are similar to those reported for O–O stretching frequencies of the metal–superoxide complexes including mononuclear non-heme iron(III) species.^[1] Furthermore, Fe–O stretching bands were observed at 592 cm^{-1} on the $^{16}\text{O}_2$ derivative and 568 cm^{-1} on the $^{18}\text{O}_2$ derivative,

Table 1: O–O and M–O stretching frequencies [cm^{-1}].

Compound	$\nu(\text{O}=\text{O})$ ($\Delta^{18}\text{O}$)	$\nu(\text{M}=\text{O})$ ($\Delta^{18}\text{O}$)
3	1168 ^[a] (78)	592 (24)
$[\text{Fe}(\text{BDPP})(\text{O}_2)]^{[b]}$	1125 (63)	not reported
$[\text{Fe}(\text{TAML})(\eta^2\text{-O}_2)]^{[b]}$	1260 (77)	not reported
3 _{Co,Ph}	1150 (60)	543 (21)
5	778 (40)	598 (25)
Heme $\text{Fe}^{\text{III}}\text{-OOH}^{[c]}$	799–810 (39–47)	566–617 (25–27)
LS $\text{Fe}^{\text{III}}\text{-OOH}^{[d]}$	784–811 (31–51)	609–632 (16–26)
HS $[\text{Fe}^{\text{III}}(\text{TMC})(\text{OOH})]^{[e]}$	868 (48)	658 (25)


[a] Center of Fermi doublet. [b] Ref. [5]. [c] Ref. [18]. [d] Ref. [3b].

[e] Ref. [20a].

respectively.^[18,19] These features suggest that **3** is a mononuclear iron(III)–superoxo species.

We did not meet with success in the isolation and structure determination of **3** so far. Therefore, we explored the synthesis and structural characterization of cobalt analogues. Mononuclear cobalt(II) complexes, $[\text{Co}^{\text{II}}(\text{L}^{\text{X}})(\text{Tp}^{\text{Me}_2})]$ (**2**^{Co,X}; X = Ph and OiPr), also reacted with O₂ reversibly (Figures S1, S17, S18). The resulting O₂ adducts **3**^{Co,X} exhibited O–O stretching bands at 1150 cm^{−1} for the ¹⁶O₂ derivative and 1090 cm^{−1} for the ¹⁸O₂ derivative of **3**^{Co,Ph} observed by resonance Raman with 413 nm excitation, and 1147 cm^{−1} for the ¹⁶O₂ derivative and 1088 cm^{−1} for the ¹⁸O₂ derivative of **3**^{Co,OiPr} measured by IR, respectively (Figures S5, S7).^[19] EPR spectra of **3**^{Co,X} bearing a signal at about *g* = 2 with eight-line hyperfine coupling were characteristic of mononuclear low-spin cobalt(III)–superoxo species of which the spin state is *S* = 1/2 (Figure S11). Finally, the molecular structure of **3**^{Co,OiPr}, $[\text{Co}^{\text{III}}(\eta^1\text{-O}_2)(\text{L}^{\text{OiPr}})(\text{Tp}^{\text{Me}_2})]$, could be determined by X-ray crystallography. An O–O bond length (1.301(5) Å) is typical for superoxide. The superoxo ligand is coordinated to the sixth site of the cobalt center with end-on fashion (Scheme 2). On the basis of structural analogy of the iron(II) precursor **2** and the cobalt analogues **2**^{Co,X} as well as the similarity of *v*(O–O) values, the iron(III)–superoxo species **3** can be assigned to a mononuclear end-on O₂[−] complex, formulated as $[\text{Fe}^{\text{III}}(\eta^1\text{-O}_2)(\text{L}^{\text{Ph}})(\text{Tp}^{\text{Me}_2})]$. Furthermore, the relatively high *v*(Fe–O) frequency of **2** is similar to those observed on $\eta^1\text{-O}_2$ adducts of heme iron species.^[2a] Moreover, frequency of *v*(Fe–O) is committed to the coordination mode of peroxides in non-heme iron complexes. η^1 -hydroperoxo species shows a higher *v*(Fe–O) frequency (> 550 cm^{−1}) in comparison to η^2 -peroxo species (< 500 cm^{−1}).^[3b,18,20]

Investigation on H-abstraction ability revealed that **3** reacted with substrates having a weak X–H bond (where X = O or N; BDE of X–H < 72.6 kcal mol^{−1}) such as 2-hydroxy-2-azaadamantane (AZADOL) and phenylhydrazine. The reaction of **3** with these substrates afforded a new species assigned to an iron(III)–hydroperoxo adduct, $[\text{Fe}^{\text{III}}(\text{OOH})(\text{L}^{\text{Ph}})]$.



Scheme 2. Formation of superoxide complexes of cobalt analogues.^[23]

(Tp^{Me2}) (**5**), which exhibited a characteristic absorption band at 542 nm, and resonance Raman features at 778 and 598 cm^{−1} with 514 nm excitation corresponding to the vibration of the O–O and Fe–O bonds, respectively (Figures S2, S8, S9). The bands of this ¹⁶O₂ derivative were shifted to 738 and 573 cm^{−1} on the ¹⁸O₂ derivative of **5**. The observed values are close to those of *v*(O–O) and *v*(Fe–O) reported for the heme and non-heme low-spin ferric hydroperoxo species.^[3b,18,20] Thermal decomposition of complex **5** was confirmed by UV/Vis spectroscopy upon raising the temperature to −20 °C. Conversion of **3** to **5** was also observed by EPR spectroscopy (Figures S10, S11). A frozen THF solution of **3** was EPR-silent at 77 K. Addition of AZADOL at −60 °C led to the appearance of signals at the *g* = 2 region attributed to a rhombic low-spin iron(III) species with a N–O• radical of the resulting 2-oxyl-2-azaadamantane. Noteworthy, relatively large kinetic isotope effect was observed during the formation of **5** by using deuterated AZADOL (*k*_H/*k*_D ≈ 18 at 213 K) under low concentration of this substrate, and that suggests the O–H bond cleavage is involved in the rate-determining step (Supporting Information, Figure S2). However, the formation rates of **5** were saturated with increasing the concentration of AZADOL. Such kinetic behavior supports the existence of an equilibrium process for the formation of adduct of **3** with AZADOL. Unfortunately, **3** was unable to abstract hydrogen atoms from C–H bonds of alkanes, alkenes, and BNAH (1-benzyl-1,4-dihydronicotinamide), the last of which the C–H BDE is only 67.9 kcal mol^{−1}.

Furthermore, **3** was inert toward oxidation of exogenous electrophilic substrates including aldehydes and acyl halides at low temperature.^[21] Sluggishness of **3** in nucleophilic reaction might be attributed to the low electron density on the superoxo moiety owing to strong antiferromagnetic coupling with low-spin iron(III) center. In contrast to **3**, high-spin iron(III)–superoxo complexes are more reactive. $[\text{Fe}(\text{BDPP})(\text{O}_2)]$ abstracts H atom from 9,10-dihydroanthracene.^[5a] Although $[\text{Fe}(\text{TAML})(\eta^2\text{-O}_2)]$ (*S* = 3/2 on Fe) cannot activate hydrocarbon substrates with weak C–H bonds (for example, 1,4-cyclohexadiene), this side-on superoxo complex reacts with phenols and aldehydes showing its H-abstrating and nucleophilic reactivity, respectively.^[5b] Although 1e[−] reduction of **3** by permethylferrocene did not occur without any additive, **3** reacted with permethylferrocene in the presence of strong acid such as trifluoroacetic acid (Supporting Information, Figure S3). Accordingly, PCET or PT/ET would proceed on the superoxo ligand of **3**.^[22]

In summary, the mononuclear superoxo complex of non-heme low-spin iron(III) was characterized successfully. This low-spin ferric superoxo compound is converted into the corresponding low-spin ferric–hydroperoxo species. The iron(II) center of **2** is supported by the non-heme ligands Tp^{Me2} and L^{Ph}; however, its reactivity (the high-spin iron(II) center reacts with O₂ and CO to give the corresponding adducts of low-spin iron(III) and iron(II), respectively) is close to that of the heme iron systems as well as the non-heme ferrous center of Fe-Bleomycin. In Fe-Bleomycin, an equatorial pyrimidine ligand is critical to control the chemical properties of the iron center.^[3,16] In our system, strong electron donation from the equatorial locating L^{Ph} seems to

be responsible for the above described nature of the iron center. The square-pyramidal iron(II) supported by five azoles is found in the O₂-binding center of hemoglobin/myoglobin. Interestingly, reproducing such coordination environment results in mimicking the reversible O₂-binding/releasing rather than substrate oxidizing functions in a non-heme system. Further investigations including theoretical analysis are undergoing.

Keywords: bioinorganic chemistry · cobalt · dioxygen ligands · iron · O–O activation

How to cite: *Angew. Chem. Int. Ed.* **2015**, *54*, 7336–7339
Angew. Chem. **2015**, *127*, 7444–7447

- [1] a) K. Ray, F. F. Pfaff, B. Wang, W. Nam, *J. Am. Chem. Soc.* **2014**, *136*, 13942; b) A. S. Borovik, P. J. Zinn, M. K. Zart in *Activation of Small Molecules: Organometallic and Bioinorganic Perspectives* (Ed.: W. B. Tolman), Wiley-VCH, Weinheim, **2006**, pp. 187–234.
- [2] a) M. Sono, M. P. Roach, E. D. Coulter, J. H. Dawson, *Chem. Rev.* **1996**, *96*, 2841; b) J. P. Collman, R. Boulton, C. J. Sunderland, L. Fu, *Chem. Rev.* **2004**, *104*, 561; c) T. L. Poulos, *Chem. Rev.* **2014**, *114*, 3919.
- [3] a) E. I. Solomon, T. C. Brunold, M. I. Davis, J. N. Kemsley, S.-K. Lee, N. Lehnert, F. Neese, A. J. Skulan, Y.-S. Yang, J. Zhou, *Chem. Rev.* **2000**, *100*, 235; b) M. Costas, M. P. Mehn, M. P. Jensen, L. Que, Jr., *Chem. Rev.* **2004**, *104*, 939; c) D. Buongiorno, G. D. Straganz, *Coord. Chem. Rev.* **2013**, *257*, 541.
- [4] a) Y. Zhang, K. L. Colabroy, T. P. Begley, S. E. Ealick, *Biochemistry* **2005**, *44*, 7632; b) M. M. Mbughuni, M. Chakrabarti, J. A. Hayden, E. L. Bominaar, M. P. Hendrich, E. Münck, J. D. Lipscomb, *Proc. Natl. Acad. Sci. USA* **2010**, *107*, 16788; c) G. Xing, Y. Diao, L. M. Hoffart, E. W. Barr, K. S. Prabhu, R. J. Arner, C. C. Reddy, C. Krebs, J. M. Bollinger, Jr., *Proc. Natl. Acad. Sci. USA* **2006**, *103*, 6130; d) J. M. Bollinger, Jr., C. Krebs, *Curr. Opin. Chem. Biol.* **2007**, *11*, 151.
- [5] a) C.-W. Chiang, S. T. Kleespies, H. D. Stout, K. K. Meier, P.-Y. Li, E. L. Bominaar, L. Que, Jr., E. Münck, W.-Z. Lee, *J. Am. Chem. Soc.* **2014**, *136*, 10846; b) S. Hong, K. D. Sutherlin, J. Park, E. Kwon, M. A. Siegler, E. I. Solomon, W. Nam, *Nat. Commun.* **2014**, *5*, 5440.
- [6] S. Kundu, E. Matito, S. Walleck, F. F. Pfaff, F. Heims, B. Ráby, J. M. Luis, A. Company, B. Braun, T. Glaser, K. Ray, *Chem. Eur. J.* **2012**, *18*, 2787.
- [7] M. Zhao, B. Helms, E. Slonkina, S. Friedle, D. Lee, J. DuBois, B. Hedman, K. O. Hodgson, J. M. J. Fréchet, S. J. Lippard, *J. Am. Chem. Soc.* **2008**, *130*, 4352.
- [8] X. Shan, L. Que, Jr., *Proc. Natl. Acad. Sci. USA* **2005**, *102*, 5340.
- [9] E. D. Bloch, L. J. Murray, W. L. Queen, S. Chavan, S. N. Maximoff, J. P. Bigi, R. Krishna, V. K. Peterson, F. Grandjean, G. J. Long, B. Smit, S. Bordiga, C. M. Brown, J. R. Long, *J. Am. Chem. Soc.* **2011**, *133*, 14814.
- [10] In the following synthetic mononuclear non-heme iron complex systems, iron(III)–superoxo intermediates are suggested to be involved during the formation of iron(III)–hydroperoxo species: a) R. J. Guajardo, S. E. Hudson, S. J. Brown, P. K. Mascharak, *J. Am. Chem. Soc.* **1993**, *115*, 7971; b) Y.-M. Lee, S. Hong, Y. Morimoto, W. Shin, S. Fukuzumi, W. Nam, *J. Am. Chem. Soc.* **2010**, *132*, 10668; c) M. Martinho, G. Blain, F. Banse, *Dalton Trans.* **2010**, *39*, 1630; d) Q. Zhang, C. R. Goldsmith, *Inorg. Chem.* **2014**, *53*, 5206.
- [11] a) S. Trofimenko, *Scorpionates: The Coordination Chemistry of Polypyrazolylborate Ligands*, Imperial College Press, London, **1999**; b) C. Pettinari, *Scorpionates II: Chelating Borate Ligands*, Imperial College Press, London, **2008**.
- [12] a) K. Fujita, S. Hikichi, M. Akita, Y. Moro-oka, *J. Chem. Soc. Dalton Trans.* **2000**, 1255; b) S. Hikichi, K. Fujita, Y. Manabe, M. Akita, J. Nakazawa, H. Komatsuzaki, *Eur. J. Inorg. Chem.* **2010**, 5529.
- [13] a) N. Kitajima, N. Tamura, H. Amagai, H. Fukui, Y. Moro-oka, Y. Mizutani, T. Kitagawa, R. Mathur, K. Heerwegh, C. A. Reed, C. R. Randall, L. Que, Jr., K. Tatsumi, *J. Am. Chem. Soc.* **1994**, *116*, 9071; b) K. Kim, S. J. Lippard, *J. Am. Chem. Soc.* **1996**, *118*, 4914; c) E. H. Ha, R. Y. N. Ho, J. F. Kiesel, J. S. Valentine, *Inorg. Chem.* **1995**, *34*, 2265; d) A. Mukherjee, M. Martinho, E. L. Bominaar, E. Münck, L. Que, Jr., *Angew. Chem. Int. Ed.* **2009**, *48*, 1780; *Angew. Chem.* **2009**, *121*, 1812; e) A. Mukherjee, M. A. Cranswick, M. Chakrabarti, T. K. Paine, K. Fujisawa, E. Münck, L. Que, Jr., *Inorg. Chem.* **2010**, *49*, 3618; f) I. Siewert, C. Limberg, *Angew. Chem. Int. Ed.* **2008**, *47*, 7953; *Angew. Chem.* **2008**, *120*, 8071; g) H. Park, M. M. Bittner, J. S. Baus, S. V. Lindeman, A. T. Fiedler, *Inorg. Chem.* **2012**, *51*, 10279; h) M. Sallmann, I. Siewert, L. Fohlmeister, C. Limberg, C. Knispel, *Angew. Chem. Int. Ed.* **2012**, *51*, 2234; *Angew. Chem.* **2012**, *124*, 2277; i) S. Paria, S. Chatterjee, T. K. Paine, *Inorg. Chem.* **2014**, *53*, 2810.
- [14] The signal pattern of **3** is relatively similar to that of **4** except peak broadness, which may due to the presence of excess O₂ or the effect from the coupled spins.
- [15] H. Chen, M. Ikeda-Saito, S. Shaik, *J. Am. Chem. Soc.* **2008**, *130*, 14778.
- [16] a) R. M. Burger, *Chem. Rev.* **1998**, *98*, 1153; b) C. A. Claussen, E. C. Long, *Chem. Rev.* **1999**, *99*, 2797; c) S. V. Kryatov, E. V. Rybak-Akimova, *Chem. Rev.* **2005**, *105*, 2175; d) N. J. Oppenheimer, L. O. Rodriguez, S. M. Hecht, *Proc. Natl. Acad. Sci. USA* **1979**, *76*, 5616; e) S. Takahashi, J. W. Sam, J. Peisach, D. L. Rousseau, *J. Am. Chem. Soc.* **1994**, *116*, 4408; f) R. M. Burger, J. Peisach, S. B. Horwitz, *J. Biol. Chem.* **1981**, *256*, 11636; g) R. M. Burger, T. A. Kent, S. B. Horwitz, E. Münck, J. Peisach, *J. Biol. Chem.* **1983**, *258*, 1559.
- [17] R. K. Afshar, A. K. Patra, E. Bill, M. M. Olmstead, P. K. Mascharak, *Inorg. Chem.* **2006**, *45*, 3774.
- [18] T. Ohta, J.-G. Liu, Y. Naruta, *Coord. Chem. Rev.* **2013**, *257*, 407.
- [19] Isotope sensitive bands at 425 (**2**) and 384 cm⁻¹ (**2**^{Co,Ph}) can be assigned to M–O–O bending mode: S. Hirota, T. Ogura, E. H. Appleman, K. Shinzawa-Itoh, S. Yoshikawa, T. Kitagawa, *J. Am. Chem. Soc.* **1994**, *116*, 10564.
- [20] a) J. Cho, S. Jeon, S. A. Wilson, L. V. Liu, E. A. Kang, J. J. Braymer, M. H. Lim, B. Hedman, K. O. Hodgson, J. S. Valentine, E. I. Solomon, W. Nam, *Nature* **2011**, *478*, 502; b) G. Roelfes, V. Vrajmasu, K. Chen, R. Y. N. Ho, J.-U. Rohde, C. Zondervan, R. M. la Crois, E. P. Schudde, M. Lutz, A. L. Spek, R. Hage, B. L. Feringa, E. Münck, L. Que, Jr., *Inorg. Chem.* **2003**, *42*, 2639.
- [21] Nucleophilic reactivity of a copper(II)–superoxo complex has been reported: P. Pirovano, A. M. Magherusan, C. McGlynn, A. Ure, A. Lynes, A. R. McDonald, *Angew. Chem. Int. Ed.* **2014**, *53*, 5946; *Angew. Chem.* **2014**, *126*, 6056.
- [22] R. L. Peterson, J. W. Ginsbach, R. E. Cowley, M. F. Qayyum, R. A. Himes, M. A. Siegler, C. D. Moore, B. Hedman, K. O. Hodgson, S. Fukuzumi, E. I. Solomon, K. D. Karlin, *J. Am. Chem. Soc.* **2013**, *135*, 16454.
- [23] CCDC 1053457 (**2**), 1053458 (**2**^{Co,Ph}), 1053459 (**2**^{Co,OPr}), and 1053460 (**3**^{Co,OPr}) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

Received: March 13, 2015

Published online: May 4, 2015