

# Intramolecular Aminoboration of Unfunctionalized Olefins


Chun-Hua Yang, Yu-Shi Zhang, Wen-Wen Fan, Gong-Qing Liu, and Yue-Ming Li\*

**Abstract:** A direct and catalyst-free method for the intramolecular aminoboration of unfunctionalized olefins is reported. In the presence of  $\text{BCl}_3$  (1 equiv) as the sole boron source, intramolecular aminoboration of sulfonamide derivatives of 4-penten-1-amines, 5-hexen-1-amines, and 2-allylanilines proceeded readily without the use of any catalyst. The boronic acids obtained after hydrolysis could be converted into the corresponding pinacol borates in a straightforward manner by treatment with pinacol under anhydrous conditions.


Organoboron compounds have found widespread application in a variety of carbon–carbon and carbon–heteroatom coupling reactions.<sup>[1]</sup> Furthermore, boronic acid moieties have emerged as important functional groups in biologically active compounds. For example, an  $\alpha$ -aminoboronic acid is a key structure in the proteasome inhibitor Velcade (bortezomib),<sup>[2]</sup> and  $\beta$ -aminoboronic acids have been used as a key subunit in peptidomimetics with antitubercular activity.<sup>[3]</sup>

Traditionally, organoboron compounds have been prepared by transmetalation reactions between organometallic compounds, such as organolithium and organomagnesium reagents, and borates,<sup>[4]</sup> by the hydroboration of alkenes and alkynes,<sup>[5]</sup> by the haloboration or aminoboration of alkynes,<sup>[6]</sup> by the borylation of  $\text{C}=\text{X}$  double bonds,<sup>[7]</sup> by the borylation of  $\text{C}-\text{H}/\text{C}-\text{X}$  bonds,<sup>[8]</sup> and by other miscellaneous methods.<sup>[9]</sup> Recently, Hirano, Miura, and co-workers<sup>[10]</sup> and Tortosa and co-workers<sup>[11]</sup> described the copper(I)-catalyzed simultaneous addition of nitrogen and boron atoms to  $\text{C}-\text{C}$  multiple bonds or their equivalents for the preparation of aminoboron compounds. In the presence of a catalytic amount of  $\text{CuCl}$ , the intermolecular aminoboration of olefins with bis(pinacolato)diboron and *O*-benzoyl *N,N*-dialkyl hydroxylamines produced the corresponding  $\beta$ -aminoboron compounds in good to excellent yields (Scheme 1a). We are interested in developing new, straightforward procedures for the aminoboration of  $\text{C}-\text{C}$  multiple bonds. Herein, we report our recent results on the direct and catalyst-free intramolecular aminoboration of unfunctionalized  $\text{C}=\text{C}$  double bonds (Scheme 1b).

a)  $\text{Cu}^{\text{I}}$ -catalyzed intermolecular aminoboration (previously reported studies)




b) Direct intramolecular aminoboration (this study)



**Scheme 1.** Inter- and intramolecular aminoboration reactions.

Bz = benzoyl, pin = pinacolato.

In the course of searching for new methods for the amination of  $\text{C}-\text{C}$  multiple bonds, we found that the intramolecular fluoroamination of unfunctionalized olefins was possible with  $\text{BF}_3$  as the fluorine source and  $\text{PhI}(\text{OAc})_2$  as the reaction promoter.<sup>[12]</sup> We assumed that the direct aminoboration of  $\text{C}=\text{C}$  double bonds should be possible if an  $\text{N}-\text{B}$  intermediate could be generated during the reaction. In this context, an aminoboration reaction was proposed with  $\text{BF}_3 \cdot \text{Et}_2\text{O}$  as the boron source. However, no desired aminoboration product was detected when **1a** was treated with  $\text{BF}_3 \cdot \text{Et}_2\text{O}$ . Instead, a very slow intramolecular hydroamination reaction occurred, which was complete after 24 h at  $60^\circ\text{C}$  (Scheme 2).<sup>[13]</sup>




**Scheme 2.** Amination reactions involving  $\text{BF}_3$ . Ts = *p*-toluenesulfonyl.

We reasoned that  $\text{BF}_3$  would interact with the sulfonamide substrate **1a** to form a Lewis pair with either the nitrogen or oxygen atom of the sulfonamido group. The formation of an  $\text{N}-\text{BF}_2$  bond is difficult owing to the high dissociation energy of  $\text{B}-\text{F}$  bonds. Instead, Lewis acid mediated reactions took place, and the hydroamination product was isolated. Given that the dissociation energy of a  $\text{B}-\text{Cl}$  bond is lower than that of a  $\text{B}-\text{F}$  bond, we reasoned that different reactions should take place if  $\text{BCl}_3$  was used instead of  $\text{BF}_3$  under similar reaction conditions. To test this assumption, we added a stoichiometric amount of  $\text{BCl}_3$  to a solution of **1a**. After 12 h at room temperature, the desired aminoboration product **2a** was obtained (Scheme 3). The structure of **2a** was confirmed by X-ray diffraction.

[\*] C.-H. Yang, Y.-S. Zhang, W.-W. Fan, Dr. G.-Q. Liu, Prof. Dr. Y.-M. Li  
Stake Key Laboratory of Medicinal Chemical Biology, College of Pharmacy and Tianjin Key Laboratory of Molecular Drug Research  
Nankai University  
Tianjin 300071 (P.R. China)  
E-mail: ymli@nankai.edu.cn  
Prof. Dr. Y.-M. Li  
CAS Key Laboratory of Synthetic Chemistry of Natural Substances  
Shanghai Institute of Organic Chemistry  
Chinese Academy of Sciences  
345 Lingling Road, Shanghai 200032 (P.R. China)

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/ange.201505489>.



**Scheme 3.** Direct intramolecular aminoboration of **1a**.

Encouraged by this result, we optimized the conditions for this direct intramolecular aminoboration reaction. Preliminary results indicated that the reaction medium was crucial for the reaction. Haloalkane solvents were generally suitable, and 1,2-dichloroethane (DCE) gave the most promising result. This solvent was then used as the reaction medium for further studies. Polar aprotic solvents, such as THF, 1,4-dioxane, DMF, and acetonitrile, were not suitable for the reaction, possibly as a result of their strong interaction with  $\text{BCl}_3$  and subsequent deactivation of the latter. When the reaction was carried out in toluene, the expected aminoboration product was also obtained. However, the removal of toluene was difficult as compared to the removal of DCE, and reaction in toluene was not pursued further. No significant rate difference was observed when the reaction was carried out at 30 °C instead of 25 °C. Therefore, subsequent reactions were carried out at room temperature without special control of the reaction temperature. A reaction with  $\text{BBr}_3$  as the boron source gave a similar result. Studies with  $\text{BBr}_3$  were not continued owing to the relatively difficult conditions needed to handle the reagent.

Having established optimal reaction conditions, we also studied the effect of the protecting group on the course of the reaction. We found that the basicity and nucleophilicity of the nitrogen atom were crucial factors. When the amino group was protected with a tosyl group, the aminoboration product was obtained in good yield. Other protecting groups, such as benzyl, acetyl, trifluoroacetyl, and *tert*-butoxycarbonyl (Boc) groups, were not suitable, and no aminoboration product was detected when substrates bearing these protecting groups were subjected to the reaction. Deprotection occurred with the isolation of the corresponding primary amine when a Boc-protected substrate was used. These results indicated that for the aminoboration to proceed, the amino/amido group must have balanced basicity/nucleophilicity. Strongly nucleophilic amino/amido groups will interact more strongly with  $\text{BCl}_3$ , thus leading to the deactivation of both the nitrogen and the boron atom.

As the purification of boronic acids on a silica-gel column is sometimes problematic, the obtained boronic acids were converted into the corresponding boronates<sup>[14]</sup> to facilitate product purification. A variety of *N*-(4-pentenyl)sulfonamide substrates **1** were subjected to the intramolecular aminoboration reaction. After complete consumption of the starting material, the reaction mixture was carefully treated with water, and the crude product was separated from the mixture and treated with pinacol and magnesium sulfate to give the corresponding pinacol boronate. Reactions of *p*-toluene-, methane-, 2-nitrobenzene-, and 4-nitrobenzenesulfonamide substrates all proceeded readily, and the corresponding

boronates were isolated in good overall yields (Table 1). The reaction was not affected significantly by the Thorpe–Ingold effect (compare the formation of products **3a**, **3f**, **3h**,

**Table 1:** Direct intramolecular aminoboration of unfunctionalized *N*-(4-pentenyl)sulfonamides.<sup>[a]</sup>

Entry	R <sup>1</sup>	R <sup>2</sup>	Product	Yield [%] <sup>[b]</sup>
1	Ph	Ts	<b>3a</b>	74
2	Ph	Ms	<b>3b</b>	86
3	Ph	<i>o</i> -NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub> SO <sub>2</sub>	<b>3c</b>	44
4	Ph	<i>p</i> -NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub> SO <sub>2</sub>	<b>3d</b>	84
5			<b>3e</b>	86
6	-(CH <sub>2</sub> ) <sub>5</sub> -	Ts	<b>3f</b>	83
7	-(CH <sub>2</sub> ) <sub>5</sub> -	Ms	<b>3g</b>	45
8	Me	Ts	<b>3h</b>	61
9	Me	Ms	<b>3i</b>	44
10	Me	<i>p</i> -NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub> SO <sub>2</sub>	<b>3j</b>	68
11	H	Ts	<b>3k</b>	55
12	H	PhSO <sub>2</sub>	<b>3l</b>	63
13	H	<i>p</i> -NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub> SO <sub>2</sub>	<b>3m</b>	40


[a] Reactions conditions: **1** (1 mmol),  $\text{BCl}_3$  (1 mmol), DCE (10 mL), argon atmosphere. [b] Yield of the isolated product. Ms = methanesulfonyl.

and **3k**).<sup>[15]</sup> Substrates with substituents on the main chain, such as 2,2-diphenyl (substrate **1a**),  $-(\text{CH}_2)_5-$  (substrate **1f**), or 2,2-dimethyl (substrate **1h**), were transformed efficiently into the desired aminoboration products, and the aminoboration of substrates without substituents on the main chain (substrates **1k–m**) also gave the corresponding aminoboration products in satisfactory yields (Table 1, entries 11–13). Substrate **1e** did not undergo the aminoboration reaction. Instead, the Friedel–Crafts alkylation product **3e** was isolated in high yield (Table 1, entry 5).<sup>[13]</sup> A gram-scale synthesis of **2a** was also carried out to test the scalability of the method: The aminoboration of **1a** on a 5.0 mmol scale afforded boronic acid **2a** in 83 % yield. Furthermore, functional-group transformations by oxidation, amination, and Suzuki coupling reactions were possible.<sup>[16]</sup>

We next turned our attention to the intramolecular aminoboration of *N*-(5-hexenyl)sulfonamide substrates **4**. Reactions of *N*-(5-hexenyl)sulfonamide substrates **4** generally proceeded less efficiently than those of *N*-(4-pentenyl)sulfonamide substrates **1**, possibly as a result of the disfavored entropic nature of the cyclization reactions.<sup>[17]</sup> However, aminoboration products **5** were still isolated in moderate to good yields under the optimized conditions (Table 2).

Following the synthesis of [(4,4-diphenyl-1-tosylpyrrolidin-2-yl)methyl]boronic acid (**2a**), 2-(4,4,5,5-tetramethyl-1,3,2]dioxaborolan-2-ylmethyl]pyrrolidines (**3**), and 2-(4,4,5,5-tetramethyl-1,3,2]dioxaborolan-2-ylmethyl]piperi-

**Table 2:** Direct intramolecular aminoboration of unfunctionalized *N*-(5-hexenyl)sulfonamides.<sup>[a]</sup>




Entry	R <sup>1</sup>	R <sup>2</sup>	Product	Yield [%] <sup>[b]</sup>
1	Ph	Ts	<b>5 a</b>	58
2	Ph	Ms	<b>5 b</b>	70
3	Ph	<i>p</i> -NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub> SO <sub>2</sub>	<b>5 c</b>	50
4	-(CH <sub>2</sub> ) <sub>5</sub> -	Ts	<b>5 d</b>	54
5	-(CH <sub>2</sub> ) <sub>5</sub> -	Ms	<b>5 e</b>	65
6	-(CH <sub>2</sub> ) <sub>5</sub> -	<i>p</i> -NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub> SO <sub>2</sub>	<b>5 f</b>	67
7	Me	Ts	<b>5 g</b>	86
8	Me	Ms	<b>5 h</b>	56
9	Me	<i>o</i> -NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub> SO <sub>2</sub>	<b>5 i</b>	61
10	H	Ts	<b>5 j</b>	63
11	H	PhSO <sub>2</sub>	<b>5 k</b>	76
12	H	<i>p</i> -NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub> SO <sub>2</sub>	<b>5 l</b>	54

[a] Reaction conditions: **4** (1 mmol), BCl<sub>3</sub> (1 mmol), DCE (10 mL), argon atmosphere. [b] Yield of the isolated product.

dines (**5**), we also investigated the synthesis of different 2-(4,4,5,5-tetramethyl[1,3,2]dioxaborolan-2-ylmethyl)-2,3-dihydro-1*H*-indoles **7** by this method. The desired aminoboration products **7 a–f** were isolated in moderate overall yields, and the reaction was not overly sensitive to the electronic effect of substituents on the benzene ring (Table 3).

**Table 3:** Direct intramolecular aminoboration of *N*-(2-allylphenyl)-sulfonamides.<sup>[a]</sup>




Entry	R <sup>1</sup>	R <sup>2</sup>	Product	Yield [%] <sup>[b]</sup>
1	H	Ts	<b>7 a</b>	78
2	Me	Ts	<b>7 b</b>	69
3	MeO	Ts	<b>7 c</b>	29
4	Cl	Ts	<b>7 d</b>	41
5	Me	<i>p</i> -NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub> SO <sub>2</sub>	<b>7 e</b>	51
6	Cl	<i>p</i> -NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub> SO <sub>2</sub>	<b>7 f</b>	67

[a] Reaction conditions: **6** (1 mmol), BCl<sub>3</sub> (1 mmol), DCE (10 mL), argon atmosphere. [b] Yield of the isolated product.

The formation of the Friedel–Crafts alkylation product **3 e** rather than the corresponding aminoboration product when the trisubstituted substrate **1 e** was subjected to the reaction (Table 1) indicated the formation of a carbenium ion intermediate. The carbenium cation could be formed by protonation of the substituted C=C double bond with HCl when the corresponding carbenium cation intermediate showed enough stability. Reactions of terminal alkenes could proceed through the aminoboration pathway since the formation of the corresponding secondary carbenium cation would be slow and less favored.

We carried out control experiments to study the possible formation of HCl during the aminoboration reaction. Two reactions were examined under the standard aminoboration conditions, and sodium carbonate was added to one of the two reaction mixtures. The reactions were carried out in an ice–water bath and were sampled every 10 min. The study showed that reactions in the presence of Na<sub>2</sub>CO<sub>3</sub> proceeded slightly faster than reactions without the base (Table 4). This result

**Table 4:** Control experiments with and without a base.<sup>[a]</sup>




Entry	Reaction time [min]	Yield [%] <sup>[b,c]</sup>	
		A	B
1	10	43	47
2	20	48	55
3	30	53	58
4	40	57	61
5	50	61	66
6	60	64	72

[a] Reactions were carried out with **1 a** (0.5 mmol) and BCl<sub>3</sub> (0.5 mmol) in CDCl<sub>3</sub> (5 mL), with or without Na<sub>2</sub>CO<sub>3</sub>, in an ice–water bath. [b] The yield of **2 a** was determined by NMR spectroscopy with 1,3,5-trimethoxybenzene (0.33 mmol) as the internal standard. [c] Conditions A: The reaction was carried out in the absence of Na<sub>2</sub>CO<sub>3</sub>. Conditions B: The reaction was carried out in the presence of Na<sub>2</sub>CO<sub>3</sub>.

possibly supports the assumption of the formation of HCl during the reaction. Removal of HCl from the reaction system should help to move the equilibrium to the N–BCl<sub>2</sub> side and speed up the reaction.

NMR spectroscopic experiments were also carried out to study the interaction between BCl<sub>3</sub> and the sulfonamide functional group. As a direct NMR spectroscopic study on the interaction between the substrate and BCl<sub>3</sub> was difficult owing to the fast conversion of the starting material into the aminoboration product, *N*-methyl-*p*-toluenesulfonamide (TsNHMe) was used as a model substrate. <sup>1</sup>H NMR spectra showed that the sulfonamide signals changed significantly after the addition of BCl<sub>3</sub>. Similarly, the <sup>11</sup>B NMR signal of BCl<sub>3</sub> changed significantly after the addition of TsNHMe. We reasoned that this change was due to the strong interaction between BCl<sub>3</sub> and the sulfonamide. DFT calculations were also carried out. Preliminary results indicated that the departure of HCl from the Lewis adduct was the rate-limiting step. However, this step was energetically favored both kinetically and thermodynamically. The intramolecular aminoboration step was found to be a fast step with a low energy barrier.<sup>[16]</sup>

On the basis of previously reported results and these preliminary studies, we propose a possible reaction pathway for the intramolecular aminoboration reaction in Scheme 4: When BCl<sub>3</sub> is mixed with the substrate, it interacts with the sulfonamide nitrogen atom to form an LA⋯NHRTs adduct **A**. The elimination of HCl from **A** to give an N–BCl<sub>2</sub> intermediate **B** is the rate-limiting step. It is also a fast step,



**Scheme 4.** A tentative reaction pathway for the intramolecular aminoboration reaction.

and the addition of a base had little effect on the course of the reaction. A downfield shift of the  $^1\text{H}$  NMR signals for the methylene hydrogen atoms adjacent to the nitrogen atom was observed owing to the attachment of an electron-withdrawing boron group to the nitrogen atom. Intramolecular aminoboration similar to a hydroboration reaction then occurred to give product **C**, which could be hydrolyzed to afford the final aminoboration product **D**. We propose that when the substituted substrate **1e** was used, protonation of the  $\text{C}=\text{C}$  bond by the  $\text{HCl}$  generated in situ led to the formation of a carbenium cation intermediate, which was captured by the phenyl group to yield the Friedel–Crafts alkylation product **3e**.<sup>[13]</sup>

In summary, we have described the direct intramolecular aminoboration of a variety of *N*-(4-pentenyl)-, *N*-(5-hexenyl)-, and *N*-(2-allylphenyl)sulfonamide substrates by treatment with  $\text{BCl}_3$ . In the presence of  $\text{BCl}_3$  (1 equiv) as the sole boron source, direct intramolecular aminoboration of these unfunctionalized olefins proceeded readily at room temperature without the use of a catalyst to give the corresponding boronic acids in good yields. The boronic acids could be readily converted into boronates, and the obtained boronates could be further functionalized by oxidation, amination, and Suzuki coupling reactions. The good yields, mild reaction conditions, and straightforward reaction procedure make this transformation an attractive method for the synthesis of a variety of useful *N*-heterocyclic boronic acids and boronates.

## Acknowledgements

We acknowledge financial support from the National Natural Science Foundation of China (NSFC 20972072, NSFC 21272121).

**Keywords:** aminoboration · boronic acids · boronic esters · Suzuki cross-coupling · synthetic methods

**How to cite:** *Angew. Chem. Int. Ed.* **2015**, *54*, 12636–12639  
*Angew. Chem.* **2015**, *127*, 12827–12830

- [1] a) N. Miyaura, A. Suzuki, *Chem. Rev.* **1995**, *95*, 2457–2483; b) A. J. J. Lennox, G. C. Lloyd-Jones, *Chem. Soc. Rev.* **2014**, *43*, 412–443; c) S. R. Chemler, D. Trauner, S. J. Danishefsky, *Angew. Chem. Int. Ed.* **2001**, *40*, 4544–4568; *Angew. Chem.* **2001**, *113*, 4676–4701; d) H. Doucet, *Eur. J. Org. Chem.* **2008**, 2013–2030; e) A. Suzuki, *Angew. Chem. Int. Ed.* **2011**, *50*, 6722–6737; *Angew. Chem.* **2011**, *123*, 6854–6869.

- [2] J. Adams, M. Kauffman, *Cancer Invest.* **2004**, *22*, 304–311.  
[3] A. S. Gorovoy, O. V. Gozhina, J. S. Svendsen, A. A. Domorad, G. V. Tetz, V. V. Tetz, T. Lejon, *Chem. Biol. Drug Des.* **2013**, *81*, 408–413.  
[4] a) H. C. Brown, T. E. Cole, *Organometallics* **1983**, *2*, 1316–1319; b) T. Ishiyama, M. Murata, N. Miyaura, *J. Org. Chem.* **1995**, *60*, 7508–7510.  
[5] a) H. C. Brown, S. K. Gupta, *J. Am. Chem. Soc.* **1971**, *93*, 1816–1818; b) H. C. Brown, S. K. Gupta, *J. Am. Chem. Soc.* **1972**, *94*, 4370–4371; c) H. C. Brown, N. G. Bhat, V. Somayaji, *Organometallics* **1983**, *2*, 1311–1316; d) I. D. Gridnev, N. Miyaura, A. Suzuki, *Organometallics* **1993**, *12*, 589–592; e) H. Yoshida, I. Kageyuki, K. Takaki, *Org. Lett.* **2014**, *16*, 3512–3515.  
[6] a) Y. Satoh, H. Serizawa, N. Miyaura, S. Hara, A. Suzuki, *Tetrahedron Lett.* **1988**, *29*, 1811–1814; b) C. Mazal, M. Vaultier, *Tetrahedron Lett.* **1994**, *35*, 3089–3090; c) E. Chong, S. A. Blum, *J. Am. Chem. Soc.* **2015**, *137*, 10144–10147.  
[7] a) M. A. Beenen, C. An, J. A. Ellman, *J. Am. Chem. Soc.* **2008**, *130*, 6910–6911; b) N. Hu, G. Zhao, Y. Zhang, X. Liu, G. Li, W. Tang, *J. Am. Chem. Soc.* **2015**, *137*, 6746–6749.  
[8] a) M. A. Larsen, J. F. Hartwig, *J. Am. Chem. Soc.* **2014**, *136*, 4287–4299; b) T. C. Atack, R. M. Lecker, S. P. Cook, *J. Am. Chem. Soc.* **2014**, *136*, 9521–9523.  
[9] a) D. S. Matteson, R. J. Moody, *J. Org. Chem.* **1980**, *45*, 1091–1095; b) T. Ishiyama, N. Matsuda, N. Miyaura, A. Suzuki, *J. Am. Chem. Soc.* **1993**, *115*, 11018–11019.  
[10] a) N. Matsuda, K. Hirano, T. Satoh, M. Miura, *J. Am. Chem. Soc.* **2013**, *135*, 4934–4937; b) R. Sakae, N. Matsuda, K. Hirano, T. Satoh, M. Miura, *Org. Lett.* **2014**, *16*, 1228–1231; c) R. Sakae, K. Hirano, T. Satoh, M. Miura, *Angew. Chem. Int. Ed.* **2015**, *54*, 613–617; *Angew. Chem.* **2015**, *127*, 623–627; d) R. Sakae, K. Hirano, M. Miura, *J. Am. Chem. Soc.* **2015**, *137*, 6460–6463.  
[11] A. Parra, L. Amenós, M. Guisán-Ceinos, A. López, J. L. García Ruano, M. Tortosa, *J. Am. Chem. Soc.* **2014**, *136*, 15833–15836.  
[12] G.-Q. Liu, Y.-M. Li, *J. Org. Chem.* **2014**, *79*, 10094–10109.  
[13] C.-H. Yang, W.-W. Fan, G.-Q. Liu, L. Duan, L. Li, Y.-M. Li, *RSC Adv.* **2015**, *5*, 61081–61093.  
[14] a) B. M. Partridge, L. Chausset-Boissarie, M. Burns, A. P. Pulis, V. K. Aggarwal, *Angew. Chem. Int. Ed.* **2012**, *51*, 11795–11799; *Angew. Chem.* **2012**, *124*, 11965–11969; b) A. P. Pulis, D. J. Blair, E. Torres, V. K. Aggarwal, *J. Am. Chem. Soc.* **2013**, *135*, 16054–16057.  
[15] a) P. G. Sammes, D. J. Weller, *Synthesis* **1995**, 1205–1222; b) J. Kaneti, A. J. Kirby, A. H. Koedjikov, I. G. Pojarlieff, *Org. Biomol. Chem.* **2004**, *2*, 1098–1103; c) M. E. Jung, G. Piizzi, *Chem. Rev.* **2005**, *105*, 1735–1766; d) M. R. Crimmin, M. Arrowsmith, A. G. M. Barrett, I. J. Casely, M. S. Hill, P. A. Procopiou, *J. Am. Chem. Soc.* **2009**, *131*, 9670–9685; e) N. T. Patil, V. S. Raut, R. D. Kavthé, V. V. N. Reddy, P. V. K. Raju, *Tetrahedron Lett.* **2009**, *50*, 6576–6579; f) J. Kostal, W. L. Jorgensen, *J. Am. Chem. Soc.* **2010**, *132*, 8766–8773; g) C. Duncan, A. V. Biradar, T. Asefa, *ACS Catal.* **2011**, *1*, 736–750.  
[16] Please refer to the Supporting Information for details.  
[17] a) G. Illuminati, L. Mandolini, *Acc. Chem. Res.* **1981**, *14*, 95–102; b) A. J. Kirby in *Advances in Physical Organic Chemistry*, Vol. 17 (Eds.: V. Gold, D. Bethell), Academic Press, London, New York, Toronto, Sydney, San Francisco, **1981**, pp. 183–278; c) T. G. Denisova, E. T. Denisov, *Russ. Chem. Bull.* **2002**, *51*, 949–960; d) J.-S. Ryu, T. J. Marks, F. E. McDonald, *J. Org. Chem.* **2004**, *69*, 1038–1052; e) A. Dzudza, T. J. Marks, *Chem. Eur. J.* **2010**, *16*, 3403–3422.

Received: June 15, 2015

Revised: August 4, 2015

Published online: September 1, 2015