

Cuprous Oxide Catalyzed Oxidative C–C Bond Cleavage for C–N Bond Formation: Synthesis of Cyclic Imides from Ketones and Amines


Min Wang, Jianmin Lu, Jiping Ma, Zhe Zhang, and Feng Wang*

Abstract: Selective oxidative cleavage of a C–C bond offers a straightforward method to functionalize organic skeletons. Reported herein is the oxidative C–C bond cleavage of ketone for C–N bond formation over a cuprous oxide catalyst with molecular oxygen as the oxidant. A wide range of ketones and amines are converted into cyclic imides with moderate to excellent yields. In-depth studies show that both α -C–H and β -C–H bonds adjacent to the carbonyl groups are indispensable for the C–C bond cleavage. DFT calculations indicate the reaction is initiated with the oxidation of the α -C–H bond. Amines lower the activation energy of the C–C bond cleavage, and thus promote the reaction. New insight into the C–C bond cleavage mechanism is presented.

The selective cleavage of C–C bonds offers a straightforward method for functionalizing the organic skeletons, and thus shows promising applications in organic synthesis and biomass conversion.^[1] But the nonpolar, thermodynamically stable, and kinetically inert character of C–C bonds makes it a challenge to increasing product selectivity.^[2] Nowadays chemical processes are being upgraded from traditional uncatalyzed reactions with stoichiometric oxidants, such as peroxides and metal salts,^[3] to catalyzed reactions using molecular oxygen.^[4]

Cyclic imides are widely used in biological, medicinal, and polymer chemistry.^[5] They have been synthesized by heating dicarboxylic acids or anhydrides with an amine (Scheme 1 a).^[6] In general, harsh thermal reaction conditions and activation reagents are mandatory. Recently studied oxidation of cyclic amines and carbonylation reactions have broadened the substrate scope using precious metal and carbonyl catalysts (Scheme 1 b and c).^[7]

Herein we report a new reaction for the synthesis of cyclic imides by oxidative coupling of a cyclic ketone with an amine (Scheme 1 d). To obtain good imide selectivity, each activation steps in the reaction, including C–H bond, C–C bond, and O=O bond (molecular oxygen) activation, should be kept selective over one catalyst.^[8] Very recently, we found that Cu₂O can activate oxygen and more interestingly, it enables the selective cleavage of strong C–C bonds over weak C–N bonds in tertiary amines.^[9] This ability indicates that the Cu₂O



Scheme 1. The methods for the synthesis of the cyclic imide. TBHP = *tert*-butyl hydroperoxide.

catalyst is unique for C–C bond-cleavage reactions. Inspired by this result, we discovered that Cu₂O catalyzed the C–C bond cleavage of ketones to synthesize cyclic imides. The method can be used in synthesizing various cyclic imides from anilines, benzylamines, aliphatic, and heterocyclic amines. Moreover, the combination of experiments and DFT results has revealed that the presence of both α -C–H and β -C–H bonds are essential for C–C bond cleavage, a transient carbanion intermediate is involved, and that the α -C–H bond is initially oxidized. To the best of our knowledge, this is the first report on conversion of ketones into cyclic imides.

We commenced our study with 1-indanone and aniline in the presence of a Cu₂O catalyst, which can either be obtained from a commercial source or prepared according to the literature.^[10] The solvent polarity remarkably affects the catalytic performance. Reactions in DMSO show higher activity (Table 1, entries 1–7). The catalyst was reused and gave a comparably good result (entry 7), and no reaction occurred without either oxygen or the catalyst (entries 8 and 9).

The substrate scope of the amines was then tested (Table 2). Under the optimized reaction conditions, a wide

[*] Dr. M. Wang, Dr. J. M. Lu, Dr. J. P. Ma, Z. Zhang, Prof. F. Wang
State Key Laboratory of Catalysis (SKLC)
Dalian National Laboratory for Clean Energy (DNL)
Dalian Institute of Chemical Physics (DICP)
Chinese Academy of Sciences, Dalian 116023 (China)
E-mail: wangfeng@dicp.ac.cn

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/ange.201508071>.

Table 1: Screening of the reaction conditions.^[a]

Entry	Solvent	Imide [%]
1	octane	25
2	toluene	38
3	THF	40
4	MeCN	40
5	1,4-dioxane	48
6	methanol	57
7	DMSO	87 (89 in 2nd use)
8	DMSO, no catalyst	0
9	DMSO, in N ₂	0

[a] Reaction conditions: 0.5 mmol 1-indanone, 1 mmol aniline, 10 mg Cu₂O, 2 mL solvent, 0.6 MPa O₂, 110 °C, 6 h. The yield was determined by GC analysis using n-dodecane as the internal standard. The main byproduct is *o*-phthalic anhydride. DMSO = dimethylsulfoxide, THF = tetrahydrofuran.

Table 2: Substrate scope of amines.^[a]

1	2	3
4	5	6
7	8	9
10	11	12
13	14	15
16	17	18
19	20	21
22	23	24
25	26	27
28	29	30
31	32	33
34	35	36
37	38	39
40	41	42
43	44	45
46	47	48
49	50	51
52	53	54
55	56	57
58	59	60
61	62	63
64	65	66
67	68	69
70	71	72
73	74	75
76	77	78
79	80	81
82	83	84
85	86	87
88	89	90
91	92	93
94	95	96
97	98	99
100	101	102
103	104	105
106	107	108
109	110	111
112	113	114
115	116	117
118	119	120
121	122	123
124	125	126
127	128	129
130	131	132
133	134	135
136	137	138
139	140	141
142	143	144
145	146	147
148	149	150
151	152	153
154	155	156
157	158	159
160	161	162
163	164	165
166	167	168
169	170	171
172	173	174
175	176	177
178	179	180
181	182	183
184	185	186
187	188	189
190	191	192
193	194	195
196	197	198
199	200	201
202	203	204
205	206	207
208	209	210
211	212	213
214	215	216
217	218	219
220	221	222
223	224	225
226	227	228
229	230	231
232	233	234
235	236	237
238	239	240
241	242	243
244	245	246
247	248	249
250	251	252
253	254	255
256	257	258
259	260	261
262	263	264
265	266	267
268	269	270
271	272	273
274	275	276
277	278	279
280	281	282
283	284	285
286	287	288
289	290	291
292	293	294
295	296	297
298	299	300
301	302	303
304	305	306
307	308	309
310	311	312
313	314	315
316	317	318
319	320	321
322	323	324
325	326	327
328	329	330
331	332	333
334	335	336
337	338	339
340	341	342
343	344	345
346	347	348
349	350	351
352	353	354
355	356	357
358	359	360
361	362	363
364	365	366
367	368	369
370	371	372
373	374	375
376	377	378
379	380	381
382	383	384
385	386	387
388	389	390
391	392	393
394	395	396
397	398	399
400	401	402
403	404	405
406	407	408
409	410	411
412	413	414
415	416	417
418	419	420
421	422	423
424	425	426
427	428	429
430	431	432
433	434	435
436	437	438
439	440	441
442	443	444
445	446	447
448	449	450
451	452	453
454	455	456
457	458	459
460	461	462
463	464	465
466	467	468
469	470	471
472	473	474
475	476	477
478	479	480
481	482	483
484	485	486
487	488	489
490	491	492
493	494	495
496	497	498
499	500	501
502	503	504
505	506	507
508	509	510
511	512	513
514	515	516
517	518	519
520	521	522
523	524	525
526	527	528
529	530	531
532	533	534
535	536	537
538	539	540
541	542	543
544	545	546
547	548	549
550	551	552
553	554	555
556	557	558
559	560	561
562	563	564
565	566	567
568	569	570
571	572	573
574	575	576
577	578	579
580	581	582
583	584	585
586	587	588
589	590	591
592	593	594
595	596	597
598	599	600
601	602	603
604	605	606
607	608	609
610	611	612
613	614	615
616	617	618
619	620	621
622	623	624
625	626	627
628	629	630
631	632	633
634	635	636
637	638	639
640	641	642
643	644	645
646	647	648
649	650	651
652	653	654
655	656	657
658	659	660
661	662	663
664	665	666
667	668	669
670	671	672
673	674	675
676	677	678
679	680	681
682	683	684
685	686	687
688	689	690
691	692	693
694	695	696
697	698	699
700	701	702
703	704	705
706	707	708
709	710	711
712	713	714
715	716	717
718	719	720
721	722	723
724	725	726
727	728	729
730	731	732
733	734	735
736	737	738
739	740	741
742	743	744
745	746	747
748	749	750
751	752	753
754	755	756
757	758	759
760	761	762
763	764	765
766	767	768
769	770	771
772	773	774
775	776	777
778	779	780
781	782	783
784	785	786
787	788	789
790	791	792
793	794	795
796	797	798
799	800	801
802	803	804
805	806	807
808	809	810
811	812	813
814	815	816
817	818	819
820	821	822
823	824	825
826	827	828
829	830	831
832	833	834
835	836	837
838	839	840
841	842	843
844	845	846
847	848	849
850	851	852
853	854	855
856	857	858
859	860	861
862	863	864
865	866	867
868	869	870
871	872	873
874	875	876
877	878	879
880	881	882
883	884	885
886	887	888
889	890	891
892	893	894
895	896	897
898	899	900
901	902	903
904	905	906
907	908	909
910	911	912
913	914	915
916	917	918
919	920	921
922	923	924
925	926	927
928	929	930
931	932	933
934	935	936
937	938	939
940	941	942
943	944	945
946	947	948
949	950	951
952	953	954
955	956	957
958	959	960
961	962	963
964	965	966
967	968	969
970	971	972
973	974	975
976	977	978
979	980	981
982	983	984
985	986	987
988	989	990
991	992	993
994	995	996
997	998	999
1000	1001	1002

[a] 0.5 mmol 1-indanone, 1 mmol amines, 10 mg Cu₂O, 0.6 MPa O₂, 2 mL DMSO, 110 °C, 8 h. The yield was determined by GC analysis using n-dodecane as the internal standard. The data within parentheses shows the yield of the isolated product.


range of amines were coupled with a ketone to form N-substituted cyclic imides with moderate to excellent yields. Electron-donating or electron-withdrawing substituents show

little effect on the reaction (entries 1–11). 1-Naphthylamine and *tert*-butylamine give moderate yield because of the steric hindrance for C–N bond formation (entries 12 and 19). Amines bearing halo, Me, OH, or MeO groups afford the corresponding imides in yields ranging from 76 to 92 % (entries 2–11, 15, 16, 24). The S- and N-containing heterocyclic amines, which usually poison noble-metal centers, can also be converted into imides with yields ranging from 77 to 89 % (entries 13 and 17).


The scope of the reaction with respect to ketone was then investigated (Table 3). Substituted 1-indanones, including halo, Me, and MeO substituents were all converted into the corresponding imides with yields in the 86–90 % range. Apparent substituent effects are observed. Different to the oxidation reaction, such as alcohol and amine oxidation,^[11] ketones with electron-withdrawing substituents are more easily oxidized. A higher reaction temperature and longer reaction time is needed for the conversion of methoxy-substituted 1-indanones.

Table 3: Substrate scope of ketones.^[a]

1	2	3
4	5	6
7	8	9
10	11	12
13	14	15
16	17	18
19	20	21
22	23	24
25	26	27
28	29	30
31	32	33
34	35	36
37	38	39
40	41	42
43	44	45
46	47	48
49	50	51
52	53	54
55	56	57
58	59	60
61	62	63
64	65	66
67	68	69
70	71	72
73	74	75
76	77	78
79	80	81
82	83	84
85	86	87
88	89	90
91	92	93
94	95	96
97	98	99
100	101	102
103	104	105
106	107	108
109	110	111
112	113	114
115	116	117
118	119	120
121	122	123
124	125	126
127	128	129
130	131	132
133	134	135
136	137	138</



trometry (MS; see Figure S2). *o*-Phthalic anhydride was detected (see Figure S3). We then found that the reaction of *o*-phthalic anhydride with aniline quickly generated *N*-phenylphthalimide [Eqs. (3) and (4)]. This result infers that *o*-phthalic anhydride is an intermediate en route to the imide. Now it becomes clear that there are two steps: the oxidation of 1-indanone to *o*-phthalic anhydride [Eq. (3)] and the imidization reaction of *o*-phthalic anhydride with amines [Eq. (4)]. In first step, 1-indanone is oxidized to *o*-phthalic anhydride with 93 % GC yield within 6 hours. The second step is much faster and 96 % GC yield of the imide is obtained after 2 hours. Step 2 could also occur without catalyst and oxygen. Cu₂O and oxygen show little effect on the step 2.



No reaction takes place if the ketone is lacking both two α -C-H bonds and two β -C-H bonds (Table 4, entries 2–4). By employing 1,2-indandione and 1,2,3-indantrione as reactants, phthalic anhydrides are obtained with 99 % conversion after 2 hours, which is much faster than that with 1-indanone (entries 5 and 6). Therefore, 1,2-indandione and 1,2,3-indantrione are two intermediates formed by the C-H bond oxidation of 1-indanone.

Correlating $-\ln(1-C)$ (C is the conversion of substituted 1-indanones) against reaction time (ks) indicates a linear relationship and a pseudo-first-order reaction with respect to the substituted 1-indanones. The apparent activation energy, E_a , is determined to be 78 kJ mol⁻¹ (Figure 1 a). The addition of aniline lowers activation energy ($E_a = 70$ kJ mol⁻¹) and increases the reaction rate (see Figure S4). This change may result from the fact that aniline either as basic molecule helps deprotonate the C-H bond, or as a ligand coordinates to the surface copper sites to increase reactivity.^[14] A linear relationship between $\log(k_X/k_H)$ and substituent constant σ was established for substituted 1-indanone ($X = p\text{-OCH}_3, p\text{-H}, p\text{-$

Table 4: Oxidation of the derivatives of 1-indanone.^[a]

1	2	3
conv. 93% (6 h)	no reaction	no reaction
4	5	6
no reaction	conv. 99% (2 h)	conv. 99% (2 h)

[a] 0.5 mmol substrate, 10 mg Cu₂O, 2 mL DMSO, 110 °C, 0.6 MPa O₂, 8 h. Conversions were determined by GC analysis using *n*-dodecane as the internal standard.




Figure 1. a) Arrhenius plot with or without the addition of aniline. b) Hammett plot of the oxidation of substituted 1-indanones.

$p\text{-Cl}, p\text{-Br}$ (Figure 1 b). The resulting Hammett parameter, ρ , is 0.56, thus suggesting the reaction is substituent-sensitive and carbanions are involved. The α -C-H and β -C-H bond are cleaved by a proton transfer on the catalyst surface. Electron-withdrawing substituents can decrease the negative charge on the carbanions and thus stabilize the intermediates. DFT calculations find that 1-indanone adsorbs on the Cu₂O-(111) surface in a horizontal manner (Figure 2), where three aryl carbon atoms bind to three adjacent copper atoms on the top sites. The oxygen atom in the carbonyl group binds to the surface on the bridge site. The adsorption is exothermic with -1.86 eV. Two possible pathways for the oxidation of 1-indanone are calculated (Figure 3; the detailed adsorption and transition state are provided in the Supporting Information). One is the oxidation of the α -C-H bond (Path A), and the other is the oxidation of the β -C-H bond (Path B). The overall the E_a value of the α -C-H oxidation is 2.91 eV, which is much lower than that of the β -C-H bond oxidation (3.56 eV). In the initial step for the C-H bond oxidation, the activation energy for deprotonating one α -H at the Cu₂O surface is only 0.44 eV, which is much lower than that of removing a β -H (1.67 eV). Furthermore, Path A is thermodynamically favored as evidenced by a more-negative reaction enthalpy (-0.75 eV vs 0.01 eV). These results indicate that the α -C-H bond oxidation is preferred and Path A is the





Figure 2. The adsorption of 1-indanone on $\text{Cu}_2\text{O}(111)$. a) Top view. b) Side view.

main reaction route. In addition, α -hydroxy-1-indanone is unstable and is further oxidized to 1,2-indandione with merely 0.14 eV activation energy.

Based on the above-mentioned results and the following experiments, we may come to a tentative reaction mechanism (Scheme 2). In the initial step, one of the α -C-H bonds in 1-indanone is oxidized by the activated oxygen to give α -hydroxy-1-indanone, which is further oxidized to 1,2-indandione by the oxidation of the other α -C-H bond.^[15] Then, the oxidation of the α -C-H bond adjacent to carbonyl group is repeated for 1,2-indandione with the formation of the 1,2,3-indantrione intermediate. This step shows why the β -C-H bond is indispensable. The 1,2,3-indantrione intermediate is further converted into *o*-phthalic anhydride, thus releasing CO_2 . We detected 1,2,3-indantrione by in situ IR. It first appears and then disappears as the reaction proceeds (see Figure S5). Finally, the fast reaction of *o*-phthalic anhydride with an amine forms the imide. The acceleration of the reaction by the amine may be due either to the base activation of the α -C-H bond or to adjusting the catalytic properties



Scheme 2. Possible reaction mechanism for the Cu_2O -catalyzed oxidation of ketones with amines to generate imides.

by coordination on the Cu_2O surface.^[14] It is under investigation.

In summary, we report a novel strategy for the Cu_2O -catalyzed oxidative C-C bond cleavage of a ketone for the C-N bond formation with molecular oxygen as the oxidant. A wide range of cyclic imides are synthesized from 1-indanone and amines with moderate to excellent yields. The presence of both the α -C-H and β -C-H bond is essential for C-C bond cleavage. Carbanion intermediates are involved, and the α -C-H is initially oxidized.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (21422308 21303183) and the Doctor Startup Foundation of Liaoning Province.




Figure 3. DFT calculation of the oxidation of the α -C-H and β -C-H bond of 1-indanone to 1,2-indandione and 1,3-indandione, respectively. Red O, blue Cu, grey C, white H.

Keywords: copper · density functional calculations · heterocycles · heterogeneous catalysis · oxidation

How to cite: *Angew. Chem. Int. Ed.* **2015**, *54*, 14061–14065
Angew. Chem. **2015**, *127*, 14267–14271

- [1] a) A. Rahimi, A. Azarpira, H. Kim, J. Ralph, S. S. Stahl, *J. Am. Chem. Soc.* **2013**, *135*, 6415–6418; b) C. J. Allpress, L. M. Berreau, *Coord. Chem. Rev.* **2013**, *257*, 3005–3029; c) M. Murakami, T. Tsuruta, Y. Ito, *Angew. Chem. Int. Ed.* **2000**, *39*, 2484–2486; *Angew. Chem.* **2000**, *112*, 2600–2602; d) H. Liu, M. H. Feng, X. F. Jiang, *Chem. Asian J.* **2014**, *9*, 3360–3389; e) F. Chen, T. Wang, N. Jiao, *Chem. Rev.* **2014**, *114*, 8613–8661; f) L. Soullart, N. Cramer, *Chem. Rev.* **2015**, *115*, 9410–9464.
- [2] a) L. Liu, N. Ishida, M. Murakami, *Angew. Chem. Int. Ed.* **2012**, *51*, 2485–2488; *Angew. Chem.* **2012**, *124*, 2535–2538; b) M. Sai, H. Yorimitsu, K. Oshima, *Angew. Chem. Int. Ed.* **2011**, *50*, 3294–3298; *Angew. Chem.* **2011**, *123*, 3352–3356; c) H. Li, Y. Li, X. S. Zhang, K. Chen, X. Wang, Z. J. Shi, *J. Am. Chem. Soc.* **2011**, *133*, 15244–15247; d) C. Qin, W. Zhou, F. Chen, Y. Ou, N. Jiao, *Angew. Chem. Int. Ed.* **2011**, *50*, 12595–12599; *Angew. Chem.* **2011**, *123*, 12803–12807; e) C. Qin, P. Feng, Y. Ou, T. Shen, T. Wang, N. Jiao, *Angew. Chem. Int. Ed.* **2013**, *52*, 7850–7854; *Angew. Chem.* **2013**, *125*, 8004–8008; f) A. Dermenci, J. W. Coe, G. Dong, *Org. Chem. Front.* **2014**, *1*, 567–581.
- [3] a) C. Öztürk, K. Topal, V. Aviyente, N. S. Tuzun, E. S. Fernandez, S. Arseniyadis, *J. Org. Chem.* **2005**, *70*, 7080–7086; b) W. S. Trahanovsky, J. R. Gilmore, P. C. Heaton, *J. Org. Chem.* **1973**, *38*, 760–763; c) W. S. Trahanovsky, L. H. Young, M. H. Bierman, *J. Org. Chem.* **1969**, *34*, 869–871; d) P. Spanning, P. C. A. Bruijninx, B. M. Weckhuysen, R. J. M. K. Gebbink, *RSC Adv.* **2013**, *3*, 6606–6613; e) Z. Elkhayat, I. Safir, M. Dakir, S. Arseniyadis, *Tetrahedron: Asymmetry* **2007**, *18*, 1589–1602; f) S. Rup, F. Zimmermann, E. Meux, M. Schneider, M. Sindt, N. Oget, *Ultrason. Sonochem.* **2009**, *16*, 266–272.
- [4] a) S. Paria, P. Halder, T. K. Paine, *Angew. Chem. Int. Ed.* **2012**, *51*, 6195–6199; *Angew. Chem.* **2012**, *124*, 6299–6303; b) H. Liu, C. Dong, Z. Zhang, P. Wu, X. Jiang, *Angew. Chem. Int. Ed.* **2012**, *51*, 12570–12574; *Angew. Chem.* **2012**, *124*, 12738–12742; c) B. Sedai, C. Díaz-Urrutia, R. T. Baker, R. Wu, L. A. P. Silks, S. K. Hanson, *ACS Catal.* **2011**, *1*, 794–804; d) C. J. Allpress, K. Grubel, E. Szajna-Fuller, A. M. Arif, L. M. Berreau, *J. Am. Chem. Soc.* **2013**, *135*, 659–668; e) X. Q. Huang, X. Y. Li, M. C. Zou, S. Song, C. H. Tang, Y. Z. Yuan, N. Jiao, *J. Am. Chem. Soc.* **2014**, *136*, 14858–14865; f) C. H. Tang, N. Jiao, *Angew. Chem. Int. Ed.* **2014**, *53*, 6528–6532; *Angew. Chem.* **2014**, *126*, 6646–6650; g) C. Zhang, P. Feng, N. Jiao, *J. Am. Chem. Soc.* **2013**, *135*, 15257–15262; h) J. Wang, W. Chen, S. Zuo, L. Liu, X. Zhang, J. Wang, *Angew. Chem. Int. Ed.* **2012**, *51*, 12334–12338; *Angew. Chem.* **2012**, *124*, 12500–12504; i) B. Tiwari, J. Zhang, Y. R. Chi, *Angew. Chem. Int. Ed.* **2012**, *51*, 1911–1914; *Angew. Chem.* **2012**, *124*, 1947–1950; j) W. Zhou, W. Y. Fan, Q. J. Jiang, Y. F. Lang, N. Jiao, *Org. Lett.* **2015**, *17*, 2542–2545.
- [5] a) Y. J. Bian, C. Y. Chen, Z. Z. Huang, *Chem. Eur. J.* **2013**, *19*, 1129–1133; b) J. A. Letizia, M. R. Salata, C. M. Tribout, A. Facchetti, M. A. Ratner, T. J. Marks, *J. Am. Chem. Soc.* **2008**, *130*, 9679–9694; c) Q. T. Zhang, J. M. Tour, *J. Am. Chem. Soc.* **1997**, *119*, 5065–5066.
- [6] a) M. A. Ali, S. M. A. H. Siddiki, K. Kon, J. Hasegawa, K. Shimizu, *Chem. Eur. J.* **2014**, *20*, 14256–14260; b) M. Nasr-Esfahani, M. Montazerzohori, N. Filvan, *J. Serb. Chem. Soc.* **2012**, *77*, 415–421.
- [7] a) K. M. Driller, H. Klein, R. Jackstell, M. Beller, *Angew. Chem. Int. Ed.* **2009**, *48*, 6041–6044; *Angew. Chem.* **2009**, *121*, 6157–6160; b) X. Y. Yan, K. Fang, H. L. Liu, C. J. Xi, *Chem. Commun.* **2013**, *49*, 10650–10652.
- [8] a) Z. Zhang, F. Wang, M. Wang, S. Xu, H. Chen, C. Zhang, J. Xu, *Green Chem.* **2014**, *16*, 2523–2527; b) A. Grirrane, A. Corma, H. Garcia, *Science* **2008**, *322*, 1661–1664.
- [9] M. Wang, X. Gu, H. Su, J. Lu, J. Ma, M. Yu, Z. Zhang, F. Wang, *J. Catal.* **2015**, *330*, 458–464.
- [10] P. Lignier, R. Bellabarba, R. P. Tooze, *Chem. Soc. Rev.* **2012**, *41*, 1708–1720.
- [11] M. Wang, F. Wang, J. P. Ma, M. R. Li, Z. Zhang, Y. H. Wang, X. C. Zhang, J. Xu, *Chem. Commun.* **2014**, *50*, 292–294.
- [12] a) I. Efremenko, R. Neumann, *J. Am. Chem. Soc.* **2012**, *134*, 20669–20680; b) Q. Hua, T. Cao, X. K. Gu, J. Q. Lu, Z. Q. Jiang, X. R. Pan, L. F. Luo, W. X. Li, W. X. Huang, *Angew. Chem. Int. Ed.* **2014**, *53*, 4856–4861; *Angew. Chem.* **2014**, *126*, 4956–4961.
- [13] a) P. Pirovano, A. M. Magherusan, C. McGlynn, A. Ure, A. Lynes, A. R. McDonald, *Angew. Chem. Int. Ed.* **2014**, *53*, 5946–5950; *Angew. Chem.* **2014**, *126*, 6056–6060; b) S. Kim, J. W. Ginsbach, A. I. Billah, M. A. Siegler, C. D. Moore, E. I. Solomon, K. D. Karlin, *J. Am. Chem. Soc.* **2014**, *136*, 8063–8071; c) M. Che, A. J. Tench, *Adv. Catal.* **1983**, *32*, 1–148.
- [14] a) Y. Z. Huang, J. Gao, H. Ma, H. Miao, J. Xu, *Tetrahedron Lett.* **2008**, *49*, 948–951; b) M. S. DeClue, J. S. Siegel, *Org. Biomol. Chem.* **2004**, *2*, 2287–2298; c) D. J. Berrisford, C. Bolm, K. B. Sharpless, *Angew. Chem. Int. Ed. Engl.* **1995**, *34*, 1059–1070; *Angew. Chem.* **1995**, *107*, 1159–1171.
- [15] L. Zhang, X. Bi, X. Guan, X. Li, Q. Liu, B. D. Barry, P. Liao, *Angew. Chem. Int. Ed.* **2013**, *52*, 11303–11307; *Angew. Chem.* **2013**, *125*, 11513–11517.

Received: August 28, 2015

Published online: October 23, 2015