

Direct Trifluoromethylthiolation and Perfluoroalkylthiolation of C(sp²)–H Bonds with CF₃SO₂Na and R_fSO₂Na

Lvqi Jiang, Jinlong Qian, Wenbin Yi,* Guoping Lu, Chun Cai, and Wei Zhang*

Abstract: A new method for CF₃SO₂Na-based direct trifluoromethylthiolation of C(sp²)–H bonds has been developed. CF₃SSCF₃ is generated in situ from cheap and easy-to-handle CF₃SO₂Na, and in the presence of CuCl can be used for electrophilic trifluoromethylthiolation of indoles, pyrroles, and enamines. The method has been extended to perfluoroalkylthiolation reactions using R_fSO₂Na.

Organofluorination is an area of active research because incorporation of either fluorine or fluorine-containing groups can alter the physical, chemical, and biological properties of the parent molecules.^[1] In the organofluorine family, the trifluoromethylthiol (CF₃S) group holds a special position because of its high electron-withdrawing capability, good lipophilicity,^[2] and bioavailability.^[3] It has a great potential in the development of new pharmaceutical and agrochemical chemicals.^[4]

Direct trifluoromethylthiolation using CF₃SH, CF₃SCl, and CF₃SSCF₃ has been reported in the literature.^[5] Since they are highly reactive and hard-to-handle gaseous chemicals, a series of CF₃S-based reagents, including quaternary ammonium Me₄NSCF₃,^[6] metallic CuSCF₃ and AgSCF₃,^[7] and the shelf-stable chemicals **1a–f**, have been developed for trifluoromethylthiolation (Figure 1).^[8–13] However, they have

to be prepared by trifluoromethylthiolation or trifluoromethylation reactions, and could be very expensive. In 2009, the group of Magnier reported a CF₃SO₂K-initiated trifluoromethylation reaction. A small amount of trifluoromethylthiolation product was detected in the reaction mixture.^[14] The group of Shibata recently used the trifluoromethanesulfonyl hypervalent iodonium ylide **2a** as a CF₃SO₂-based trifluoromethylthiolation agent.^[15] It was proposed that CF₃SSCF₃ is generated in the reaction system.^[15c] Sodium trifluoromethanesulfonate (CF₃SO₂Na; **2b**), known as the Langlois reagent, is a readily available and is a stable benchtop solid. It is a good source for the generation of CF₃ radical.^[16] We envisioned that it could be a CF₃SO₂-based agent for direct electrophilic trifluoromethylthiolation through the in situ generation of CF₃SSCF₃.^[5m]

Development of a system for efficient conversion of CF₃SO₂Na into CF₃SSCF₃ was the key for this project. We first examined the CF₃SO₂Na reaction of indole^[17] in the presence of hypophosphite,^[18] phosphite ester,^[19] and iodo-trimethylsilane^[20] since they are known reducing agents for sulfonate, sulfonate, and sulfoxide (Table 1, entries 1–3). A reaction of 2 equivalents of (EtO)₂P(O)H and CF₃SO₂Na at 110 °C without using any oxidant gave the trifluoromethyl-




Figure 1. User-friendly trifluoromethylthiolation agents. Ts = 4-toluene-sulfonyl.

[*] L. Jiang, J. Qian, Prof. Dr. W. Yi, G. Lu, C. Cai
School of Chemical Engineering
Nanjing University of Science and Technology
200 Xiao Ling Wei Street, Nanjing 210094 (China)
E-mail: yiw@njust.edu.cn

Prof. Dr. W. Zhang
Department of Chemistry, University of Massachusetts Boston
100 Morrissey Boulevard, Boston, MA 02125 (USA)
E-mail: wei2.zhang@umb.edu

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201508495>.

Table 1: Optimization of trifluoromethylthiolation with NaSO₂CF₃.

Entry	Reductant	Oxidant	Cat. (equiv)	Yield [%] ^[a]
1	EtOP(O)H ₂	–	–	–
2	(EtO) ₂ P(O)H	–	–	32
3	Me ₃ SiI	–	–	–
4	(EtO) ₂ P(O)H	DMSO	–	trace
5	(EtO) ₂ P(O)H	TMSO	–	–
6	(EtO) ₂ P(O)H	I ₂	–	–
7	(EtO) ₂ P(O)H	–	CuCl (0.5)	31
8	(EtO) ₂ P(O)H	DMSO	CuI (0.5)	13
9	(EtO) ₂ P(O)H	DMSO	CuOAc (0.5)	20
10	(EtO) ₂ P(O)H	DMSO	CuCl ₂ (0.5)	14
11	(EtO) ₂ P(O)H	DMSO	AgNO ₃ (0.5)	–
12	(EtO) ₂ P(O)H	DMSO	I ₂ (0.2)	trace
13	(EtO) ₂ P(O)H	DMSO	CuCl (0.5)	55
14	(EtO) ₂ P(O)H	DMSO	CuCl (0.8)	77
15	(EtO) ₂ P(O)H	DMSO	CuCl (1.0)	94
16 ^[b]	(EtO) ₂ P(O)H	DMSO	CuCl (1.0)	52
17 ^[c]	(EtO) ₂ P(O)H	DMSO	CuCl (1.0)	74

Reaction conditions: **3a** (0.2 mmol), PhMe (1 mL), 110 °C, under N₂.
[a] Determined by GC analysis. [b] **2b** (1 equiv). [c] **2b** (1.5 equiv).
DMSO = dimethylsulfoxide, TMSO = tetramethylene sulfoxide.

thiolation product **4a** in 32% yield (entry 2). Electrophilic trifluoromethylthiolation of CF_3SSCF_3 generates 1 equivalent of CF_3S^- , which could be oxidized back to CF_3SSCF_3 with agents such as DMSO.^[21]

We then performed reactions by adding either DMSO, TMSO, or I_2 to the reaction system (Table 1, entries 4–6). The reaction with DMSO afforded a trace amount of **4a**. Metal salts are able to catalyze trifluoromethylthiolation with **2b**,^[15] while I_2 is able to promote sulfonylation of sodium sulfonates.^[19] After testing copper and silver salts, as well as I_2 for the $\text{CF}_3\text{SO}_2\text{Na}$ reactions (entries 7–13), we found that the reaction with CuCl (0.5 equiv) together with DMSO afforded a good amount of **4a** (55%; entry 13). Optimization of the reaction conditions (entries 10–17) revealed that using $(\text{EtO})_2\text{P}(\text{O})\text{H}$ (2.0 equiv), DMSO (3.0 equiv), and CuCl (1.0 equiv) in toluene at 110 °C for 12 hours could increase the product yield to 94% (entry 14).



Reactions of a series of substituted indoles under the optimized reaction conditions were conducted. Trifluoromethylthiolation took place selectively at the 3-position of indoles to form products bearing methyl (**4b**), methoxy (**4c**), halogens (**4d–f**), nitro (**4g**), esters (**4h, 4o**), and hydroxy (**4n**) groups at the 4-, 5-, 6-, and 7-positions in 42–93% yields (Table 2). It was found that indoles with an electron-donating group gave better results than those with an electron-withdrawing group, for example, **4n** and **4o**. The bulky substituent in **4o** and **4r** affected the yields. N-substituted indoles with either Me, Et, or Ph groups afforded the products **4s–v** in 77, 71, and 63% yield, respectively. The structure of **4h** was confirmed by single-crystal X-ray analysis (see the Supporting Information).

Similar to indoles, pyrroles are important nitrogen-containing heterocyclic rings existing in many natural products, biologically active molecules, and dyes for solar cells.^[22] We were able to do direct trifluoromethylthiolation reaction of pyrroles (Scheme 1). Under the same reaction conditions for indoles, the reaction of pyrroles afforded **5a–c** in good yields. Enamines, an important building block for a variety of biologically and synthetically interested nitrogen-containing heterocycles,^[23] were also used for the synthesis of **6a–b**.


It was found in the study of glycosidase inhibitors that the introduction of CF_3S and R_fS with a fluorine chain could stabilize the parent molecules and change their amphiphilic properties.^[24] The R_fS groups were introduced by the reaction of a thiophenol or sulfide with perfluoroalkyl iodides.^[25] We have successfully extended the direct trifluoromethylthiolation for perfluoroalkylthiolation. Sodium perfluoroalkanesulfonates ($\text{R}_f\text{SO}_2\text{Na}$), with a different R_f group, were prepared following the procedure reported by Hu and DesMarteau.^[26] These compounds were used for the perfluoroalkylthiolation reactions with indole. The products **7a–d**, bearing $\text{C}_2\text{F}_5\text{S}$, $\text{C}_4\text{F}_9\text{S}$, $\text{C}_6\text{F}_{13}\text{S}$, and $\text{C}_8\text{F}_{17}\text{S}$ groups, respectively, were obtained in good yields (Scheme 2).

To understand the mechanism of $\text{CF}_3\text{SO}_2\text{Na}$ -based direct trifluoromethylthiolation, a reaction of indole under the optimized reaction conditions was closely monitored by ^{19}F NMR spectroscopy using PhCF_3 as an internal standard. During the 12 hour reaction process, five fluorine peaks, including those for the reagent **2b** ($\delta = -84.46$ ppm), three

Table 2: Copper-catalyzed trifluoromethylthiolation of indoles.


 3	$\text{CF}_3\text{SO}_2\text{Na}$ (2 equiv) 2b	$(\text{EtO})_2\text{P}(\text{O})\text{H}$ DMSO CuCl, toluene 110 °C, 12 h	 4
 4a , 89%	 4b , 93%	 4c , 94%	 4d , 72%
 4e , 70%	 4f , 67%	 4g , 71%	 4h , 80%
 4i , 81%	 4j , 78%	 4k , 89%	 4l , 80%
 4m , 82%	 4n , 42%	 4o , 51%	 4p , 94%
 4q , 78%	 4r , 54%	 4s , 77%	 4t , 71%
 4u , 63%	 4v , 80%		

Reaction conditions: **3** (0.2 mmol), **2b** (0.4 mmol), $(\text{EtO})_2\text{P}(\text{O})\text{H}$ (0.4 mmol), DMSO (0.6 mmol), CuCl (0.2 mmol) in PhMe (1 mL) under N_2 at 110 °C for 12 h. Yields are those for the isolated products.



Scheme 1. Trifluoromethylthiolation of pyrroles and enamines. Reaction conditions: pyrroles or enamines (0.2 mmol), **2b** (0.4 mmol), $(\text{EtO})_2\text{P}(\text{O})\text{H}$ (0.4 mmol), DMSO (0.6 mmol), CuCl (0.2 mmol) in PhMe (1 mL) under N_2 at 110 °C for 12 h. [a] Yield of isolated product. [b] Yield determined by GC-MS.

intermediates (**8** at $\delta = -78.58$ ppm, **9** at $\delta = -62.14$ ppm, and **10** at $\delta = -45.48$ ppm), and product **4a** ($\delta = -44.50$ ppm) were observed (Figure 2). The $\text{CF}_3\text{SO}_2\text{Na}$ peak disappeared after 2 hours. The amount of **4a** increased steadily during the reaction process. The structure of **10** was confirmed to be CF_3SSCF_3 by its ^{19}F NMR data.^[15b,c] To gain more information for the intermediates, we performed four control reactions in



Scheme 2. Perfluoroalkylthiolation of indole. Reaction conditions: **3a** (0.2 mmol), $R_f\text{SO}_2\text{Na}$ (0.4 mmol), $(\text{EtO})_2\text{P}(\text{O})\text{H}$ (0.4 mmol), DMSO (0.6 mmol), CuCl (0.2 mmol) in PhMe (1 mL) under N_2 at 110°C for 12 h. Yields are those of the isolated products.




Figure 2. Progress of the reaction of **3a**, for up to 12 h, by ^{19}F NMR spectroscopy. The peaks (in CDCl_3) represent $\text{CF}_3\text{SO}_2\text{Na}$ (**2b**, $\delta = -84.46$ ppm), $\text{CF}_3\text{S}(\text{O})\text{H}$ (**8**, $\delta = -78.58$ ppm), the internal standard (PhCF_3 , $\delta = -62.43$ ppm), $\text{CF}_3\text{S}-\text{OH}$ (**9**, $\delta = -62.14$ ppm), CF_3SSCF_3 (**10**, $\delta = -45.48$ ppm), and **4a** ($\delta = -44.50$ ppm).

the absence of indole. From the reaction of $\text{CF}_3\text{SO}_2\text{Na}$ and $(\text{EtO})_2\text{P}(\text{O})\text{H}$, a small peak for $\text{CF}_3\text{SO}_2\text{Na}$ and a big peak for **8** were detected (Figure 3, I). Similar peaks were detected for the reaction with either CuCl (Figure 3, II) or DMSO (Figure 3, III). When both CuCl and DMSO were added, all fluorine peaks, including those of the intermediates **8**, **9**, **10**, were detected (Figure 3, IV). We inferred that **8** is $\text{CF}_3\text{S}(\text{O})\text{H}$, a reduction product of **2b**, and that **9** is CF_3SOH ,^[27] which was





Figure 3. Progress of reactions without indole, at 110°C , using ^{19}F NMR (in CDCl_3) spectroscopy.

generated through intramolecular nucleophilic collapse of $\text{CF}_3\text{S}(\text{O})\text{H}$.^[15c] The compound **9** is less stable than **8** and **10** and difficult for ^{19}F NMR detection. Analysis of a reaction mixture of **2b** and $(\text{EtO})_2\text{P}(\text{O})\text{H}$ showed a strong MS peak at m/z 119 ($M+1$), which matches the molecular weight of **8** and **9**, both of which have a molecular weight of 118 (see the Supporting Information). We also monitored the reaction of **3a** with **2b** and $(\text{EtO})_2\text{P}(\text{O})\text{H}$ (Figure 3, V), and were able to detect **4a** after 3 hours, along with a small amount of CF_3SSCF_3 . This result suggests path A (see Scheme 3) is possible, but not as efficient as path B. Only ^{19}F NMR data for RSO_nCF_3 compounds were found in the literature,^[11,28] and no information for $\text{CF}_3\text{S}(\text{O})\text{H}$ and CF_3SOH is available. The ^{19}F NMR data of related compounds are listed in Table 3 for comparison.

Table 3. ^{19}F NMR data for $\text{CF}_3\text{SO}_n\text{R}$ and $\text{CF}_3\text{SO}_n\text{Na/H}$.

$\text{CF}_3\text{SO}_n\text{R}$ (literature) ^{19}F NMR (δ in ppm)	$\text{CF}_3\text{SO}_n\text{Na/H}$ (this work) ^{19}F NMR (δ in ppm, in CDCl_3)
$\text{CF}_3\text{SO}_2\text{CH}_3$: $-79.95^{[28a]}$	$\text{CF}_3\text{SO}_2\text{Na}$ (2b): -84.46
$\text{CF}_3\text{S}(\text{O})\text{CH}_3$: $-76.6^{[28b]}$	$\text{CF}_3\text{S}(\text{O})\text{H}$ (8): -78.58
CF_3SOR (1d): $-51.91^{[11]}$	CF_3SOH (9): -62.15

On the bases of the experimental results shown in Table 1 as well as Figures 2 and 3, we proposed a possible mechanism for trifluoromethylthiolation with $\text{CF}_3\text{SO}_2\text{Na}$. By using 2 equivalents each of $\text{CF}_3\text{SO}_2\text{Na}$ and $(\text{EtO})_2\text{P}(\text{O})\text{H}$, 3 equivalents of DMSO, and 1 equivalent of CuCl (Scheme 3), the



Scheme 3. Proposed mechanism for $\text{CF}_3\text{SO}_2\text{Na}$ trifluoromethylthiolation.

reduction of $\text{CF}_3\text{SO}_2\text{Na}$ with $(\text{EtO})_2\text{P}(\text{O})\text{H}$ leads to the formation of $\text{CF}_3\text{S}(\text{O})\text{H}$, which is then converted into CF_3SOH though intramolecular nucleophilic collapse.^[15c] CF_3SOH is unstable and able to produce CF_3S^+ ^[27] for the formation of a small amount of **4a** (Table 1, entry 2) through path A (Scheme 3). Path B is more efficient for trifluoromethylthiolation since CuCl reduces CF_3SOH to CF_3SSCF_3 (Figure 3, IV). In this reaction DMSO assists the CuCl reduction of CF_3SOH and also serves as an oxidizing agent for the conversion of CF_3S^- back into CF_3SSCF_3 .^[21] CuCl serves as both a reducing agent for $\text{CF}_3\text{SO}_2\text{Na}$ and also a catalyst for CF_3SSCF_3 .

In summary, we have developed a new method using the cheap and stable sulfinate $\text{CF}_3\text{SO}_2\text{Na}$ for direct trifluoromethylthiolation of $\text{C}(\text{sp}^2)\text{--H}$ bonds. The reaction system consists of a reducing agent, $(\text{EtO})_2\text{P}(\text{O})\text{H}$ and CuCl , and an oxidizing agent, DMSO. CuCl is as a catalyst for the electrophilic trifluoromethylthiolation of CF_3SSCF_3 with indoles, pyrroles, and enamines. The method has been successfully extended for perfluoroalkylthiolation using $\text{R}_f\text{SO}_2\text{Na}$. This economically favorable and easy-to-handle reaction could be suitable for large-scale trifluoromethylthiolation reaction.

Acknowledgements

We gratefully acknowledge the National Natural Science Foundation of China (21476116, 21402093), Natural Science Foundation of Jiangsu (BK20141394), FRFCU (30920140122003), Priority Academic Program Development of Jiangsu Higher Education Institutions, the Centre for Green Chemistry at the University of Massachusetts Boston, and the Center for Advanced Materials and Technology in Nanjing University of Science and Technology for financial support.

Keywords: copper · fluorine · heterocycles · sulfur · synthetic methods

How to cite: *Angew. Chem. Int. Ed.* **2015**, *54*, 14965–14969
Angew. Chem. **2015**, *127*, 15178–15182

- [1] a) P. Kirsch, *Modern Fluoroorganic Chemistry: Synthesis Reactivity, Applications*, Wiley-VCH, Weinheim, **2004**; b) T. Hiyama, *Organofluorine Compounds: Chemistry and Properties*, Springer, Berlin, **2000**; c) K. Uneyama, *Organofluorine Chemistry*, Blackwell, Oxford, **2006**; d) *Fluorine in Medicinal Chemistry and Chemical Biology* (Ed.: I. Ojima), Wiley-Blackwell, Chichester, **2009**.
- [2] C. Hansch, A. Leo, R. W. Taft, *Chem. Rev.* **1991**, *91*, 165–195.
- [3] Selected reviews of trifluoromethylthiolation: a) V. N. Boiko, *Beilstein J. Org. Chem.* **2010**, *6*, 880–921; b) A. Tlili, T. Billard, *Angew. Chem. Int. Ed.* **2013**, *52*, 6818–6819; *Angew. Chem.* **2013**, *125*, 6952–6954; c) X. H. Xu, K. Matsuzaki, N. Shibata, *Chem. Rev.* **2015**, *115*, 731–764; d) X. Shao, C. Xu, L. Lu, Q. Shen, *Acc. Chem. Res.* **2015**, *48*, 1227–1236; e) K. Zhang, X.-H. Xu, F.-L. Qing, *Chin. J. Org. Chem.* **2015**, *35*, 556–569; f) J.-H. Lin, Y.-L. Ji, J.-C. Xiao, *Curr. Org. Chem.* **2015**, *19*, 1541–1553.
- [4] a) P. Laczay, G. Voros, G. Semjen, *Int. J. Parasitol.* **1995**, *25*, 753–756; b) P. Pommier, A. Keita, S. W. Robert, B. Dellac, H. C. Mundt, *Rev. Med. Vet.* **2003**, *154*, 41–46; c) J. N. Andre, L. G. Dring, G. Gillet, C. Mas-Chamberlin, *Br. J. Pharmacol.* **1979**, *66*, 506P; d) T. Silverstone, J. Fincham, J. Plumley, *Br. J. Clin. Pharmacol.* **1979**, *7*, 353–356; e) G. W. Counts, D. Gregory, D. Zeleznik, M. Turck, *Antimicrob. Agents Chemother.* **1977**, *11*, 708–711; f) N. Aswapokee, H. C. Neu, *Antimicrob. Agents Chemother.* **1979**, *15*, 444–446; g) J. Wang, M. S. Rosello, J. Acena, C. Pozo, A. E. Sorochinsky, S. Fustero, V. A. Soloshonok, H. Liu, *Chem. Rev.* **2014**, *114*, 2432–2506.
- [5] Selected examples for reaction of CF_3SH , F_3CSCl , and F_3CSSCF_3 : a) J. F. Harris, Jr., F. W. Stacey, *J. Am. Chem. Soc.* **1961**, *83*, 840–845; b) W. A. Sheppard, *J. Org. Chem.* **1964**, *29*, 895–898; c) J. F. Harris, Jr., *J. Org. Chem.* **1966**, *31*, 931–935; d) S. Andreades, J. F. Harris, W. A. Sheppard, *J. Org. Chem.* **1964**, *29*, 898–900; e) H. Bayreuther, A. Haas, *Chem. Ber.* **1973**, *106*, 1418–1422; f) M. Bauer, A. Haas, H. Muth, *J. Fluorine Chem.* **1980**, *16*, 129–136; g) A. Haas, W. Kortmann, Z. Anorg. Allg. Chem. **1983**, *501*, 79–88; h) A. Haas, M. Lieb, Y. Zhang, *J. Fluorine Chem.* **1985**, *30*, 203–210; i) A. Kolasa, *J. Fluorine Chem.* **1987**, *36*, 29–40; j) D. I. Rossman, A. J. Muller, E. O. Lewis, *J. Fluorine Chem.* **1991**, *55*, 221–224; k) R. Boese, A. Haas, M. Lieb, U. Roeske, *Chem. Ber.* **1994**, *127*, 449–455; l) S. Munavalli, D. K. Rohrbaugh, L. L. Szafraniec, H. D. Durst, *Phosphorus Sulfur Silicon Relat. Elem.* **2006**, *181*, 305–324; m) L. D. Tran, I. Popov, O. Daugulis, *J. Am. Chem. Soc.* **2012**, *134*, 18237–18240.
- [6] Selected examples for the reaction of Me_4NSCF_3 : a) V. V. Orda, L. M. Yagupolskii, V. F. Bystrov, A. U. Stepanyants, *J. Gen. Chem.* **1965**, *35*, 1628–1636; b) J. H. Clark, C. W. Jones, A. P. Kybett, M. A. McClinton, *J. Fluorine Chem.* **1990**, *48*, 249–253; c) S. T. Tavener, D. J. Adams, J. H. Clark, *J. Fluorine Chem.* **1999**, *95*, 171–176; d) D. J. Adams, J. H. Clark, *J. Org. Chem.* **2000**, *65*, 1456–1460; e) W. Tyrre, D. Naumann, B. Hoge, Y. L. Yagupolskii, *J. Fluorine Chem.* **2003**, *119*, 101–108.
- [7] Selective examples for the reaction of AgSCF_3 and CuSCF_3 : a) D. J. Adams, A. Goddard, J. H. Clark, D. J. Macquarrie, *Chem. Commun.* **2000**, 987–988; b) Q. Lefebvre, E. Fava, P. Nikolaenko, M. Rueping, *Chem. Commun.* **2014**, *50*, 6617–6619; c) P. Nikolaenko, R. Pluta, M. Rueping, *Chem. Eur. J.* **2014**, *20*, 9867–9870; d) Y. Huang, X. He, H. Li, Z. Weng, *Eur. J. Org. Chem.* **2014**, 7324–7328; e) X. Wang, Y. Zhou, G. Ji, G. Wu, M. Li, Y. Zhang, J. Wang, *Eur. J. Org. Chem.* **2014**, 3093–3096; f) S. Q. Zhu, X.-H. Xu, F. L. Qing, *Eur. J. Org. Chem.* **2014**, 4453–4456; g) W. Yin, Z. Wang, Y. Huang, *Adv. Synth. Catal.* **2014**, *356*, 2998–3006; h) K. Zhang, J. B. Liu, F. L. Qing, *Chem. Commun.* **2014**, *50*, 14157–14160.
- [8] F. Baert, J. Colomb, T. Billard, *Angew. Chem. Int. Ed.* **2012**, *51*, 10382–10385; *Angew. Chem.* **2012**, *124*, 10528–10531.
- [9] A. Ferry, T. Billard, B. R. Langlois, E. Bacqu, *Angew. Chem. Int. Ed.* **2009**, *48*, 8551–8555; *Angew. Chem.* **2009**, *121*, 8703–8707.
- [10] S. Alazet, L. Zimmer, T. Billard, *Chem. Eur. J.* **2014**, *20*, 93–97.
- [11] E. V. Vinogradova, P. Muller, S. L. Buchwald, *Angew. Chem. Int. Ed.* **2014**, *53*, 3125–3128; *Angew. Chem.* **2014**, *126*, 3189–3192.
- [12] T. Bootwicha, X. Liu, R. Pluta, I. Atodiresei, M. Rueping, *Angew. Chem. Int. Ed.* **2013**, *52*, 12856–12859; *Angew. Chem.* **2013**, *125*, 13093–13097.
- [13] C. Xu, B. Ma, Q. Shen, *Angew. Chem. Int. Ed.* **2014**, *53*, 9316–9320; *Angew. Chem.* **2014**, *126*, 9470–9474.
- [14] Y. Mace, B. Raymondeau, C. Pradet, J. C. Blazejewski, E. Magnier, *Eur. J. Org. Chem.* **2009**, 1390–1397.
- [15] a) Y. D. Yang, A. Azuma, E. Tokunaga, M. Yamasaki, M. Shiro, N. Shibata, *J. Am. Chem. Soc.* **2013**, *135*, 8782–8785; b) F. Yin, X.-S. Wang, *Org. Lett.* **2014**, *16*, 1128–1131; c) Z. Huang, Y.-D. Yang, E. Tokunaga, N. Shibata, *Org. Lett.* **2015**, *17*, 1094–1097; d) S. Arimori, M. Takada, N. Shibata, *Org. Lett.* **2015**, *17*, 1063–1065.
- [16] Selective examples for using $\text{CF}_3\text{SO}_2\text{Na}$ as a CF_3 radical source: a) B. R. Langlois, E. Laurent, N. Roidot, *Tetrahedron Lett.* **1991**, *32*, 7525–7528; b) B. R. Langlois, E. Laurent, N. Roidot, *Tetrahedron Lett.* **1992**, *33*, 1291–1294; c) B. R. Langlois, D. Montkgre, N. Roidot, *J. Fluorine Chem.* **1994**, *68*, 63–66; d) T. Billard, N. Roques, B. R. Langlois, *J. Org. Chem.* **1999**, *64*, 3813–3820; e) Y. D. Yang, K. Iwamoto, E. Tokunaga, N. Shibata, *Chem. Commun.* **2013**, *49*, 5510–5512; f) C. Zhang, *Adv. Synth. Catal.* **2014**, *356*, 2895–2906.
- [17] Selected recent examples of trifluoromethylthiolation of indoles and other aromatic compounds: a) B. Ma, X. Shao, Q. Shen, *J. Fluorine Chem.* **2015**, *171*, 73–77; b) S. Alazet, L. Zimmer, T. Billard, *J. Fluorine Chem.* **2015**, *171*, 78–81; c) K. Zhang, X.-H. Xu, F.-L. Qing, *J. Org. Chem.* **2015**, *80*, 7658–7665; d) Q. Wang, Z. Qi, F. Xie, X. Li, *Adv. Synth. Catal.* **2015**, *357*, 355–360;

- e) X. X. Shao, C.-F. Xu, L. Lu, Q. Shen, *J. Org. Chem.* **2015**, *80*, 3012–3021; f) R. Honeker, J. B. Ernst, F. Glorius, *Chem. Eur. J.* **2015**, *21*, 8047–8051; g) Q. Wang, F. Xie, X. Li, *J. Org. Chem.* **2015**, *80*, 8361–8366; h) M. Jereb, K. Gosak, *Org. Biomol. Chem.* **2015**, *13*, 3103–3115; i) S. Alazet, T. Billard, *Synlett* **2015**, 26, 76–78.
- [18] H. W. Pinnick, M. A. Reynolds, R. T. McDonald, Jr., W. D. Brewster, *J. Org. Chem.* **1980**, *45*, 930–932.
- [19] F. Xiao, H. Xie, S. Liu, G. J. Deng, *Adv. Synth. Catal.* **2014**, 356, 364–368.
- [20] G. A. Olah, S. C. Narang, L. D. Field, G. F. Salem, *J. Org. Chem.* **1980**, *45*, 4792–4793.
- [21] T. J. Wallacen, *J. Am. Chem. Soc.* **1964**, *86*, 2018–2021.
- [22] a) M. d'Ischia, A. Napolitano, A. Pezzella, In *Comprehensive Heterocyclic Chemistry III: Pyrroles and Their Benzo Derivatives: Applications*, Vol. 3 (Eds.: A. R. Katritzky, C. A. Ramsden, E. F. V. Scriven, R. J. K. Taylor), Elsevier, Amsterdam, **2008**, pp. 353–388; b) I. S. Young, P. D. Thornton, A. Thompson, *Nat. Prod. Rep.* **2010**, *27*, 1801–1839.
- [23] X. Ji, H. Huang, W. Xiong, K. Huang, W. Wu, H. Jiang, *J. Org. Chem.* **2014**, *79*, 7005–7011.
- [24] J. P. Begue, D. B. Delpon, *Bioorganic and Medicinal Chemistry of Fluorine*, Wiley, Hoboken, NJ, **2008**, chap. 6, pp. 180–222.
- [25] C. Pooput, W. R. Dolbier, Jr., M. Medebielle, *J. Org. Chem.* **2006**, *71*, 3564–3568.
- [26] L. Q. Hu, D. D. DesMarteau, *Inorg. Chem.* **1993**, *32*, 5007–5010.
- [27] R. N. Haszeldine, J. M. Kidd, *J. Chem. Soc.* **1955**, 2901–2910.
- [28] a) D. T. Saure, J. M. Shreeve, *Inorg. Chem.* **1971**, *10*, 358–362; b) N. Walker, A. J. Leffler, *Inorg. Chem.* **1974**, *13*, 484–486.

Received: September 10, 2015

Published online: October 16, 2015