


Tetraalkylammonium Salts as Hydrogen-Bonding Catalysts

Seiji Shirakawa,* Shiyao Liu, Shiho Kaneko, Yusuke Kumatabara, Airi Fukuda, Yumi Omagari, and Keiji Maruoka*

Abstract: Although the hydrogen-bonding ability of the α hydrogen atoms on tetraalkylammonium salts is often discussed with respect to phase-transfer catalysts, catalysis that utilizes the hydrogen-bond-donor properties of tetraalkylammonium salts remains unknown. Herein, we demonstrate hydrogen-bonding catalysis with newly designed tetraalkylammonium salt catalysts in Mannich-type reactions. The structure and the hydrogen-bonding ability of the new ammonium salts were investigated by X-ray diffraction analysis and NMR titration studies.

Tetraalkylammonium salts are recognized as representative organocatalysts^[1] and are often used as phase-transfer catalysts for the activation of anionic nucleophiles through the formation of an ion pair with an ammonium cation.^[2] Although the structures of tetraalkylammonium salts are commonly expressed as **1a**, their actual ionic structure is





discussed differently.^[3–5] The positive charge of ammonium salts is delocalized on the α hydrogen atoms, which are known to interact with an anionic counterion through hydrogen bonding, as shown in **1A**. Reetz and co-workers proved the delocalization of the positive charge in tetraalkylammonium salts by X-ray crystal-structure analysis of tetrabutylammonium salts, such as tetrabutylammonium enolate and phenoxide.^[3] Furthermore, DFT calculations support the delocalized structures of ammonium salts, which include chiral

ammonium salts.^[4,5] The interaction between α hydrogen atoms on the chiral tetraalkylammonium salt catalyst and the enolate oxygen atom was thought to be important in the transition-state model of asymmetric phase-transfer reactions.^[5] However, despite the interesting hydrogen-bonding ability of the α hydrogen atoms on tetraalkylammonium salts, catalysis that could utilize such properties is, to the best of our knowledge, still unknown. Herein, we report the development of tetraalkylammonium salts as hydrogen-bonding catalysts that function on the basis of the characteristic properties of the α hydrogen atoms on the catalyst.^[6]

For efficient hydrogen-bonding catalysis, we designed new tetraalkylammonium salts **2**, which were readily prepared by the methylation of the commercially available 2,6-piperidinecarboxylate **3** (Scheme 1). The carboxylate groups at the α carbon atoms of ammonium salts **2** enhance the hydrogen-bonding ability of the α hydrogen atoms. Furthermore, the six-membered piperidine ring fixes the acidic α hydrogen atoms in an arrangement that is appropriate for bidentate binding to an anionic group.^[7]

The X-ray crystal structure of ammonium iodide **2a** provided important structural information about the newly prepared tetraalkylammonium salts **2** (Figure 1).^[8] As expected, hydrogen-bonding interactions between the α hydrogen atoms and the counteranion (I^-) were clearly observed, and the iodide anion was bound by three α hydrogen atoms, including one α hydrogen atom of the methyl group. Although a similar binding mode was also observed in



Scheme 1. Design and synthesis of a tetraalkylammonium salt for use as a hydrogen-bonding catalyst.

[*] Prof. Dr. S. Shirakawa, S. Liu, S. Kaneko, Y. Kumatabara, A. Fukuda, Y. Omagari

Department of Environmental Science, Graduate School of Fisheries and Environmental Sciences, Nagasaki University
1-14 Bunkyo-machi, Nagasaki 852-8521 (Japan)
E-mail: seijishirakawa@nagasaki-u.ac.jp

Homepage: <http://seijishirakawa.wix.com/greenchemistry>

Prof. Dr. K. Maruoka

Department of Chemistry, Graduate School of Science
Kyoto University
Sakyo, Kyoto 606-8502 (Japan)
E-mail: maruoka@kuchem.kyoto-u.ac.jp

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201508659>.




Figure 1. X-ray crystal structures of tetraalkylammonium iodides **2a** and **4**.

the crystal structure of the piperidine-derived ammonium iodide **4**,^[9] the distances between the α hydrogen atoms and the iodide anion were very different, and shorter in **2a**. These results indicate that the additional carboxylate moieties of **2a** enhance the hydrogen-bonding ability of the α hydrogen atoms.

The catalytic ability of **2** as a hydrogen-bonding catalyst was investigated in a Mannich-type reaction of *N*-acyl isoquinoline **6a**, which was generated in situ from 2,2,2-trichloroethyl chloroformate (TrocCl) and isoquinoline, as a benchmark reaction (Table 1).^[10,11] In the absence of a catalyst, the reaction with ketene silyl acetal **7a** proceeded slowly at -78°C to give product **8aa** in 7% yield after 3 h (Table 1, entry 1). When, ammonium iodide **2a** (10 mol%) was used as a catalyst, the reaction was promoted to a moderate extent (38% yield; Table 1, entry 2). On the other hand, almost no catalyst acceleration was observed when the reaction was performed with ammonium iodide **4** (Table 1, entry 3) or tertiary amine **5** (entry 4). These results clearly indicate that both the quaternary ammonium moiety and the carboxylate groups in **2a** were important in promoting the reaction. Catalyst **2b** with tetrakis[3,5-bis(trifluoromethyl)phenyl]borate (BArF^-) as a noncoordinating counteranion promoted the reaction more efficiently: Product **8aa** was formed in 61% yield within 3 h (Table 1, entry 5), and in 90% yield within 6 h (entry 6). The reaction profiles with ammonium salts **2a**, **2b**, and **4** are shown in Figure 2.

To obtain further information about ammonium salts **2** as hydrogen-bonding catalysts, we performed the reaction with catalyst **2a** or **2b** in the presence of tetrabutylammonium chloride (Table 1, entries 7 and 8). In these experiments, strong inhibition of the reaction was observed. We expect that anion exchange occurred between **2** and tetrabutylammonium chloride, and that the chloride anion coordinated (relatively) strongly to the α hydrogen atoms of catalysts **2**.

Table 1: Effect of different catalysts on the Mannich-type reaction of *N*-acyl isoquinoline **6a** with **7a**.^[a]

Entry	Catalyst	Yield [%] ^[b]
1	None	7
2	2a	38
3	4	9
4	5	6
5	2b	61
6 ^[c]	2b	90
7	2a + $\text{Bu}_4\text{N}^+\text{Cl}^-$ (20 mol%)	10
8	2b + $\text{Bu}_4\text{N}^+\text{Cl}^-$ (20 mol%)	23
9	PhCO_2H	16
10	TsOH	12

[a] Reaction conditions: **6a** (0.20 mmol), **7a** (0.30 mmol), catalyst (0.020 mmol, 10 mol%), THF (4.0 mL), -78°C , 3 h. [b] Yield of the isolated product **8aa**. [c] The reaction was performed for 6 h.





Figure 2. Reaction profiles.

Consequently, catalysts **2** could not efficiently activate substrate **6a** (Scheme 2). Catalysis of the reaction by representative Brønsted acids, such as benzoic acid and *p*-toluenesulfonic acid, was also examined. Under these conditions, product **8aa** was formed in only 16 and 12% yield, respectively (Table 1, entries 9 and 10). Although the acidity of the α hydrogen atoms of catalysts **2** was not much higher than that of the representative Brønsted acids, the reaction was efficiently promoted by catalysts **2** through multidentate hydrogen bonding.

To gain evidence for the hydrogen-bonding interaction between the α hydrogen atoms of **2** and the chloride substrate, we performed NMR titration studies of **2b** with chlorodiphenylmethane (**9**) as a relatively stable chloride compound (Figure 3). As a result of the titration with **9**, clear upfield



Scheme 2. Proposed inhibition mode with tetrabutylammonium chloride.

shifts of the NMR signals for the tetraalkylammonium core of **2b** were observed in both the ^1H and ^{13}C NMR spectrum in CDCl_3 . In the ^1H NMR titration studies, the signals for the α hydrogen atoms of **2b** (H_b , H_c , and H_d in Figure 3a) were shifted more significantly than those for the hydrogen atoms of the methyl esters (H_a in Figure 3a). Similar trends were observed in ^{13}C NMR spectra, and the carbon-atom signals





Figure 3. NMR titration studies of **2b**.

that shifted the most were those of the α carbon atoms of **2b** (Figure 3b). Almost no chemical-shift changes were observed for the carbon atoms of the BARF^- counteranion in the ^{13}C NMR titration studies.


The scope of the Mannich-type reaction of *N*-acyl isoquinolines **6** under the catalysis of ammonium salts **2** was investigated (Scheme 3). The catalysts **2** accelerated the reactions of **6a** with ketone-derived silyl enol ethers **7b** and **7c** to give the corresponding products **8ab** and **8ac**. Reactions of 3-methylisoquinoline derivative **6b** and 8-bromoisoquinoline derivative **6c** were also efficiently accelerated by catalyst **2b** to afford products **8ba** and **8ca** in good yield. On the other hand, poor acceleration by catalysts **2** was observed in reactions of 5-nitroisoquinoline derivative **6d** to produce product **8da**, probably owing to the suppression of coordina-



Scheme 3. Scope and limitation of the Mannich-type reaction of *N*-acyl isoquinolines **6**.

tion to the chloride atom in **6** as a result of hydrogen-bonding interactions between catalysts **2** and the nitro group of **6** (Scheme 3).

To expand the utility of hydrogen-bonding catalyst **2b**, we also examined a reaction of 2-methylpyridine (**10**; Scheme 4).^[12] The reaction was efficiently promoted by catalyst **2b** to give product **11** in high yield with high regioselectivity. This transformation is a valuable example of the highly regioselective dearomatization of a pyridine derivative.^[13]



Scheme 4. Regioselective reaction of 2-methylpyridine (**10**).

In summary, we have introduced the concept of tetraalkylammonium salts as hydrogen-bonding catalysts. The structure and binding ability of newly designed tetraalkylammonium salts **2** were investigated by X-ray diffraction analysis and NMR titration studies. The hydrogen-bond-donor ability of catalysts **2** was evaluated in Mannich-type reactions of *N*-acyl isoquinolines **6**. Furthermore, catalyst **2b** promoted a highly regioselective reaction of 2-methylpyridine (**10**). We are currently investigating further applications of catalysts **2** as well as the development of new (chiral) hydrogen-bonding catalysts of this type.

Acknowledgements

This research was partially supported by a Grant-in-Aid for Scientific Research on Innovative Areas ("Advanced Molecular Transformations by Organocatalysts") from MEXT (Japan), a Grant-in-Aid for Specially Promoted Research from JSPS, and Nagase Science and Technology Foundation.

Keywords: acidity · hydrogen bonds · hydrogen-bonding catalysts · organocatalysis · tetraalkylammonium salts

How to cite: *Angew. Chem. Int. Ed.* **2015**, *54*, 15767–15770
Angew. Chem. **2015**, *127*, 15993–15996

- [1] For reviews on organocatalysis, see: a) P. I. Dalko, L. Moisan, *Angew. Chem. Int. Ed.* **2001**, *40*, 3726; *Angew. Chem.* **2001**, *113*, 3840; b) P. I. Dalko, L. Moisan, *Angew. Chem. Int. Ed.* **2004**, *43*, 5138; *Angew. Chem.* **2004**, *116*, 5248; c) D. W. C. MacMillan, *Nature* **2008**, *455*, 304; d) *Comprehensive Enantioselective Organocatalysis: Catalysts, Reactions, and Applications* (Ed.: P. I. Dalko), Wiley-VCH, Weinheim, **2013**.
[2] For reviews on phase-transfer catalysis, see: a) E. V. Dehmlow, S. S. Dehmlow, *Phase Transfer Catalysis*, 3rd ed., VCH, Weinheim, **1993**; b) C. M. Starks, C. L. Liotta, M. Halpern, *Phase-*

Transfer Catalysis, Chapman & Hall, New York, **1994**; c) *Handbook of Phase-Transfer Catalysis* (Eds.: Y. Sasson, R. Neumann), Blackie Academic & Professional, London, **1997**; d) *Phase-Transfer Catalysis* (Ed.: M. E. Halpern), American Chemical Society, Washington, DC, **1997** (ACS Symposium Series 659); e) M. J. O'Donnell, *Aldrichimica Acta* **2001**, *34*, 3; f) T. Ooi, K. Maruoka, *Angew. Chem. Int. Ed.* **2007**, *46*, 4222; *Angew. Chem.* **2007**, *119*, 4300; g) *Asymmetric Phase-Transfer Catalysis* (Ed.: K. Maruoka), Wiley-VCH, Weinheim, **2008**; h) S. Shirakawa, K. Maruoka, *Angew. Chem. Int. Ed.* **2013**, *52*, 4312; *Angew. Chem.* **2013**, *125*, 4408.

- [3] a) M. T. Reetz, S. Hütte, R. Goddard, *J. Am. Chem. Soc.* **1993**, *115*, 9339; b) M. T. Reetz, S. Hütte, R. Goddard, *J. Phys. Org. Chem.* **1995**, *8*, 231; c) M. T. Reetz, S. Hütte, R. Goddard, C. Robyr, *Chem. Eur. J.* **1996**, *2*, 382; d) M. T. Reetz, S. Hütte, R. Goddard, *J. Prakt. Chem.* **1999**, *341*, 297; e) R. Goddard, H. M. Herzog, M. T. Reetz, *Tetrahedron* **2002**, *58*, 7847.
[4] a) M. T. Reetz, *Angew. Chem. Int. Ed. Engl.* **1988**, *27*, 994; *Angew. Chem.* **1988**, *100*, 1026; b) C. E. Cannizzaro, K. N. Houk, *J. Am. Chem. Soc.* **2002**, *124*, 7163.
[5] a) T. Ohshima, T. Shibuguchi, Y. Fukuta, M. Shibasaki, *Tetrahedron* **2004**, *60*, 7743; b) T. C. Cook, M. B. Andrus, D. H. Ess, *Org. Lett.* **2012**, *14*, 5836; c) T. Kamachi, K. Yoshizawa, *Org. Lett.* **2014**, *16*, 472.
[6] For reviews on hydrogen-bonding catalysis, see: a) P. R. Schreiner, *Chem. Soc. Rev.* **2003**, *32*, 289; b) P. M. Pihko, *Angew. Chem. Int. Ed.* **2004**, *43*, 2062; *Angew. Chem.* **2004**, *116*, 2110; c) A. G. Doyle, E. N. Jacobsen, *Chem. Rev.* **2007**, *107*, 5713; d) T. J. Auvil, A. G. Schafer, A. E. Mattson, *Eur. J. Org. Chem.* **2014**, 2633.
[7] It was difficult to prepare and isolate the *trans* isomer of ammonium salt **2**.
[8] The crystal structure of **2a** has been deposited at the Cambridge Crystallographic Data Centre (CCDC 1423221). These data can be obtained free of charge from The Cambridge Crystallographic Data Centre.
[9] W. A. Henderson, V. G. Young, Jr., P. Fylstra, H. C. De Long, P. C. Trulove, *Cryst. Growth Des.* **2006**, *6*, 1645.
[10] a) M. S. Taylor, N. Tokunaga, E. N. Jacobsen, *Angew. Chem. Int. Ed.* **2005**, *44*, 6700; *Angew. Chem.* **2005**, *117*, 6858; b) A. G. Schafer, J. M. Wieting, T. J. Fisher, A. E. Mattson, *Angew. Chem. Int. Ed.* **2013**, *52*, 11321; *Angew. Chem.* **2013**, *125*, 11531; c) J. M. Wieting, T. J. Fisher, A. G. Schafer, M. D. Visco, J. C. Gallucci, A. E. Mattson, *Eur. J. Org. Chem.* **2015**, 525.
[11] a) M. Takamura, K. Funabashi, M. Kanai, M. Shibasaki, *J. Am. Chem. Soc.* **2000**, *122*, 6327; b) Y. Yamaoka, H. Miyabe, Y. Takemoto, *J. Am. Chem. Soc.* **2007**, *129*, 6686; c) S. E. Reisman, A. G. Doyle, E. N. Jacobsen, *J. Am. Chem. Soc.* **2008**, *130*, 7198; d) F. Kniep, S. H. Jungbauer, Q. Zhang, S. M. Walter, S. Schindler, I. Schnapperelle, E. Herdtweck, S. M. Huber, *Angew. Chem. Int. Ed.* **2013**, *52*, 7028; *Angew. Chem.* **2013**, *125*, 7166; e) A. Berkessel, S. Das, D. Pekel, J. M. Neudörfl, *Angew. Chem. Int. Ed.* **2014**, *53*, 11660; *Angew. Chem.* **2014**, *126*, 11846; f) M. Zurro, S. Asmus, S. Beckendorf, C. Mück-Lichtenfeld, O. García Mancheño, *J. Am. Chem. Soc.* **2014**, *136*, 13999.
[12] a) K. Akiba, A. Ohtani, Y. Yamamoto, *J. Org. Chem.* **1986**, *51*, 5328; b) T. Itoh, M. Miyazaki, K. Nagata, A. Ohsawa, *Heterocycles* **1997**, *46*, 83; c) S. Yamada, T. Misono, M. Ichikawa, C. Morita, *Tetrahedron* **2001**, *57*, 8939; d) O. García Mancheño, S. Asmus, M. Zurro, T. Fischer, *Angew. Chem. Int. Ed.* **2015**, *54*, 8823; *Angew. Chem.* **2015**, *127*, 8947.
[13] J. A. Bull, J. J. Mousseau, G. Pelletier, A. B. Charette, *Chem. Rev.* **2012**, *112*, 2642.

Received: September 15, 2015

Published online: November 13, 2015