

Diazo Compounds

Deutsche Ausgabe: DOI: 10.1002/ange.201508804
Internationale Ausgabe: DOI: 10.1002/anie.201508804

Metal-Free Ring-Expansion Reaction of Six-membered Sulfonylimines with Diazomethanes: An Approach toward Seven-Membered Enesulfonamides


An-Jie Xia[†], Tai-Ran Kang,^{*} Long He,^{*} Lian-Mei Chen, Wen-Ting Li[†], Jin-Liang Yang,^{*} and Quan-Zhong Liu^{*}

Abstract: A new metal-free, ring-expansion reaction of six-membered *N*-sulfonylimines with unstable diazomethanes, generated in situ from the *N*-tosylhydrazones, has been developed. This reaction delivers valuable seven-membered enesulfonamides by a Tiffeneau–Demjanov rearrangement and intramolecular proton transfer tautomerization process. Moreover, this ring-expansion reaction can be carried out in a one-pot fashion and scaled up to the gram scale by using aryl aldehydes, without the need to isolate the *N*-tosylhydrazone.

Seven-membered sulfonamides or sulfamates are found to possess prominent biological activities,^[1] such as, calcium sensing receptor agonists,^[1c] HIV-1 protease inhibitors,^[1d] and ASBT inhibitors.^[1e] In addition, they are also used as synthetic intermediates for the synthesis of biologically active molecules.^[1c,2–5] Despite being appealing structural motifs, only few metal-catalyzed methods for the synthesis of these compounds have been reported.^[1f,2–5] Major synthetic methods consist of: 1) Rhodium- or iron-catalyzed intramolecular allylic C–H amination;^[2] 2) Rhodium-catalyzed intramolecular alkyne or allene amination;^[3,4] 3) Copper-catalyzed intramolecular nucleophilic ring-opening of sulfamate-derived aziridines;^[5] 4) Scandium- or lutetium-catalyzed haloamino-cyclization of primary sulfamate ester derivatives.^[1f] Despite great achievements in metal-catalyzed reactions, the discovery of efficient metal-free reactions will also be of great importance because of the reduced toxicity and elimination of the cost of the metal.^[9a] For these reasons, the metal-free reaction is highly desirable, and indeed, significant progress

has been made in the past years.^[6,8–10] The metal-free coupling reactions of diazo compounds generated in situ from *N*-tosylhydrazones under basic reaction conditions (Bamford–Stevens reaction)^[7] have appeared in the recent literature.^[8] In pioneering studies, Barluenga, Valdés, and co-workers discovered metal-free reductive cross-coupling of boronic acids,^[9a] phenols, and alcohols^[9b] with *N*-tosylhydrazones. The group of Wang also reported a metal-free reaction to convert *N*-tosylhydrazones into pinacol boronates.^[10]

The diazo carbon insertion into either a keto C–C bond or formyl C–H bond is a valuable tool in organic synthesis.^[11,12] For example, the Lewis acid catalyzed ring-expansion reaction of cyclohexanones with diazo compounds have become a classical strategy for synthesis of seven-membered rings through the Tiffeneau–Demjanov rearrangement (Scheme 1 a).^[13,14] Although it is clear that the reaction between



Scheme 1. The ring-expansion reaction of cyclohexanone or cyclic *N*-sulfonylimine. Ts = 4-toluenesulfonyl.

acyclic *N*-tosylimines and acyldiazomethanes could afford *N*-tosyldiazoketamines^[15a] or *N*-tosylaziridines^[15b] under various reaction conditions (Scheme 1 b), we reasoned that if the ring-expansion reaction were conducted with a six-membered imine, the resulting product might form a seven-membered imine, based on a similar Tiffeneau–Demjanov rearrangement mechanism under certain reaction conditions (Scheme 1 c).

The ring-expansion reactions between cyclic ketones and diazo compounds have been widely used in organic synthesis.^[16] However, to the best of our knowledge, there has been no systematic study on the ring-expansion reaction of

[*] Dr. A.-J. Xia,^[†] Prof. Dr. T.-R. Kang, Prof. Dr. L. He, Prof. Dr. L.-M. Chen, Prof. Dr. Q.-Z. Liu
Chemical Synthesis and Pollution Control Key Laboratory of Sichuan Province/College of Chemistry & Chemical Engineering, Collaborative Innovation Center of Tissue Repair Material of Sichuan Province, China West Normal University
Nanchong, 637009 (China)
E-mail: kangtairan@sina.com
longhe@cwnu.edu.cn
quanzhongliu@sohu.com

Prof. Dr. T.-R. Kang, Dr. W.-T. Li,^[†] Prof. Dr. J.-L. Yang
State Key Laboratory of Biotherapy/Collaborative Innovation Center of Biotherapy, Cancer Center, West China Hospital
Sichuan University, Chengdu (P.R. China)
E-mail: jlyang01@163.com

[†] These authors contributed equally to this work.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/ange.201508804>.

cyclic imines with diazo compounds so far. Herein, we report a metal-free ring-expansion reaction of six-membered imines with diazomethanes, thus providing a new and promising route to seven-membered enesulfonamides.


To test our hypothesis, we employed the *N*-tosylhydrazone **1a** and benzoxathiazine 2,2-dioxide **2a**^[17] as model substrates (for structures see Table; see Table S1 in the Supporting Information). After some optimization work (entries 1–12), it was found that the best yield of the expected seven-membered sulfonamide **3a** was achieved by employing Cs₂CO₃ as a base in 1,4-dioxane at 60 °C (entry 2). It should be noted that the diazoketamine and aziridine products were not found.

With the optimal reaction conditions established, a variety of *N*-tosylhydrazones (**1**) derived from aldehydes and benzoxathiazine 2,2-dioxides (**2**) were subjected to the ring-expansion reaction (Table 1). The electronic nature of substituents on the aryl ring attached on the α -position of *N*-tosylhydrazone had no apparent effects on the reaction. Aromatic groups bearing electron-donating and electron-

withdrawing substituents were tolerated. The higher reactivity of the substrates containing an electron-withdrawing group at the 4-position of the aryl ring could lead to formation of the products in higher yields (**3b–g**). Steric hindrance had a slight effect on this transformation, both *ortho*- and *meta*-substituted phenyl *N*-tosylhydrazones presented lower reactivity and led to the formation of the products in moderate yields (**3j–o**). Notably, 1-naphthyl *N*-tosylhydrazone participated in the reaction to give the corresponding product **3p** in 81 % yield. The heteroaromatic 3-thiophenyl *N*-tosylhydrazone provided the expected products **3q** in 45 % yields (entry 17), but the 2-furanyl or 2-thiophenyl *N*-tosylhydrazones failed (entries 18–19). To our delight, when R² = methyl, the corresponding products **3r–t** were afforded in moderate yield (entries 20–22). Using an *N*-tosylhydrazone derived from an aliphatic aldehyde as a substrate resulted in no new products (entry 23). Considering that the alkyl diazomethane generated in situ from the corresponding *N*-tosylhydrazones at 60 °C are highly unstable, we turned to the direct utilization of alkyl diazomethanes. By using diazomethane as a substrate, the same seven-membered product **3u** was afforded at 0 °C in excellent yield, irrespective of R² being a hydrogen or methyl group (entries 24 and 25). The use of diazoethane gave the product **3v** at room temperature in 90 % yield (entry 26).







This reaction was not significantly affected by the substituents on the aromatic ring of the benzoxathiazine 2,2-dioxides **2** (Table 2). Although a slightly lower yield was observed when the substituent was 8-Me or 6,8-*t*Bu (**4h**, **4i**), both electron-donor and electron-withdrawing substituents were effective, thus giving the corresponding products **4** in moderate to good yield (**4a–j**). The reaction also worked well with a 1-naphthyl *N*-sulfonylimine (**4k**).

Table 1: Reaction scope of the *N*-tosylhydrazones.^[a]

				
Entry	R ¹	R ²	3	Yield [%] ^[b]
1	Ph	H	3a	87
2	4-FC ₆ H ₄	H	3b	95
3	4-ClC ₆ H ₄	H	3c	94
4	4-BrC ₆ H ₄	H	3d	92
5	4-CF ₃ C ₆ H ₄	H	3e	89
6	4-CNC ₆ H ₄	H	3f	90
7	4-NO ₂ C ₆ H ₄	H	3g	88
8	4-MeC ₆ H ₄	H	3h	70
9	4-MeOC ₆ H ₄	H	3i	64
10	3-BrC ₆ H ₄	H	3j	78
11	3-MeC ₆ H ₄	H	3k	62
12	3-MeOC ₆ H ₄	H	3l	60
13	2-ClC ₆ H ₄	H	3m	68
14	2-BrC ₆ H ₄	H	3n	65
15	2-MeOC ₆ H ₄	H	3o	63
16	1-naphthyl	H	3p	81
17	3-thienyl	H	3q	45
18	2-thienyl	H	—	— ^[c]
19	2-furanyl	H	—	— ^[c]
20	Ph	Me	3r	64
21	4-BrC ₆ H ₄	Me	3s	70
22	3-BrC ₆ H ₄	Me	3t	67
23	<i>n</i> Bu	H	—	— ^[d]
24 ^[e]	CH ₂ N ₂	H	3u	99
25 ^[e]	CH ₂ N ₂	Me	3u	98
26 ^[f]	CH ₃ CHN ₂	Me	3v	90


[a] Reaction conditions: *N*-tosylhydrazone (**1**; 0.21 mmol), cyclic *N*-sulfonylimine (**2**; 0.2 mmol), and Cs₂CO₃ (75 mol %) in 0.5 mL of 1,4-dioxane at 60 °C for 8–16 h under argon. [b] Yield of isolated product. [c] Other products were obtained; see the Supporting Information. [d] No new products were detected. [e] The CH₂N₂ was used. See the Supporting Information for details. [f] The CH₃CHN₂ was used. See the Supporting Information for details.

Table 2: Scope of the cyclic *N*-sulfonylimines.^[a]

	
	
1a	2
4	
	R = F, 4a , 84% ^[b] R = Br, 4b , 79% R = Me, 4c , 89% R = OMe, 4d , 80%
	R = Cl, 4e , 70% R = Me, 4f , 77% R = OMe, 4g , 73%
	R ¹ = Me, R ² = H, 4h , 60% R ¹ = R ² = <i>t</i> Bu, 4i , 61%
	4j , 75%
	4k , 71%

[a] Reaction conditions: *N*-tosylhydrazone **1a** (0.21 mmol), cyclic *N*-sulfonylimines **2** (0.2 mmol), Cs₂CO₃ (75 mol %) in 0.5 mL of 1,4-dioxane at 60 °C for 8–16 h in argon. [b] Yield of isolated product.

This ring-expansion reaction could be carried out in a one-pot fashion and scaled-up to a gram scale from the aryl aldehydes **5** without the need to isolate the tosylhydrazones, thus affording the enesulfonamides **3** in similar yield (see the Supporting Information for details; selected examples are shown in Table 3).



Scheme 2. Proposed mechanism for the ring-expansion reaction.

Table 3: One-pot reaction of the aryl aldehydes **5** with cyclic N-sulfonylimine **2a**.^[a]

Entry	Ar	3	Yield [%] ^[b]
1	Ph	3a	86
2	4-FC ₆ H ₄	3b	90
3	3-BrC ₆ H ₄	3j	75
4	2-ClC ₆ H ₄	3m	65
5	2-MeOC ₆ H ₄	3o	60
6	1-naphthyl	3p	75
7 ^[c]	Ph (10.5 mmol)	3a (2.18 g)	80

[a] Reaction conditions: 1) aryl aldehyde **5** (0.21 mmol), tosylhydrazide (0.22 mmol) in MeOH (1 mL), at 60 °C for 1–2 h; 2) cyclic N-sulfonylimine **2a** (0.2 mmol) and Cs₂CO₃ (75 mol%) in 1,4-dioxane (1 mL), at 60 °C for 8–16 h under argon. [b] Yield of isolated product. [c] 10 mmol scale; see the Supporting Information for the reaction conditions.





Figure 1. X-ray structure of **3o**; thermal ellipsoids shown at 30% probability.

The structure of **3o** was confirmed by X-ray crystal structure analysis (Figure 1).^[18] Based on the Bamford-Stevens reaction^[7] and the mechanistic study reported by the groups of Barluenga^[9a] and Maruoka,^[15] the formation of **3a** is rationalized by a ring-expansion reaction as outlined in Scheme 2. The compound **1a** undergoes deprotonation to form the tosylhydrazone salt **A**. And the diazo compound **B** is generated by decomposition of **A**. From **B**, two pathways are possible: a) **B** could react with **2a** via a similar Tiffeneau-Demjanov-type intermediate **C**, and a rearrangement of the carbon skeleton with release of N₂, subsequently affords **F**. b) The carbene **D**, generated by thermally induced N₂ release, reacts with **2a** via a zwitterionic intermediate **E**, thus smoothly affording **F** by a rearrangement of the carbon



Scheme 3. Transformations of **3a**. Boc = *tert*-butoxycarbonyl, dba = dibenzylideneacetone, DCC = dicyclohexylcarbodiimide, DCM = dichloromethane, THF = tetrahydrofuran.

skeleton. Finally, intramolecular proton transfer tautomerization of **F** leads to the expected product **3a**.

The synthetic versatility of the product **3a** is demonstrated by a series of transformations (Scheme 3). Hydrogenation of **3a** using Pd/C gave the cyclic sulfamate **6** in 90% yield. 2-(2-Amino-1-phenylethyl)phenol (**7**), which is an important structural unit in the agricultural and pharmaceutical agents,^[19] was easily derived from the cyclic sulfonamide **6** in 85% yield. Acidic hydrolysis of **3a** produced the valuable 3-phenylbenzofuran (**8**) in 70% yield. Palladium-catalyzed hydroamination reaction between dimethyl 2-vinylcyclopropane-1,1-dicarboxylate and **3a** produced the allylic sulfonamide **9** in 95% yield. This approach is an effective and valuable hydroamination protocol. The N-Boc-(*S*)-2-amino-2-phenylacetic acid reacting with **3a** afforded **10** in 95% yield. The diazo carbon atom could be selectively inserted into the N–H bond of **3a**, thus giving **11** in 99% yield.

In conclusion, we developed a new metal-free ring-expansion reaction of cyclic N-sulfonylimines with diazo-methanes by a tandem reaction similar to a Tiffeneau-Demjanov rearrangement/proton-transfer tautomerization process. This transformation represents an extremely simple way to afford seven-membered enesulfonamides. Moreover, the reaction can be conducted in one pot and on gram scale using aryl aldehydes without isolation of the tosylhydrazones. We believe that this new ring-expansion reaction could become a widely used transformation in organic synthesis.

Acknowledgments

We thank Science & Technology Department of Sichuan Province (2013JY0095), Education Department of Sichuan Province (15CZ0016), the Innovative Research Team in College of Sichuan Province (14TD0016), and NSFC (21572183) for financial support.

Keywords: diazo compounds · heterocycles · hydrazones · rearrangements · ring expansion

How to cite: *Angew. Chem. Int. Ed.* **2016**, *55*, 1441–1444
Angew. Chem. **2016**, *128*, 1463–1466

- [1] For reviews on sulfamates and their therapeutic potential, see: a) J.-Y. Winum, A. Scozzafava, J.-L. Montero, C. T. Supuran, *Med. Res. Rev.* **2005**, *25*, 186; b) S. J. Williams, *Expert Opin. Ther. Pat.* **2013**, *23*, 79. For selected examples of the potent bioactive seven-membered sulfonamides, see: c) L. Kiefer, T. Gorojankina, P. Dauban, H. Faure, M. Ruat, R. H. Dodd, *Bioorg. Med. Chem. Lett.* **2010**, *20*, 7483; d) A. K. Ganguly, S. S. Alluri, D. Caroccia, D. Biswas, C.-H. Wang, E. Kang, Y. Zhang, A. T. McPhail, S. S. Carroll, C. Burlein, V. Munshi, P. Orth, C. Strickland, *J. Med. Chem.* **2011**, *54*, 7176; e) M. B. Tollefson, S. A. Kolodziej, T. R. Fletcher, W. F. Vernier, J. A. Beaudry, B. T. Keller, D. B. Reitz, *Bioorg. Med. Chem. Lett.* **2003**, *13*, 3727; f) Y. Cai, P. Zhou, X. Liu, J. Zhao, L. Lin, X. Feng, *Chem. Eur. J.* **2015**, *21*, 6386.
- [2] a) D. N. Zalatan, J. D. Bois, *J. Am. Chem. Soc.* **2008**, *130*, 9220; b) S. M. Paradine, M. C. White, *J. Am. Chem. Soc.* **2012**, *134*, 2036.
- [3] a) A. R. Thornton, S. B. Blakey, *J. Am. Chem. Soc.* **2008**, *130*, 5020; b) N. Mace, A. R. Thornton, S. B. Blakey, *Angew. Chem. Int. Ed.* **2013**, *52*, 5836; *Angew. Chem.* **2013**, *125*, 5948; c) R. D. Grigg, J. W. Rigoli, S. D. Pearce, J. M. Shomaker, *Org. Lett.* **2012**, *14*, 280; d) R. A. Brawn, K. Zhu, J. S. Panek, *Org. Lett.* **2014**, *16*, 74.
- [4] a) A. R. Thornton, V. I. Martin, S. B. Blakey, *J. Am. Chem. Soc.* **2009**, *131*, 2434; b) A. H. Stoll, S. B. Blakey, *J. Am. Chem. Soc.* **2010**, *132*, 2108; c) G. C. Feast, L. W. Page, J. Robertson, *Chem. Commun.* **2010**, *46*, 2835; d) C. S. Adams, L. A. Boralsky, I. A. Guzei, J. M. Schomaker, *J. Am. Chem. Soc.* **2012**, *134*, 10807; e) L. A. Boralsky, D. Marston, R. D. Grigg, J. C. Hersherberger, J. M. Schomaker, *Org. Lett.* **2011**, *13*, 1924.
- [5] a) F. Duran, L. Leman, A. Ghini, G. Burton, P. Dauban, R. H. Dodd, *Org. Lett.* **2002**, *4*, 2481; b) G. Malik, A. Estéoule, P. Retailleau, P. Dauban, *J. Org. Chem.* **2011**, *76*, 7438.
- [6] For selected examples of metal-free reactions, see: a) C. Sun, H. Li, D. Yu, M. Yu, X. Zhou, X. Lu, K. Huang, S. Zheng, B. Li, Z. Shi, *Nat. Chem.* **2010**, *2*, 1044; b) E. Shirakawa, K. Itoh, T. Higashino, T. Hayashi, *J. Am. Chem. Soc.* **2010**, *132*, 15537.
- [7] W. R. Bamford, T. S. Stevens, *J. Chem. Soc.* **1952**, 4735.
- [8] For selected reviews on N-tosylhydrazones, see: a) J. Barluenga, C. Valdés, *Angew. Chem. Int. Ed.* **2011**, *50*, 7486; *Angew. Chem.* **2011**, *123*, 7626; b) Z. Shao, H. Zhang, *Chem. Soc. Rev.* **2012**, *41*, 560; c) Q. Xiao, Y. Zhang, J. Wang, *Acc. Chem. Res.* **2013**, *46*, 236; d) Z. Liu, J. Wang, *J. Org. Chem.* **2013**, *78*, 10024; e) Y. Xia, Y. Zhang, J. Wang, *ACS Catal.* **2013**, *3*, 2586; f) Y. Zhang, J. Wang, *Top. Curr. Chem.* **2012**, *327*, 239; g) J. R. Fulton, V. K. Aggarwal, J. de Vicente, *Eur. J. Org. Chem.* **2005**, 1479. For selected examples of N-tosylhydrazones, see: h) S. Xu, G. Wu, F. Ye, X. Wang, H. Li, X. Zhao, Y. Zhang, J. Wang, *Angew. Chem. Int. Ed.* **2015**, *54*, 4669; *Angew. Chem.* **2015**, *127*, 4752; i) W. Xiong, C. Qi, H. He, L. Ouyang, M. Zhang, H. Jiang, *Angew. Chem. Int. Ed.* **2015**, *54*, 3084; *Angew. Chem.* **2015**, *127*, 3127; j) Z. Liu, H. Tan, L. Wang, T. Fu, Y. Xia, Y. Zhang, J. Wang, *Angew. Chem. Int. Ed.* **2015**, *54*, 3056; *Angew. Chem.* **2015**, *127*, 3099; k) Z. Zhang, Q. Zhou, W. Yu, T. Li, G. Wu, Y. Zhang, J. Wang, *Org. Lett.* **2015**, *17*, 2474; l) S. Mao, Y.-R. Gao, X.-Q. Zhu, D.-D. Guo, Y.-Q. Wang, *Org. Lett.* **2015**, *17*, 1692.
- [9] a) J. Barluenga, M. Tomás-Gamasa, F. Aznar, C. Valdés, *Nat. Chem.* **2009**, *1*, 494; b) J. Barluenga, M. Tomás-Gamasa, F. Aznar, C. Valdés, *Angew. Chem. Int. Ed.* **2010**, *49*, 4993; *Angew. Chem.* **2010**, *122*, 5113.
- [10] H. Li, L. Wang, Y. Zhang, J. Wang, *Angew. Chem. Int. Ed.* **2012**, *51*, 2943; *Angew. Chem.* **2012**, *124*, 2997.
- [11] a) W. Li, J. Wang, X. L. Hu, K. Shen, W. T. Wang, Y. Y. Chu, L. L. Lin, X. H. Liu, X. M. Feng, *J. Am. Chem. Soc.* **2010**, *132*, 8532; b) L. Gao, B. C. Kang, G.-S. Hwang, D. H. Ryu, *Angew. Chem. Int. Ed.* **2012**, *51*, 8322; *Angew. Chem.* **2012**, *124*, 8447; c) T. Hashimoto, Y. Naganawa, K. Maruoka, *J. Am. Chem. Soc.* **2011**, *133*, 8834; d) V. L. Rendina, D. C. Moebius, J. S. Kingsbury, *Org. Lett.* **2011**, *13*, 2004; e) W. Li, X. Liu, X. Hao, Y. Cai, L. Lin, X. Feng, *Angew. Chem. Int. Ed.* **2012**, *51*, 8644; *Angew. Chem.* **2012**, *124*, 8772.
- [12] a) Y. Xia, P. Qu, Z. Liu, R. Ge, Q. Xiao, Y. Zhang, J. Wang, *Angew. Chem. Int. Ed.* **2013**, *52*, 2543; *Angew. Chem.* **2013**, *125*, 2603; b) D. M. Allwood, D. C. Blakemore, S. V. Ley, *Org. Lett.* **2014**, *16*, 3064.
- [13] For reviews, see: a) T. Ye, M. A. McKerver, *Chem. Rev.* **1994**, *94*, 1091; b) M. P. Doyle, T. Ye, M. A. McKerver, *Modern Catalytic Methods for Organic Synthesis with Diazo Compounds*, Wiley, New York, **1998**.
- [14] a) H. J. Dauben, Jr., H. J. Ringold, R. H. Wade, D. L. Pearson, A. G. Anderson, Jr., *Org. Synth. Coll.* **1963**, *4*, 221; b) T. Hashimoto, Y. Naganawa, K. Maruoka, *J. Am. Chem. Soc.* **2009**, *131*, 6614. And see references therein.
- [15] a) N. Jiang, Z. Ma, Z. Qu, X. Xing, L. Xie, J. Wang, *J. Org. Chem.* **2003**, *68*, 893; b) V. K. Aggarwal, M. Ferrar, C. J. O'Brien, A. Thompson, R. V. H. Jones, R. Fieldhouse, *J. Chem. Soc. Perkin Trans. 1* **2001**, 1635.
- [16] B. M. Trost, J. Xie, N. Maulide, *J. Am. Chem. Soc.* **2008**, *130*, 17258.
- [17] Examples of cyclic N-sulfonylimines employed as substrates. See: a) P. Hu, J. Hu, J. Jiao, X. Tong, *Angew. Chem. Int. Ed.* **2013**, *52*, 5319; *Angew. Chem.* **2013**, *125*, 5427; b) H. Wang, T. Jiang, M.-H. Xu, *J. Am. Chem. Soc.* **2013**, *135*, 971; c) Y. Luo, H. B. Hepburn, N. Chotsaeng, H. W. Lam, *Angew. Chem. Int. Ed.* **2012**, *51*, 8309; *Angew. Chem.* **2012**, *124*, 8434; d) Y. Luo, A. J. Carnell, H. W. Lam, *Angew. Chem. Int. Ed.* **2012**, *51*, 6762; *Angew. Chem.* **2012**, *124*, 6866; e) Y.-Q. Wang, Y. Zhang, H. Dong, J. Zhang, J. Zhao, *Eur. J. Org. Chem.* **2013**, 764; f) A. G. Kravina, J. Mahatthananchai, J. W. Bode, *Angew. Chem. Int. Ed.* **2012**, *51*, 9433; *Angew. Chem.* **2012**, *124*, 9568; g) N. D. Litvinas, B. H. Brodsky, J. D. Bois, *Angew. Chem. Int. Ed.* **2009**, *48*, 4513; *Angew. Chem.* **2009**, *121*, 4583; h) M. Rommel, T. Fukuzumi, J. W. Bode, *J. Am. Chem. Soc.* **2008**, *130*, 17266; i) Y. Liu, T.-R. Kang, Q.-Z. Liu, L.-M. Chen, Y.-C. Wang, J. Liu, Y.-M. Xie, J.-L. Yang, L. He, *Org. Lett.* **2013**, *15*, 6090.
- [18] CCDC 1423532 (**3o**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [19] a) Y.-Q. Wang, C.-B. Yu, D.-W. Wang, X.-B. Wang, Y.-G. Zhou, *Org. Lett.* **2008**, *10*, 2071; b) M. T. Reetz, *Angew. Chem. Int. Ed. Engl.* **1991**, *30*, 1531; *Angew. Chem.* **1991**, *103*, 1559.

Received: September 19, 2015

Revised: November 13, 2015

Published online: December 14, 2015