

Total Synthesis

Deutsche Ausgabe: DOI: 10.1002/ange.201510861
Internationale Ausgabe: DOI: 10.1002/anie.201510861

Total Synthesis of Isodaphnolongamine H: A Possible Biogenetic Conundrum

Amit Kumar Chattopadhyay, Vu Linh Ly, Shashidhar Jakkepally, Gilles Berger, and Stephen Hanessian*

Abstract: Herein we describe the first synthetic efforts toward the total synthesis of isodaphnolongamine H, a calyciphylline B-type alkaloid. The strategy employs a chemoenzymatic process for the preparation of a functionalized cyclopentanol with a quaternary center. This molecule is elaborated to form an enantiopure 1-aza-perhydrocyclopentylene core, representing rings A and E of all calyciphylline B-type alkaloids. Further transformations involve the formation of a cyclic enaminone, 1,4-conjugate addition with a cyclopentenyl subunit, and intramolecular aldol cyclization to achieve a pentacyclic intermediate, ultimately forming isodaphnolongamine H in a total of 24 steps from the commercially available compound 2-carbethoxycyclopentanone. Isodaphnolongamine H exhibits promising inhibitory activity against a panel of human cancer cell lines.

The *Daphniphyllum* alkaloids are among the structurally most diverse group of polyazacyclic natural products belonging to a single genus of the family *Daphniphyllaceae*.^[1] To date, over 300 different structurally distinct alkaloids have been isolated and characterized, representing complex polyazacyclic cage-like architectures. Besides their biological activities,^[1] their biosynthesis, starting with mevalonic acid and proceeding via squalene dialdehyde to progressively complex intermediates, is a fascinating example of the ingenuity of nature. Following pioneering efforts by Suzuki, Yamamura, and co-workers,^[2] a unifying biosynthetic pathway to prepare the *Daphniphyllum* alkaloids was proposed by Heathcock and Ruggeri.^[3] This seminal contribution paved the way to the elegant total syntheses of several members of this family by the Heathcock group.^[4] Inspired by these landmark feats in the total synthesis of complex *Daphniphyllum* alkaloids, a number of groups have reported creative approaches toward the synthesis of a variety of core structures.^[5] However, efforts toward the total synthesis of other complex *Daphniphyllum* alkaloids have been sparse. Only relatively recently have the total syntheses of daphmanidin E, daphenylline, and calyciphylline N been reported by the groups of Carreira,^[6] Li,^[7] and Smith,^[8] respectively.

In 2003, Kobayashi and Morita isolated calyciphylline B (1) from the leaves of *D. calycinum* and the tentative structure

was assigned by NMR spectroscopic analysis (Figure 1).^[9] In the same year, deoxycalyciphylline B (2) and deoxyisocalyciphylline B (3) were isolated from the stem of *D. subverticillatum* by Yue and Yang.^[10] In 2009, Hao and co-workers reported the isolation of daphnolongamine H (4), a new calyciphylline B-type alkaloid with an unprecedented C6/C7 cis-ring junction, from the leaf extracts of the evergreen tree *D. longeracemosum* Rosenth.^[11] The structure and stereochemistry of daphnolongamine H was proposed based on NMR spectroscopy and its biogenetic relationship with deoxycalyciphylline B, whose structure had been confirmed by X-ray crystallography.^[10] We now report the total synthesis of isodaphnolongamine H, the biogenetically related 5-*epi* isomer of daphnolongamine H (Figure 1).




Figure 1. Representative calyciphylline B-type alkaloids and their synthetic analogues.

The biosynthetic pathway proposed by Yue and Yang^[10] for deoxycalyciphylline B and deoxyisocalyciphylline B, which differ only in the spatial disposition of the C5 methyl group, presents a possible conundrum (Figure 2). Thus, it is proposed that biosynthetic carbocation intermediate **A**, harboring *cis*-oriented hydrogens at the C6 and C7 positions, loses a hydrogen atom to give the neutral tetrasubstituted olefin intermediate **B**, which would capture a proton in an undefined process to give carbocation **C**, containing *trans*-oriented hydrogens at the C6 and C7 positions. Lactone formation with the appended propionic acid chain would deliver deoxycalyciphylline B (2) and deoxyisocalyciphylline B (3).

[*] Dr. A. K. Chattopadhyay, V. L. Ly, Dr. S. Jakkepally, Dr. G. Berger, Prof. Dr. S. Hanessian
Department of Chemistry, Université de Montréal
Station Centre Ville, C.P. 6128, Montréal, QC, H3C 3J7 (Canada)
E-mail: stephen.hanessian@umontreal.ca

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201510861>.




Figure 2. Proposed biosynthetic pathway to deoxycalciphylline B and deoxyisocalciphylline B^[10] and the anticipated pathway to form daphlongamine H (natural) and isodaphlongamine H (synthetic). The structures of intermediates A–C were redrawn in the perspectives shown to correspond to the drawing of the natural products. For the original drawings, see Ref. [10].

We propose that daphlongamine H can result from the direct C5 capture of the carbocation **A** with the propionic acid chain. Although not isolated from the extracts of the same plant source as yet, one would also expect lactonization of carbocation **A** to provide the 5-*epi* isomer, that is, isodaphlongamine H (**5**), in analogy with the isolation of deoxycalciphylline B (**2**) and its 5-*iso* epimer (**3**). In this respect our synthetic isodaphlongamine H could be the “missing” fourth component in the biosynthetically related calciphylline B-type quartet of *Daphniphyllum* alkaloids.

The unique hexacyclic framework harboring an unprecedented C6/C7 *cis*-fused stereochemistry in the deoxycalciphylline B family, as well as the intriguing biosynthetic intermediates, encouraged us to undertake the total synthesis of daphlongamine H (**4**) and its 5-*epi* isomer (**5**). We were also cognizant that a strategy which would produce a common advanced intermediate could also be applicable toward the total synthesis of the biogenetically related deoxycalciphylline B (**2**) and deoxyisocalciphylline B (**3**) (Figure 2). The hexacyclic framework of daphlongamine H contains eight stereogenic carbon atoms of which one is quaternary at the C8 position. A schematic representation of the key bond-forming reactions is shown in Figure 3. We assumed that an enolate alkylation and an intramolecular cyclization would be used to access rings A and E. The central ring B could be generated from a cyclic enaminone which would undergo 1,4-conjugate addition with a cyclopentenyl organometallic subunit. Subsequent intramolecular aldol cyclization of a keto aldehyde would generate the pentacyclic framework harboring rings A–E. Finally, a late-stage lactonization would provide daphlongamine H and/or isodaphlongamine H.





Figure 3. Key synthetic steps toward daphlongamine H (**4**).


We commenced our synthetic efforts with 2-carbethoxycyclopentanone (**6**) which was transformed to enantiopure cyclopentanol **7** in two consecutive steps using a known chemoenzymatic process^[12] (Scheme 1). Swern oxidation of **7** provided β -ketoester **8** in 96% yield. A diastereoselective alkylation of the corresponding Na enolate with triflate **9** prepared from D-lactic acid afforded a 1:1 inseparable mixture of **10** in 41% yield.^[13,14] Reduction of **10** was best achieved under Luche conditions to give the corresponding cyclopentanol as a single isomer after chromatographic separation, which was then converted into diol **11** by reduction of the ester with DIBAL-H. Bis-mesylation, followed by selective monoazidation using Bu₄NN₃ in toluene afforded **12** in 76% yield over two steps. Treatment of **12**



Scheme 1. Synthesis of tricyclic enaminone **16**. Reagents and conditions: a) (COCl)₂, (Me)₂SO, CH₂Cl₂, Et₃N, –78 to 0 °C, 96%; b) NaHMDS, toluene, –78 to –40 to –78 °C, then **9**, –78 to –40 °C, 41% yield (75% brsm), 1:1 mixture of diastereomers; c) NaBH₄, CeCl₃·7 H₂O, MeOH, 0 °C, 43%; d) DIBAL-H, CH₂Cl₂, –78 °C, 69%; e) MsCl, pyridine, DMAP, CH₂Cl₂, RT, 92%; f) Bu₄NN₃, toluene, RT, 83%; g) PPh₃, THF, 1 N aqueous NaOH, RT, then (Boc)₂O, 91%; h) DIBAL-H, CH₂Cl₂, –78 to –40 °C, 75%; i) Dess–Martin periodinane, CH₂Cl₂, RT, 91%; j) Ethynyl MgBr, THF, 0 °C to RT; k) Dess–Martin periodinane, CH₂Cl₂, RT, 85% (two steps); l) formic acid, NaI, RT, then evaporated to dryness, then K₂CO₃, MeOH, RT, 80%. HMDS = 1,1,1,3,3,3-hexamethyldisilazane, THF = tetrahydrofuran, DIBAL-H = diisobutylaluminum hydride, Ms = methane sulfonyl, DMAP = 4-dimethylaminopyridine, Boc = *tert*-butyl carbonyl; brsm = based on recovered starting material.


under Staudinger conditions led to the primary amine, which underwent in situ intramolecular cyclization to give the azaoctahydropentalene core unit **13** as the N-Boc derivative in 91 % yield. DIBAL-H reduction of the ethyl ester, followed by Dess–Martin oxidation of the corresponding alcohol, afforded aldehyde **14** which was treated with ethynylmagnesium bromide, and the resulting alcohol was oxidized to ynone **15**. Cyclization in the presence of formic acid, NaI and K₂CO₃ according to the procedure of Georg et al.^[15] afforded the cyclic enaminone **16** with an overall yield of 68 % for the three steps.

The synthesis of the cyclopentene subunit corresponding to ring D started with the known enantiopure alcohol **18** (Scheme 2).^[16] A Johnson–Claisen rearrangement in the presence of catalytic propionic acid at 145 °C led to the homoallylic ester which was reduced with DIBAL-H, and the resulting alcohol **19** was protected as the TBS ether **20** in a maximum of 37 % overall yield.



Scheme 2. Synthesis of the cyclopentene subunit. Reagents and conditions: a) CH₃C(OCH₃)₃, propionic acid, 145 °C; b) DIBAL-H, CH₂Cl₂, –78 to –40 °C, 25–40 % (two steps); c) TBSCl, Et₃N, CH₂Cl₂, RT, 93 %. TBS = *tert*-butyl dimethyl silyl.


With the tricyclic enaminone **16** and cyclopentenyl iodide **20** in hand, we proceeded with the intended 1,4-conjugate addition. Thus, treatment of the iodide **20** with *n*BuLi generated the vinyl lithium reagent **21**, which was added to the enaminone **16** in the presence of BF₃·Et₂O and CuBr·Me₂S, based on related precedents by Comins and co-workers,^[17] to give adduct **22** in 92 % yield and 90 % d.e. (Scheme 3). Silyl deprotection to the alcohol **23** gave a single diastereomer which was oxidized with PCC to the keto-aldehyde intermediate **24** in 59 % yield over two steps. Following several trials with various bases, we found that TBD (1,5,7-triazabicyclo[4.4.0]dec-5-ene) led to intramolecular aldol cyclization to give **25** in 90 % yield.^[18] A single crystal X-ray structure analysis confirmed the stereochemistry of the cyclization product including the *cis*-orientation at the B/C ring junction.^[14,19] Treatment of **25** with *p*-TSA in 1,1,2-trichloroethane at 65 °C resulted in smooth elimination to give the enone **26** which was subjected to conjugate reduction with L-Selectride to afford the pentacyclic intermediate **27** in 42 % yield for two steps. Addition of MeLi afforded exclusively the tertiary alcohol **28** whose structure and stereochemistry were determined by X-ray crystallography of the corresponding methiodide salt.^[14,19] Considering the *cis*-junction of rings B and C in **27**, it is not surprising that the trajectory of approach of the MeLi reagent toward the carbonyl group favored the less hindered α -face. In view of this result, we chose to complete the synthesis of isodaphlongamine H (**5**). Thus, the allylic double bond in **28** was hydroborated to the primary alcohol and the latter was oxidized with PCC to the lactol which was further trans-




Scheme 3. Synthesis of isodaphlongamine H. Reagents and conditions: a) **20**, *n*BuLi, Et₂O, –78 °C; then CuBr·Me₂S, BF₃·Et₂O, THF, –78 °C, then **16**, –78 °C to RT, 92 %; b) (±)-CSA, CH₂Cl₂, MeOH, RT, 91 %; c) PCC, CH₂Cl₂, RT, 65 %; d) TBD, THF, RT, 90 %; e) *p*-TSA, 1,1,2-trichloroethane, 65 °C, 70 %; f) L-Selectride, Et₂O, –78 to –20 °C, 60 %; g) MeLi, THF, –78 to +10 °C, 85 %; h) 9-BBN, THF, RT; then 2 N NaOH, H₂O₂, RT; i) PCC, CH₂Cl₂, RT, 65 % (two steps). CSA = camphorsulfonic acid, PCC = pyridinium chlorochromate, TBD = 1,5,7-triazabicyclo[4.4.0]dec-5-ene, *p*-TSA = *p*-toluenesulfonic acid, 9-BBN = 9-borabicyclo[3.3.1]nonane; L-Selectride = lithium tri-*sec*-butylborohydride.

formed in situ to lactone **5**. An X-ray crystal structure analysis of **5** confirmed the absolute stereochemistry.^[14,19] It is interesting to note that in the crystal structure of deoxycalciphylline B, ring C adopts a boat conformation,^[14] whereas the crystal structure of our synthetic isodaphlongamine H (or 6-*epi*-deoxycalciphylline B) shows a chair conformation (Figure 4).^[14] DFT calculations also suggest that isodaphlongamine H is 3.3 kcal mol^{–1} more stable than deoxycalciphylline B.^[14]

Considering the proposed biosynthetic pathway, it is intriguing that intermediate **A** would preferentially give the tetrasubstituted intermediate olefin **B**, which would be subject to severe A^{1,3}-strain only to reprotonate to **C**, then to cyclize to deoxycalciphylline B (**2**) and deoxyisocalciphylline B (**3**). In an attempt to reproduce the synthetic equivalent of the proposed tetrasubstituted intermediate **B** in the biosynthetic pathway^[10] (Figure 2), we investigated the tertiary alcohol **28** under a variety of elimination reaction conditions. These led exclusively to the exocyclic methylene product **29** (Scheme 4). All attempts to isomerize it to the endocyclic pentacycle **30** corresponding to the intermediate **B** in the biosynthesis




Isodaphlongamine H



Deoxycalcyciphylline B

Figure 4. ORTEP diagrams of isodaphlongamine H and deoxycalcyciphylline B.^[9]



Scheme 4. Attempted synthesis of the tetrasubstituted intermediate **B** reported by Yue and Yang (Ref. [10]). Reagents and conditions:

a) $\text{BF}_3 \cdot \text{Et}_2\text{O}$, CH_2Cl_2 , 80%; or SOCl_2 , pyridine, THF, -50 to $+10^\circ\text{C}$, 85%; b) oxalic acid, toluene, 115°C ; or *p*-TSA, 1,1,2-trichloroethane, 115°C .

failed, most likely because of severe $\text{A}^{1,3}$ -strain. DFT calculations concur with these results, showing that structure **30** is $2.5 \text{ kcal mol}^{-1}$ higher in energy compared to **29**.^[14] The value for the corresponding propionic acid (represented by intermediate **B** in Figure 2) is $3.7 \text{ kcal mol}^{-1}$ higher compared to that of an exocyclic methylene isomer.^[14]

Clearly, the biosynthetic steps involving the transformation of C6/C7 *cis*-pentacyclic intermediate **A** to the *trans* isomer **C** as found in deoxycalcyciphylline B and its 5-*iso* congener (Figure 2) remain unresolved^[9] and warrant further

study. Based on the carbocation intermediate **A** proposed by Yue, it seems highly probable that daphlongamine H and isodaphlongamine H can be formed by direct lactonization.

The only biological activity in this unique family of hexacyclic alkaloids has been reported for calyciphylline B ($\text{IC}_{50} = 12 \mu\text{M}$ against L1210 cells).^[9] With the new synthetic alkaloid isodaphlongamine H (**5**; 6-*epi*-deoxycalcyciphylline B) in hand, we have obtained *in vitro* data on a panel of NCI human cancer cell lines. Preliminary studies showed that isodaphlongamine H exhibited good cytotoxicity against HOP-92 (lung), SNB-75 (central nervous system), MDA_MB-435 (melanoma), and UO-31 (renal) cell lines with GI_{50} (50% growth inhibition) values of 38, 35, 48, and $43 \mu\text{M}$ respectively.^[14] The natural product deoxycalcyciphylline B, tested for the first time, was found to be only two times more active than isodaphlongamine H.

In conclusion, we have reported a highly convergent total synthesis of isodaphlongamine H accomplished in 24 linear steps from the commercially available 2-carbethoxycyclopentanone **6**. Isodaphlongamine H, the 5-*epi* isomer of daphlongamine H (the only *cis*-fused naturally occurring *Daphniphyllum* alkaloid isolated to date among the calyciphylline B family), is believed to be the missing component in this quartet of structurally unique hexacyclic alkaloids. Their hitherto unreported antitumor activities against a variety of cancer cell lines warrants further effort toward the total synthesis of related congeners.

Acknowledgements

We thank Steven Cummings (McGill University) and Gabriel Meilleur (Université de Montréal) as undergraduate research participants for Baker's yeast resolution experiments. We are thankful to Dr. Michel Simard for X-ray crystallography and to Dr. Cédric Malveau for high-resolution NMR studies. We are also very grateful to Professor Yue for an X-ray quality sample of deoxycalcyciphylline B as a reference compound. Anticancer activity was determined at the National Cancer Institute (Washington, D.C.) for which we are highly appreciative.

Keywords: alkaloids · deoxycalcyciphylline B · enaminones · natural products · total synthesis

How to cite: *Angew. Chem. Int. Ed.* **2016**, *55*, 2577–2581
Angew. Chem. **2016**, *128*, 2623–2627

- [1] a) B. Kang, P. Jakubec, D. J. Dixon, *Nat. Prod. Rep.* **2014**, *31*, 550–562; b) H. Wu, X. Zhang, L. Ding, S. Chen, J. Yang, X. Xu, *Planta Med.* **2013**, *79*, 1589–1598; c) J. Kobayashi, T. Kubota, *Nat. Prod. Rep.* **2009**, *26*, 936–962; d) C. H. Heathcock, *Angew. Chem. Int. Ed. Engl.* **1992**, *31*, 665–681; *Angew. Chem.* **1992**, *104*, 675–691.
- [2] a) K. T. Suzuki, S. Okuda, H. Niwa, M. Toda, Y. Hirata, S. Yamamura, *Tetrahedron Lett.* **1973**, *14*, 799–802; b) H. Niwa, Y. Hirata, K. T. Suzuki, S. Yamamura, *Tetrahedron Lett.* **1973**, *14*, 2129–2132.
- [3] R. B. Ruggeri, C. H. Heathcock, *Pure Appl. Chem.* **1989**, *61*, 289–292.

- [4] a) C. H. Heathcock, J. A. Stafford, D. L. Clark, *J. Org. Chem.* **1992**, *57*, 2575–2585; b) C. H. Heathcock, S. K. Davidsen, S. Mills, M. A. Sanner, *J. Am. Chem. Soc.* **1986**, *108*, 5650–5651; c) C. H. Heathcock, R. B. Ruggeri, K. F. McClure, *J. Org. Chem.* **1992**, *57*, 2585–2594; d) C. H. Heathcock, J. A. Stafford, D. L. Clark, *J. Org. Chem.* **1992**, *57*, 2566–2574; e) C. H. Heathcock, J. C. Kath, R. B. Ruggeri, *J. Org. Chem.* **1995**, *60*, 1120–1130.
- [5] For recent examples of *Daphniphyllum* alkaloids core structure syntheses, see: a) F. Diaba, A. Martínez-Laporta, G. Coussanes, I. Fernandez, J. Bonjoch, *Tetrahedron* **2015**, *71*, 3642–3651; b) J. L. Stockdill, A. M. Lopez, A. A. Ibrahim, *Tetrahedron Lett.* **2015**, *56*, 3503–3506; c) D. Ma, H. Cheng, C. Huang, L. Xu, *Tetrahedron Lett.* **2015**, *56*, 2492–2495; d) W. Wang, G.-P. Li, S.-F. Wang, Z.-F. Shi, X.-P. Cao, *Chem. Asian J.* **2015**, *10*, 377–382; e) I. Hayakawa, K. Niida, H. Kigoshi, *Chem. Commun.* **2015**, *51*, 11568–11571; f) J.-J. Guo, Y. Li, B. Cheng, T. Xu, C. Tao, X. Yang, D. Zhang, G. Yan, H. Zhai, *Chem. Asian J.* **2015**, *10*, 865–868; g) L. Wang, C. Xu, L. Chen, X. Hao, D. Z. Wang, *Org. Lett.* **2014**, *16*, 1076–1079; h) X. Xiong, Y. Li, Z. Lu, M. Wan, J. Deng, S. Wu, H. Shao, A. Li, *Chem. Commun.* **2014**, *50*, 5294–5297; i) A. A. Ibrahim, A. N. Golonka, A. M. Lopez, J. L. Stockdill, *Org. Lett.* **2014**, *16*, 1072–1075; j) H. Li, Y. Qiu, C. Zhao, Z. Yuan, X. Xie, X. She, *Chem. Asian J.* **2014**, *9*, 1274–1277; k) D. R. Williams, P. K. Mondal, S. A. Bawel, P. P. Nag, *Org. Lett.* **2014**, *16*, 1956–1959; l) S. Dorich, J. R. Del Valle, S. Hanessian, *Synlett* **2014**, 799–804; m) S. Hanessian, S. Dorich, H. Menz, *Org. Lett.* **2013**, *15*, 4134–4137; n) B. Fang, H. Zheng, C. Zhao, P. Jing, H. Li, X. Xie, X. She, *J. Org. Chem.* **2012**, *77*, 8367–8373; o) H. Li, J. Zheng, S. Xu, D. Ma, C. Zhao, B. Feng, X. Xie, X. She, *Chem. Asian J.* **2012**, *7*, 2519–2522; p) C. Xu, L. Wang, X. Hao, D. Z. Wang, *J. Org. Chem.* **2012**, *77*, 6307–6313; q) M. Yang, I. Wang, S.-H. Wang, S.-Y. Zhang, Y.-Q. Tu, F.-M. Zhang, *Org. Lett.* **2012**, *14*, 5114–5117; r) Y. Yao, G. Liang, *Org. Lett.* **2012**, *14*, 5499–5501; s) B. Darses, I. N. Michaelides, F. Sladojevich, J. W. Ward, P. R. Rzepa, D. Dixon, *Org. Lett.* **2012**, *14*, 1684–1687; t) G. Bélanger, J. Boudreault, F. Lévesque, *Org. Lett.* **2011**, *13*, 6204–6207; u) I. Coldham, L. Watson, H. Adams, N. G. Martin, *J. Org. Chem.* **2011**, *76*, 2360–2366; v) I. Coldham, A. J. M. Burrell, H. D. S. Guerrand, N. Oram, *Org. Lett.* **2011**, *13*, 1267–1269; w) F. Sladojevich, I. N. Michaelides, B. Darses, J. W. Ward, D. J. Dixon, *Org. Lett.* **2011**, *13*, 5132–5135; x) C. Xu, Z. Liu, H. Wang, B. Zhang, Z. Xiang, X. Hao, D. Z. Wang, *Org. Lett.* **2011**, *13*, 1812–1815; y) S. Ikeda, M. Shibuya, N. Kanoh, Y. Iwabuchi, *Org. Lett.* **2009**, *11*, 1833–1836; z) T. B. Dunn, J. M. Ellis, C. C. Kofink, J. R. Manning, L. E. Overman, *Org. Lett.* **2009**, *11*, 5658–5661; aa) S. E. Denmark, R. Y. Baiazitov, S. T. Nguyen, *Tetrahedron* **2009**, *65*, 6535–6548; ab) S. E. Denmark, S. T. Nguyen, R. Y. Baiazitov, *Heterocycles* **2008**, *76*, 143–154; ac) M. Ueda, M. A. A. Walczak, P. Wipf, *Tetrahedron Lett.* **2008**, *49*, 5986–5989; ad) A. Cordero-Vargas, X. Urbaneja, J. Bonjoch, *Synlett* **2007**, 2379–2382; ae) S. E. Denmark, R. Y. Baiazitov, *J. Org. Chem.* **2006**, *71*, 593–605; af) D. Solé, X. Urbaneja, J. Bonjoch, *Org. Lett.* **2005**, *7*, 5461–5464.
- [6] M. E. Weiss, E. M. Carreira, *Angew. Chem. Int. Ed.* **2011**, *50*, 11501–11505; *Angew. Chem.* **2011**, *123*, 11703–11707.
- [7] Z. Lu, Y. Li, J. Deng, A. Li, *Nat. Chem.* **2013**, *5*, 679–684.
- [8] a) A. Shvartsbart, A. B. Smith III, *J. Am. Chem. Soc.* **2015**, *137*, 3510–3519; b) A. Shvartsbart, A. B. Smith III, *J. Am. Chem. Soc.* **2014**, *136*, 870–873.
- [9] H. Morita, J. Kobayashi, *Org. Lett.* **2003**, *5*, 2895–2898.
- [10] S.-P. Yang, J.-M. Yue, *J. Org. Chem.* **2003**, *68*, 7961–7966.
- [11] C.-S. Li, Y.-T. Di, Q. Zhang, Y. Zhang, C.-J. Tan, X.-J. Hao, *Helv. Chim. Acta* **2009**, *92*, 653–659.
- [12] a) Z. Hng, Z. Wang, K. Ding, *Adv. Synth. Catal.* **2011**, *353*, 1584–1590; b) C. A. M. Fraga, E. J. Barreiro, *Chirality* **1996**, *8*, 305–310.
- [13] For related examples of enolate alkylation, see: a) E. Vedejs, O. Dauguils, *J. Am. Chem. Soc.* **2003**, *125*, 4166–4173; b) E. Vedejs, O. Dauguils, *J. Am. Chem. Soc.* **1999**, *121*, 5813–5814; c) C. P. Decicco, D. J. Nelson, R. L. Corbett, J. C. Dreabitt, *J. Org. Chem.* **1995**, *60*, 4782–4785; d) R. V. Hoffman, H.-O. Kim, *J. Org. Chem.* **1995**, *60*, 5107–5113.
- [14] Please see the Supporting Information.
- [15] a) M. J. Niphakis, B. J. Turunen, G. I. Georg, *J. Org. Chem.* **2010**, *75*, 6793–6805; b) B. J. Turunen, G. I. Georg, *J. Am. Chem. Soc.* **2006**, *128*, 8702–8703; For recent reviews on cyclic enamines, see: c) A. K. Chattopadhyay, S. Hanessian, *Chem. Commun.* **2015**, *51*, 16437–16449; d) A. K. Chattopadhyay, S. Hanessian, *Chem. Commun.* **2015**, *51*, 16450–16467.
- [16] a) J. Li, P. Yang, M. Yao, J. Deng, A. Li, *J. Am. Chem. Soc.* **2014**, *136*, 16477–16480; b) M. Kabat, J. Kiegiel, N. Cohen, K. Toth, P. M. Wovkulich, M. R. Uskokoric, *Tetrahedron Lett.* **1991**, *32*, 2343–2346.
- [17] J. D. Brown, M. A. Foley, D. L. Comins, *J. Am. Chem. Soc.* **1988**, *110*, 7445–7447.
- [18] C. Ghobril, C. Sabot, C. Mioskowski, R. Baati, *Eur. J. Org. Chem.* **2008**, 4104–4108.
- [19] CCDC 1437874 (**25**), 1437875 (methiodide of **28**), 1437876 (**5**), and 1437873 (**2**) contain the supplementary crystallographic data for this paper. These data are provided free of charge by The Cambridge Crystallographic Data Centre.

Received: November 23, 2015

Published online: January 14, 2016