


Heterocycle Synthesis

Deutsche Ausgabe: DOI: 10.1002/ange.201510820
Internationale Ausgabe: DOI: 10.1002/anie.201510820Ruthenium-Catalyzed C–C Bond Cleavage of 2*H*-Azirines: A Formal [3+2+2] Cycloaddition to Fused Azepine Skeletons

Tengfei Li, Fen Xu, Xincheng Li, Chunxiang Wang,* and Boshun Wan*

Abstract: 2*H*-azirines can serve as three-atom synthons by C–C bond cleavage, however, it involves a high energy barrier under thermal conditions ($> 50.0 \text{ kcal mol}^{-1}$). Reported is a ruthenium-catalyzed [3+2+2] cycloaddition reaction of 2*H*-azirines with diynes, thus leading to the formation of fused azepine skeletons. This approach features an unprecedented metal-catalyzed C–C bond cleavage of 2*H*-azirines at room temperature, and the challenging construction of aza-seven-membered rings from diynes. The results of this study provide a new reaction pattern for constructing nitrogen-containing seven-membered rings and may find applications in the synthesis of other complex heterocycles.

Transition-metal-catalyzed C–C bond activation represents a straightforward but challenging strategy to discover new transformations and construct complex molecules in modern organic synthesis.^[1] In comparison with achievements in C–H bond activation, the development of C–C bond activation still lags behind because of its inert character and poor interaction of the C–C σ bond with transition metals.^[2] In contrast, C–C bond activation of small rings has gained considerable attention in the past decade. As a paradigm, 2*H*-azirines can undergo diverse ring-opening reactions by different bond-cleavage modes.^[3] To date, the chemistry of 2*H*-azirines mainly focuses on the cleavage of a C–N single bond. For example, 2*H*-azirines can serve as synthetic equivalents of vinyl nitrenes by C–N bond cleavage to construct various N-heterocycles, such as indoles,^[4] pyrroles,^[5] pyridines^[6] and pyrazines^[7] (Scheme 1 a). In contrast, reactions of 2*H*-azirines with unsaturated compounds by C–C bond cleavage have been rarely explored.^[8–10] Recently, Xiao and Lu disclosed the visible-light-induced C–C bond cleavage of 2*H*-azirines under metal-free conditions, which allowed the divergent synthesis of polysubstituted pyrroles and oxazoles by formal [3+2] cycloadditions (Scheme 1 b).^[8] These reactions are initiated by single-electron oxidation of the 2*H*-azirines in the presence of an excited photocatalyst with subsequent homolytic cleavage of the C–C bond. To the best of our knowledge,



Scheme 1. Ring-opening reactions of 2*H*-azirines. EWG = electron-withdrawing group, Ts = 4-toluenesulfonyl.

transition-metal-catalyzed C–C bond cleavage of 2*H*-azirines remains largely unexploited, possibly because of the higher energy barrier.^[11]

Azepine derivatives are important skeletal motifs which are widely distributed in numerous natural products and pharmaceuticals (Figure 1).^[12] Owing to the significant medicinal and bioactive properties of the azepine moiety, many methods for the preparation of azepine-containing heterocycles have been developed in recent decades.^[13,14] Particularly, azepine frameworks have been constructed by [4+3], [5+2], and [3+2+2] cycloadditions of unsaturated hydrocarbons with triazoles, aziridines, azomethine imines, cyclopropyl imines, imine esters, and amino ketones.^[14] Despite the significant advances made in this subject, there are only limited reports on the preparation of ring-fused azepine derivatives.^[13k,14l,m]

Cycloadditions involving diynes enable the straightforward construction of medium-sized carbo- or heterocycles with high atom efficiency.^[15] Along with our ongoing research interest in the metal-catalyzed cycloaddition reactions by

[*] T. F. Li, Dr. F. Xu, Dr. X. C. Li, Dr. C. X. Wang, Prof. Dr. B. S. Wan
Dalian Institute of Chemical Physics, Chinese Academy of Sciences
457 Zhongshan Road, Dalian 116023 (China)
E-mail: cxwang@dicp.ac.cn
bswan@dicp.ac.cn

Dr. F. Xu
Department of Material and Chemical Engineering
Zhengzhou University of Light Industry
5 Dongfeng Road, Zhengzhou 450002 (China)

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201510820>.





Figure 1. Natural products containing fused azepine skeletons.


using diynes as C_4 units,^[16] we surmized that the azepine framework could be assembled by [3+2+2] cycloaddition of diynes with 2*H*-azirines, in which the 2*H*-azirines would serve as three-atom components. Indeed, examples of the formation of aza-seven-membered rings from diynes are extremely rare.^[17] Our previous studies demonstrated that [3+2] rather than [3+2+2] cycloadducts were preferentially constructed in the reaction of diynes with methyleneaziridines, thus leaving an alkyne unit intact.^[16c] To our delight, the reaction of diynes with 2*H*-azirines follows the [3+2+2] pathway by using an appropriate ruthenium catalyst,^[18] thus providing the desired azepine architectures (Scheme 1 c). This approach features an unprecedented metal-catalyzed C–C bond cleavage of 2*H*-azirines, and the challenging construction of aza-seven-membered rings from diynes. Herein, we report our preliminary results.

At the outset, the diyne **1a** and 2*H*-azirine **2a** were chosen as model substrates for the optimization of the reaction conditions [Eq. (1)]. The results are summarized in Table S1



in the Supporting Information. Some typical transition-metal catalysts for cycloadditions were first investigated. Unfortunately, no desired product was observed with our previously developed iron catalyst^[16a] (Table S1, entry 1). In addition, cobalt,^[19a] nickel,^[19b] rhodium,^[19c] and iridium^[19d] catalysts also failed to provide any new products (entries 2–5). However, the possibility that the above systems might be workable could not be completely excluded if an in-depth investigation is done. Gratifyingly, the product 3*H*-azepine **3a** was obtained in 71 % yield using 10 mol % $[Cp^*Ru(COD)Cl]$ (entry 6; COD = 1,5-cyclooctadiene, $Cp^* = C_5Me_5$). The structure of **3a** was unambiguously confirmed by X-ray crystal diffraction.^[20] Various ruthenium (II) catalysts were then evaluated in the reaction, however, none of them led to better results (entries 7–11). Intriguingly, ruthenium(II) complexes lacking either the Cp or Cp^* ligand exhibited no appreciable catalytic activity (entries 9–11). Subsequently, a screening of solvents revealed that both DCE and DCM (DCE = 1,2-dichloroethane, DCM = dichloromethane) showed the best performance (entries 12–15). Furthermore, a simple inspection on the reaction temperatures indicated that lowering the temperature to 25 °C can afford a better yield (entry 17). A slight improvement in product yield can be achieved by prolonging the reaction time to 20 hours (80 %, entry 19). It is noteworthy that the dimerization of diynes was commonly detected under ruthenium catalysis,^[18d,e] thus slightly lowering the yield of **3a**. Remarkably, decreasing the catalyst loading to 5 mol % displayed lower activity (entry 20).

With the optimal reaction conditions secured, we turned our attention to the scope of this reaction. The variations of the R^1 group on the C=N double bond moiety of 2*H*-azirines were first examined. As highlighted in Scheme 2, both




Scheme 2. Substrate scope. Yields of isolated products are given.

electron-donating and electron-withdrawing groups on the phenyl ring could be successfully introduced, thus providing the corresponding polysubstituted azepines (**3b–e**, **3g–i**) in moderate to excellent yields. The R^1 group is placed near to the methylene unit (CH_2) in the product, as identified by the X-ray crystal diffraction of **3g**.^[20] However, the 2*H*-azirine **2f**, bearing a strong electron-withdrawing aryl group, provided **3f** in low yield. 2*H*-azirines bearing an *ortho*-substituted phenyl ring (**2h**, **2i**) or 1-naphthyl group (**2j**) reacted smoothly to afford the corresponding azepines in high yields, thus suggesting that steric hindrance is well tolerated (**3h–j**). Moreover, the 2-thienyl group was tolerated and provided the desired product **3k** albeit with moderate yield. Replacement of the aryl group with an alkyl substituent also led to the effective formation of the corresponding azepine (**3l**), although a much lower yield was observed. Notably, the reaction was sensitive to the electronic and steric effect of the R^2 substituent. 2*H*-azirine, having an electron-donating group (Me) on the *para* position of the phenyl ring (**2m**), showed higher reactivity than that bearing an electron-withdrawing group (NO₂, **2n**). When a sterically more demanding 2-methoxyphenyl-substituted 2*H*-azirine (**2o**) was employed, a satisfactory yield can still be achieved.


Next, the reactions of various terminal diynes with 2*H*-azirines were investigated. The sulfonamide-based diynes (**1a**, **1p**) could undergo this cyclization and be transformed into the corresponding cycloadducts in moderate yields. However, malonate-tethered terminal diynes were found to be less effective, thus affording the azepines in 30 and 24 % yield,

respectively (**3q** and **3r**). Interestingly, ketones, which are unsaturated groups that also undergo cycloadditions with diynes,^[18f] did not interfere with the reaction involving the azirines (**3s** and **3t**). In addition, the O-tethered diyne **1u** reacted with 2*H*-azirine **2h** to deliver **3u** in moderate yield. Unfortunately, unlike our previous observation in the ruthenium-catalyzed [2+2+2] cycloaddition,^[16b] the methylene-linked diyne was completely unreactive, thus indicating that the Thorpe–Ingold effect^[21] was crucial for this transformation (**3v**). Furthermore, when the unsymmetrical diyne **1w**, bearing a methyl group at one alkyne terminus, was subjected to the reaction, the azepine compound **3w** was obtained albeit with low yield. However, the diyne **1x**, with a phenyl group at one alkyne terminus, showed no reactivity under the optimized reaction conditions, probably because of steric hindrance.^[22]

To explore the reaction mechanism, the reaction of **1a** with deuterated 2*H*-azirine, [D]-**2a**, was first investigated [Eq. (2)]. As expected, the desired product [D]-**3a** was





obtained with more than 99% deuterium incorporation. This observation reveals that a 1,5-sigmatropic hydrogen shift occurs in the reaction. Additionally, the reaction of **1a** and **2a** in the presence of CH_3COOD did not provide any deuterated product [Eq. (3)], thus indicating an intramolecular proton-



transfer process and an irreversible C–H functionalization. Under this premise, kinetic isotope effect (KIE) studies were performed. The intermolecular competitive reactions between **2a** and [D]-**2a** with **1a** gave a KIE value of approximately 1.0 at 5 minutes (or 10 min) on the basis of ^1H NMR analysis [Eq. (4)]. These data suggest that the C–H bond cleavage is not involved in the rate-determining step.

Consequently, a plausible mechanism was outlined (Scheme 3) on the basis of the present results. The catalytic cycle starts with the oxidative cyclization of the diyne **1** on the ruthenium center to form the ruthenacyclopentatriene inter-



Scheme 3. Proposed mechanism.

mediate **A**.^[18] Then, the insertion of the imine moiety on 2*H*-azirine **2** into the intermediate **A** affords either the intermediate **B** or **B'**. The intermediate **B'** is less favored as a result of the steric repulsion between R^1 and R . Subsequently, β -carbon elimination^[23] results in C–C bond cleavage. The bicyclic [5.1.0] skeleton of **B** is expanded with the release of the strain of three-membered aziridine ring to give the eight-membered ruthenacycle intermediate **C**. The reductive elimination of **C** provides the 2*H*-azepine **D** and regenerates the catalyst. Finally, 1,5-H shift of **D** provides the thermodynamically stable product 3*H*-azepine **3**.^[24] In the case of unsymmetrical diynes **1w** and **1x**, the steric repulsion between R and R^2 may cause the diminished yields in products.^[25]

In summary, we have successfully developed a formal [3+2+2] approach to construct seven-membered heterocycles through the ruthenium-catalyzed cycloaddition of diynes with 2*H*-azirines under mild reaction conditions, in which 2*H*-azirines undergo the cleavage of C–C single bond. This transformation provides a facile and straightforward method for the synthesis of fused azepine derivatives with broad substrate scope and a wide range of functional groups. We expect that these results may provide new insight into the chemistry of 2*H*-azirines and find applications in the synthesis of complex heterocycles.

Acknowledgments

We thank NSFC (21372219, 21572225) for the financial support.


Keywords: alkynes · C–C activation · cycloaddition · heterocycles · ruthenium

How to cite: *Angew. Chem. Int. Ed.* **2016**, 55, 2861–2865
Angew. Chem. **2016**, 128, 2911–2915

- [1] For selected reviews of C–C bond cleavage, see: a) C. Jun, *Chem. Soc. Rev.* **2004**, 33, 610; b) M. Murakami, T. Matsuda, *Chem. Commun.* **2011**, 47, 1100; c) T. Seiser, T. Saget, D. N. Tran, N. Cramer, *Angew. Chem. Int. Ed.* **2011**, 50, 7740; *Angew. Chem.* **2011**, 123, 7884; d) F. Chen, T. Wang, N. Jiao, *Chem. Rev.* **2014**, 114, 8613; e) A. Dermenci, J. W. Coe, G. Dong, *Org. Chem.*

- Front.* **2014**, *1*, 567; f) H. Liu, M. Feng, X. Jiang, *Chem. Asian J.* **2014**, *9*, 3360; g) L. Soullart, N. Cramer, *Chem. Rev.* **2015**, *115*, 9410.
- [2] a) M. Murakami, Y. Ito, *Top. Organomet. Chem.* **1999**, *3*, 97; b) B. Rybtchinski, D. Milstein, *Angew. Chem. Int. Ed.* **1999**, *38*, 870; *Angew. Chem.* **1999**, *111*, 918; c) H. Li, Y. Li, X. S. Zhang, K. Chen, X. Wang, Z. J. Shi, *J. Am. Chem. Soc.* **2011**, *133*, 15244; d) L. Liu, N. Ishida, M. Murakami, *Angew. Chem. Int. Ed.* **2012**, *51*, 2485; *Angew. Chem.* **2012**, *124*, 2535; e) C. Qin, P. Feng, Y. Ou, T. Shen, T. Wang, N. Jiao, *Angew. Chem. Int. Ed.* **2013**, *52*, 7850; *Angew. Chem.* **2013**, *125*, 8004.
- [3] For reviews on reactions of 2*H*-azirines, see: a) F. Palacios, A. M. O. de Retana, E. M. de Marigorta, J. M. de los Santos, *Eur. J. Org. Chem.* **2001**, 2401; b) A. F. Khlebnikov, M. S. Novikov, *Tetrahedron* **2013**, *69*, 3363; c) C. Y. Huang, A. G. Doyle, *Chem. Rev.* **2014**, *114*, 8153.
- [4] a) D. F. Taber, W. W. Tian, *J. Am. Chem. Soc.* **2006**, *128*, 1058; b) D. A. Candito, M. Lautens, *Org. Lett.* **2010**, *12*, 3312; c) S. Jana, M. D. Clements, B. K. Sharp, N. Zheng, *Org. Lett.* **2010**, *12*, 3736.
- [5] a) X. Qi, X. Xu, C. M. Park, *Chem. Commun.* **2012**, *48*, 3996; b) A. F. Khlebnikov, M. V. Golovkina, M. S. Novikov, D. S. Yufit, *Org. Lett.* **2012**, *14*, 3768; c) C. M. Park, Y. Jiang, *Chem. Sci.* **2014**, *5*, 2347; d) T. Li, X. Xin, C. Wang, D. Wang, F. Wu, X. Li, B. Wan, *Org. Lett.* **2014**, *16*, 4806; e) L. Zhu, Y. Yu, Z. Mao, X. Huang, *Org. Lett.* **2015**, *17*, 30; f) S. K. Pawar, R. L. Sahani, R. S. Liu, *Chem. Eur. J.* **2015**, *21*, 10843.
- [6] a) Y. F. Wang, K. K. Toh, J. Y. Lee, S. Chiba, *Angew. Chem. Int. Ed.* **2011**, *50*, 5927; *Angew. Chem.* **2011**, *123*, 6049; b) N. S. Loy, A. Singh, X. Xu, C. M. Park, *Angew. Chem. Int. Ed.* **2013**, *52*, 2212; *Angew. Chem.* **2013**, *125*, 2268; c) A. Prechter, G. Henrion, P. Faudot dit Bel, F. Gagosz, *Angew. Chem. Int. Ed.* **2014**, *53*, 4959; *Angew. Chem.* **2014**, *126*, 5059; d) Y. Jiang, C. M. Park, T. P. Loh, *Org. Lett.* **2014**, *16*, 3432.
- [7] a) F. Palacios, A. M. Ochoa de Retana, J. I. Gil, R. Lopez de Munain, *Org. Lett.* **2002**, *4*, 2405; b) N. S. Y. Loy, S. Kim, C. M. Park, *Org. Lett.* **2015**, *17*, 395; c) T. Ryu, Y. Baek, P. H. Lee, *J. Org. Chem.* **2015**, *80*, 2376.
- [8] a) J. Xuan, X. Xia, T. Zeng, Z. Feng, J. Chen, L. Lu, W. Xiao, *Angew. Chem. Int. Ed.* **2014**, *53*, 5653; *Angew. Chem.* **2014**, *126*, 5759; b) T. Zeng, J. Xuan, W. Ding, K. Wang, L. Lu, W. Xiao, *Org. Lett.* **2015**, *17*, 4070.
- [9] a) B. Singh, E. F. Ullman, *J. Am. Chem. Soc.* **1967**, *89*, 6911; b) A. Padwa, J. Smolanoff, *J. Am. Chem. Soc.* **1971**, *93*, 548; c) B. Singh, A. Zweig, J. B. Gallivan, *J. Am. Chem. Soc.* **1972**, *94*, 1199; d) E. Albrecht, J. Mattay, S. Steenken, *J. Am. Chem. Soc.* **1997**, *119*, 11605; e) C. M. Nunes, I. Reva, R. Fausto, *J. Org. Chem.* **2013**, *78*, 10657.
- [10] Lewis acid promoted C–C bond cleavage of related aziridines has been reported. For selected examples see: a) P. D. Pohlhaus, R. K. Bowman, J. S. Johnson, *J. Am. Chem. Soc.* **2004**, *126*, 2294; b) P. A. Wender, D. Strand, *J. Am. Chem. Soc.* **2009**, *131*, 7528; c) Z. Chen, L. Wei, J. Zhang, *Org. Lett.* **2011**, *13*, 1170; d) L. Li, X. Wu, J. Zhang, *Chem. Commun.* **2011**, *47*, 5049; e) L. Li, J. Zhang, *Org. Lett.* **2011**, *13*, 5940.
- [11] a) T. Pinho e Melo, A. M. R. Gonsalves, *Curr. Org. Synth.* **2004**, *1*, 275; b) S. Calvo-Losada, J. J. Quirante, D. Suárez, T. L. Sordo, *J. Comput. Chem.* **1998**, *19*, 912; c) Y. Liu, P. Guan, Y. Wang, L. Liu, J. Gao, *J. Phys. Chem. A* **2015**, *119*, 67.
- [12] a) B. Renfro, G. Harrington, G. Proctor, *Heterocyclic Compounds: Azepines*, Wiley & Interscience, New York, **1984**; b) C. R. Ganellin, D. J. Triggle, *Dictionary of Pharmacological Agents*, Chapman & Hall/CRC, London, **1996**; c) D. O'Hagan, *Nat. Prod. Rep.* **1997**, *14*, 637; d) J. A. Builla, J. Vaguero, J. Barluenga, *Modern Heterocyclic Chemistry*, Wiley-VCH, Weinheim, **2011**, pp. 1865–1988; e) C. Y. Chen, D. J. Hart, *J. Org. Chem.* **1990**, *55*, 6236; f) P. N. Tariat, P. Solomon, J. Morris, P. Kershaw, S. Lilienfeld, C. Ding, *Neurology* **2000**, *54*, 2269; g) J. Kobayashi, M. Yoshinaga, N. Yoshida, M. Shiro, H. Morita, *J. Org. Chem.* **2002**, *67*, 2283; h) D. Kakuta, Y. Hitotsuyanagi, N. Matsuura, H. Fukaya, K. Takeya, *Tetrahedron* **2003**, *59*, 7779; i) S. Berkov, C. Codina, F. Viladomat, J. Bastida, *Bioorg. Med. Chem. Lett.* **2008**, *18*, 2263.
- [13] For reports on the classical methods for azepine synthesis: Heck coupling, see: a) M. Qadir, J. Cobb, P. W. Sheldrake, N. Whittall, A. J. White, K. K. Hii, P. N. Horton, M. B. Hursthouse, *J. Org. Chem.* **2005**, *70*, 1545; radicals: b) L. Yet, *Tetrahedron* **1999**, *55*, 9349; c) C. E. Masse, A. J. Morgan, J. S. Panek, *Org. Lett.* **2000**, *2*, 2571; ring expansion: d) R. M. Kellogg, T. Van Bergen, *J. Org. Chem.* **1971**, *36*, 978; e) E. J. Kantorowski, M. J. Kurth, *Tetrahedron* **2000**, *56*, 4317; f) J. Yadav, B. S. Reddy, M. K. Gupta, A. Prabhakar, B. Jagadeesh, *Chem. Commun.* **2004**, 2124; g) E. A. Ilardi, J. T. Njardarson, *J. Org. Chem.* **2013**, *78*, 9533; ring-closing metathesis: h) M. E. Maier, *Angew. Chem. Int. Ed.* **2000**, *39*, 2073; *Angew. Chem.* **2000**, *112*, 2153; i) P. Jakubec, A. Hawkins, W. Felzmann, D. J. Dixon, *J. Am. Chem. Soc.* **2012**, *134*, 17482; Cope rearrangement: j) T. Sasaki, S. Eguchi, M. Ohno, *J. Am. Chem. Soc.* **1970**, *92*, 3192; k) E. E. Schultz, V. N. G. Lindsay, R. Sarpong, *Angew. Chem. Int. Ed.* **2014**, *53*, 9904; *Angew. Chem.* **2014**, *126*, 10062.
- [14] For reports on cycloaddition routes to azepine derivatives: [4+3] cycloaddition, see: a) B. M. Trost, C. M. Marrs, *J. Am. Chem. Soc.* **1993**, *115*, 6636; b) N. D. Shapiro, F. D. Toste, *J. Am. Chem. Soc.* **2008**, *130*, 9244; c) C. S. Jeffrey, K. L. Barnes, J. A. Eickhoff, C. R. Carson, *J. Am. Chem. Soc.* **2011**, *133*, 7688; d) X. Hu, J. Chen, S. Gao, B. Feng, L. Lu, W. Xiao, *Chem. Commun.* **2013**, *49*, 7905; e) K. Liu, H. Teng, C. Wang, *Org. Lett.* **2014**, *16*, 4508; f) H. Shang, Y. Wang, Y. Tian, J. Feng, Y. Tang, *Angew. Chem. Int. Ed.* **2014**, *53*, 5662; *Angew. Chem.* **2014**, *126*, 5768; [5+2] cycloaddition: g) E. L. Stogryn, S. J. Brois, *J. Am. Chem. Soc.* **1967**, *89*, 605; h) P. A. Wender, T. M. Pedersen, M. J. Scanio, *J. Am. Chem. Soc.* **2002**, *124*, 15154; i) C. Roscini, K. L. Cubbage, M. Berry, A. J. Orr-Ewing, K. I. Booker-Milburn, *Angew. Chem. Int. Ed.* **2009**, *48*, 8716; *Angew. Chem.* **2009**, *121*, 8872; j) E. Kanno, K. Yamanoi, S. Koya, I. Azumaya, H. Masu, R. Yamasaki, S. Saito, *J. Org. Chem.* **2012**, *77*, 2142; k) M. Zhou, R. Song, C. Wang, J. Li, *Angew. Chem. Int. Ed.* **2013**, *52*, 10805; *Angew. Chem.* **2013**, *125*, 11005; l) J. J. Feng, T. Y. Lin, H. H. Wu, J. Zhang, *J. Am. Chem. Soc.* **2015**, *137*, 3787; m) J. J. Feng, T. Y. Lin, H. H. Wu, J. Zhang, *Angew. Chem. Int. Ed.* **2015**, *54*, 15854; *Angew. Chem.* **2015**, *127*, 16080; [3+2+2] cycloaddition: n) M. Zhou, R. Song, J. Li, *Angew. Chem. Int. Ed.* **2014**, *53*, 4196; *Angew. Chem.* **2014**, *126*, 4280.
- [15] M. I. Alexandrovna, B. I. Ionin, J. C. Tebby in *Alkynes in Cycloadditions* (Ed.: J. Tebby), Wiley, New Delhi, **2014**.
- [16] a) C. Wang, X. Li, F. Wu, B. Wan, *Angew. Chem. Int. Ed.* **2011**, *50*, 7162; *Angew. Chem.* **2011**, *123*, 7300; b) F. Xu, C. Wang, X. Li, B. Wan, *ChemSusChem* **2012**, *5*, 854; c) B. Pan, C. Wang, D. Wang, F. Wu, B. Wan, *Chem. Commun.* **2013**, *49*, 5073; d) C. Wang, D. Wang, F. Xu, B. Pan, B. Wan, *J. Org. Chem.* **2013**, *78*, 3065; e) F. Xu, C. Wang, D. Wang, X. Li, B. Wan, *Chem. Eur. J.* **2013**, *19*, 2252; f) C. Wang, D. Wang, H. Yan, H. Wang, B. Pan, X. Xin, X. Li, F. Wu, B. Wan, *Angew. Chem. Int. Ed.* **2014**, *53*, 11940; *Angew. Chem.* **2014**, *126*, 12134; g) F. Xu, C. Wang, H. Wang, X. Li, B. Wan, *Green Chem.* **2015**, *17*, 799.
- [17] Construction of seven-membered carbocycles from diynes: a) K. E. Schwiebert, J. M. Stryker, *J. Am. Chem. Soc.* **1995**, *117*, 8275; b) K. Maeda, S. Saito, *Tetrahedron Lett.* **2007**, *48*, 3173; c) R. A. Bauer, C. M. DiBlasi, D. S. Tan, *Org. Lett.* **2010**, *12*, 2084; d) L. Saya, I. Fernández, F. López, J. L. Mascareñas, *Org. Lett.* **2014**, *16*, 5008; e) T. Yoshida, Y. Tajima, M. Kobayashi, K. Masutomi, K. Noguchi, K. Tanaka, *Angew. Chem. Int. Ed.* **2015**, *54*, 8241; *Angew. Chem.* **2015**, *127*, 8359. Construction of

- aza-seven-membered rings from two identical alkynes, see: ref. [14n].
- [18] Ruthenium-catalyzed cycloaddition reactions of diynes: a) M. O. Albers, D. J. De Waal, D. C. Liles, D. J. Robinson, E. Singleton, M. B. Wiege, *J. Chem. Soc. Chem. Commun.* **1986**, 1680; b) C. Gemel, A. LaPensée, K. Mauthner, K. Mereiter, R. Schmid, K. Kirchner, *Monatsh. Chem.* **1997**, *128*, 1189; c) E. Rüba, K. Mereiter, R. Schmid, V. N. Sapunov, K. Kirchner, H. Schottenberger, M. J. Calhorda, L. F. Veiros, *Chem. Eur. J.* **2002**, *8*, 3948; d) Y. Yamamoto, H. Kitahara, R. Ogawa, H. Kawaguchi, K. Tatsumi, K. Itoh, *J. Am. Chem. Soc.* **2000**, *122*, 4310; e) Y. Yamamoto, H. Takagishi, K. Itoh, *J. Am. Chem. Soc.* **2002**, *124*, 28; f) Y. Yamamoto, H. Takagishi, K. Itoh, *J. Am. Chem. Soc.* **2002**, *124*, 6844; g) J. A. Varela, L. Castedo, C. Saá, *Org. Lett.* **2003**, *5*, 2841; h) J. A. Varela, L. Castedo, C. Saá, *J. Org. Chem.* **2003**, *68*, 8595; i) Y. Yamamoto, K. Kinpara, T. Saigoku, H. Takagishi, S. Okuda, H. Nishiyama, K. Itoh, *J. Am. Chem. Soc.* **2005**, *127*, 605; j) Y. Yamamoto, K. Kinpara, H. Nishiyama, K. Itoh, *Adv. Synth. Catal.* **2005**, *347*, 1913; k) Y. Yamamoto, K. Kinpara, R. Ogawa, H. Nishiyama, K. Itoh, *Chem. Eur. J.* **2006**, *12*, 5618; l) J. A. Varela, S. G. Rubín, C. González-Rodríguez, L. Castedo, C. Saá, *J. Am. Chem. Soc.* **2006**, *128*, 9262; m) S. García-Rubín, J. A. Varela, L. Castedo, C. Saá, *Chem. Eur. J.* **2008**, *14*, 9772; n) S. Medina, G. Dominguez, J. Perez-Castells, *Org. Lett.* **2012**, *14*, 4982; o) K. Yamashita, Y. Yamamoto, H. Nishiyama, *J. Am. Chem. Soc.* **2012**, *134*, 7660; p) K. Matsui, M. Shibuya, Y. Yamamoto, *ACS Catal.* **2015**, *5*, 6468.
- [19] a) K. Kase, A. Goswami, K. Ohtaki, E. Tanabe, N. Saino, S. Okamoto, *Org. Lett.* **2007**, *9*, 931; b) M. M. McCormick, H. A. Duong, G. Zuo, J. Louie, *J. Am. Chem. Soc.* **2005**, *127*, 5030; c) K. Tanaka, N. Suzuki, G. Nishida, *Eur. J. Org. Chem.* **2006**, 3917; d) G. Onodera, Y. Shimizu, J. Kimura, J. Kobayashi, Y. Evihara, K. Kondo, K. Sakata, R. Takeuchi, *J. Am. Chem. Soc.* **2012**, *134*, 10515.
- [20] CCDC 1057257 (**3a**) and 1057258 (**3g**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre.
- [21] a) R. M. Beesley, C. K. Ingold, J. F. Thorpe, *J. Chem. Soc.* **1915**, 107, 1080; b) M. E. Jung, J. Gervay, *J. Am. Chem. Soc.* **1991**, *113*, 224; c) M. E. Jung, G. Piizzi, *Chem. Rev.* **2005**, *105*, 1735.
- [22] In the case of internal diyne bearing a methyl group at each alkyne terminus, the reaction only provided trace amounts of dimer at room temperature. The diyne can be completely transformed at higher temperature (80°C), thus leading to the formation of inseparable complex mixtures along with a small amount of dimer. However, no desired azepine product can be obtained.
- [23] For β -carbon elimination, see: a) P. Kumar, K. Zhang, J. Louie, *Angew. Chem. Int. Ed.* **2012**, *51*, 8602; *Angew. Chem.* **2012**, *124*, 8730; b) P. Kumar, J. Louie, *Org. Lett.* **2012**, *14*, 2026; c) D. Crépin, J. Dawick, C. Aïssa, *Angew. Chem. Int. Ed.* **2010**, *49*, 620; *Angew. Chem.* **2010**, *122*, 630; d) M. Murakami, S. Ashida, T. Matsuda, *J. Am. Chem. Soc.* **2006**, *128*, 2166. Also see ref. [1b].
- [24] For 1,5-H shift observed in azepine rings, see: a) Y. Kubota, K. Satake, H. Okamoto, M. Kimura, *Org. Lett.* **2005**, *7*, 5215; b) Y. Kubota, K. Satake, H. Okamoto, M. Kimura, *Org. Lett.* **2006**, *8*, 5469; c) C. E. Cordonier, K. Satake, H. Okamoto, M. Kimura, *Eur. J. Org. Chem.* **2006**, 3803.
- [25] An alternative mechanism involving the terminal alkyne addition to 2H-azirine followed by cycloisomerization might be possible. To examine this possibility, we conducted the reaction of structurally similar **1a'** with **2a**. However, no new product was observed under ruthenium(II) catalyst even at higher temperature, thus excluding this pathway.



Received: November 23, 2015
Published online: January 21, 2016