

## Halogenation

Deutsche Ausgabe: DOI: 10.1002/ange.201510909  
Internationale Ausgabe: DOI: 10.1002/anie.201510909

## Reductive Chlorination and Bromination of Ketones via Trityl Hydrazones

Julius R. Reyes and Viresh H. Rawal\*

**Abstract:** A method is presented for the direct transformation of a ketone to the corresponding reduced alkyl chloride or bromide. The process involves the reaction of a ketone trityl hydrazone with *t*BuOCl to give a diazene which readily collapses to the  $\alpha$ -chlorocarbonyl radical, reduction of which by a hydrogen atom source gives the alkyl chloride product. The use of *N*-bromosuccinimide provides the corresponding alkyl bromide. This unique transformation provides a reductive halogenation that complements Barton's redox-neutral vinyl halide synthesis.

The transformation of an alcohol into a chloride, one of the most basic reactions in organic chemistry, can present unanticipated difficulties when encountered in a complex molecule. In the course of our studies toward the synthesis of *N*-methylwelwitindolinone B isothiocyanate (**1**; Figure 1), we





Figure 1. Welwitindolinone B and potential precursors.

attempted to convert the C13 hydroxy group of **2** into the required, inverted chloride.<sup>[1]</sup> Unfortunately, the neopentyl and homoallylic nature of the alcohol conspired to trigger a skeletal rearrangement, thereby foiling the planned synthetic route.<sup>[2]</sup> Others have also observed difficulties, in completely different systems, in making alkyl chlorides by Walden inversion of suitably activated precursors.<sup>[3]</sup> Given the limitations of this transformation, combined with the prevalence of natural products<sup>[4]</sup> having a chloride attached to an  $sp^3$ -hybridized carbon atom, we set forth to devise a fundamentally different solution for the synthesis of such chlorides.<sup>[5]</sup> We report here the realization of a method for the overall reductive transformation of ketones to alkyl halides.

The reductive chlorination method was conceived to provide a conceptually new way for the installation of

a chlorine atom in high-value compounds. Rather than introducing the chloride through nucleophilic displacement of an activated alcohol, with the attendant difficulties and complications noted above, the idea was to introduce the chlorine first and then, through the generation of a reactive intermediate, add a hydrogen atom. This concept was expected to be realized through the use of hydrazone chemistry (Figure 2a).<sup>[6–8]</sup> The reaction of a trityl hydrazone

a) Concept: Reductive chlorination of trityl hydrazones



b) Precedent: Barton-type alkenyl halide synthesis






Figure 2. Synthesis of organochlorides from hydrazone precursors.

with a chloronium source was expected to give a chlorodiazene (**5**),<sup>[9]</sup> which upon thermolysis would extrude  $N_2$  to generate a trityl radical and the sought after reactive intermediate, the  $\alpha$ -chlorocarbonyl radical **6**. Provided the thermolysis was carried out in the presence of a hydrogen atom donor, then diastereoselective hydrogen abstraction would give the desired chloroalkane product. Introduction of chlorine and generation and reduction of the reactive intermediate were envisioned through a single synthetic maneuver. Realized, this transformation provides a reductive chlorination from the  $C=O$  oxidation state, a method that complements Barton-type vinyl halide synthesis (Figure 2b).<sup>[10,11]</sup>

The feasibility of the above concept was evaluated using hydrazone **11a** (Figure 3a), which is available through condensation of trityl hydrazide and benzylacetone (74%).<sup>[12]</sup> Treatment of a solution of **11a** in THF with *t*BuOCl (1.1 equiv) at  $-20^\circ C$ , followed by addition of an excess of EtSH and warming to room temperature afforded the desired

[\*] J. R. Reyes, Prof. Dr. V. H. Rawal  
Department of Chemistry, The University of Chicago  
5735 South Ellis Avenue, Chicago, IL 60637 (USA)  
E-mail: vrawal@uchicago.edu

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201510909>.

a) Preliminary observations on reductive chlorination of **11a**b) Observation of thermolysis and O<sub>2</sub> capture

**Figure 3.** Optimization of reaction parameters. [a] Reactions performed in THF. Yield determined by NMR spectroscopy.

product of reductive chlorination in 37% yield (NMR).<sup>[13]</sup> Modest yields in these early reactions were balanced significantly by the product of apparent hydrolysis of the starting hydrazone, that is, benzylacetone. Mechanistic considerations suggested that the apparent hydrolysis product likely arises by way of peroxychloroalkane intermediate **14**, the product of O<sub>2</sub> capture by the  $\alpha$ -chlorocarbonyl radical (Figure 3b). Support for this hypothesis was obtained by carrying out the thermolysis in the absence of a reducing agent and placing it under an oxygen balloon prior to warming to room temperature. The major product of the reaction under these reaction conditions was benzylacetone (**10**; 57% yield by NMR), with no evidence of chloroalkane **13a**. In contrast, scrupulous exclusion of air through two freeze-pump-thaw cycles completely eliminated formation of **10** in the reaction mixture. Variable-temperature NMR experiments provided an understanding of the thermal requirements for the different steps of the reaction.<sup>[14,15]</sup> A  $-78^{\circ}\text{C}$  sample of **11a** and *t*BuOCl was examined by NMR spectroscopy in a probe precooled to  $-30^{\circ}\text{C}$ . After 10 minutes had elapsed, a reaction was observed, and the starting hydrazone was found to be fully consumed. The resulting putative chlorodiazene **12** was found to persist as the temperature was increased from  $-30^{\circ}\text{C}$  to  $-10^{\circ}\text{C}$ . Upon further warming above  $-10^{\circ}\text{C}$ , diazene **12** decomposed to give a mixture of products. With the sequence of reagent addition and temperature control guided by the above study, as well as careful O<sub>2</sub> exclusion, the reaction was optimized to furnish **13a** in 82% isolated yield (Table 1, entry 1). A brief screen of chloronium ion sources and H-atom donors offered no improvement over *t*BuOCl and EtSH, with *N*-chlorosuccinimide yielding none of the desired chloride. Less odorous, high-molecular weight thiols were examined briefly as H-atom donors, but found to give less satisfactory results.

The capability of the reductive chlorination procedure was examined in a range of substrates, as shown in Table 1. The hydrazone of phenoxylacetone was converted into the corresponding chloride in 85% yield (entry 2). Diastereoselectivity in the reduction event displayed high substrate dependence. Substrates in which the hydrazone was part of a conformationally locked six-membered ring favored axial hydrogen abstraction to give the equatorial chloride. Reductive chlorination of **11c** and **11d** gave a mixture of chlorides,

**Table 1:** Reductive chlorination of trityl hydrazones.

$\text{R} \begin{array}{c} \text{H} \\ \text{N} \end{array} \text{N} \text{CPh}_3 \xrightarrow[4:1 \text{ CyH/DCM, } -20^{\circ}\text{C}]{t\text{BuOCl (1.1 equiv)}} \text{R} \begin{array}{c} \text{H} \\ \text{Cl} \end{array} \text{R}' \xrightarrow[\text{then EtSH; warm to } 40^{\circ}\text{C}]{} \text{R} \begin{array}{c} \text{H} \\ \text{Cl} \end{array} \text{R}'$			
Entry	Hydrazone	Chloride	Yield [%] <sup>[a]</sup>
1			82
2			85 <sup>[b]</sup>
3			57 (3.4:1)
4			69 (2.9:1)
5			71 (1.3:1)
6			56 <sup>[b]</sup>
7			50 <sup>[c]</sup>
8			83 (21:1)
9			70 (2.6:1)
10			67 <sup>[d]</sup>

[a] Yield of the isolated product. Diastereomeric ratio (d.r.; given within parentheses) determined by <sup>1</sup>H NMR analysis of purified chlorides.

[b] Yield determined by NMR spectroscopy. [c] Yield for isolated diastereomer shown (major), 2.8:1 crude d.r. [d] Lithiated hydrazone

treated with dichloramine-T. Cy=cyclohexyl, DCM=dichloromethane, TBS=*tert*-butyldimethylsilyl.

with a preference for the equatorial chlorides by approximately 3:1 (entries 3 and 4). Diminished selectivity was observed for the reaction of the hydrazone of *trans*-1-decalone, wherein the three 1,3-diaxial interactions may disfavor axial hydrogen abstraction (entry 5). The neopentyl, homoallylic hydrazone **11g**, comprising the cyclohexane core of welwitindolinone B, gave a 2.8:1 mixture of chloride diastereomers, from which the major component **13g** was isolated in 50% yield. Notably, this reductive chlorination occurs without 1,2-migration of the vinyl group, possibly reflecting the radical stabilizing effect of chlorine.<sup>[16]</sup>

Among cyclopentanone-derived trityl hydrazones, the facial bias of the [2.2.1]-bridged system in **11h** engendered high selectivity for the *endo* chloride **13h**, which was isolated

in 83 % yield, with greater than 20:1 diastereoselectivity (Table 1, entry 8). The hydrazone of (–)- $\alpha$ -thujone (**11i**) gave a mixture of chlorides (**13i**) in 70 % yield, wherein hydrogen abstraction had taken place predominantly from the face opposite the  $\alpha$ -methyl substituent (entry 9).<sup>[17]</sup> The utility of this method was further demonstrated by the reductive chlorination of the sterically encumbered hydrazone **11j**, derived from *O*-methyl estrone (entry 10). The lower reactivity of **11j** to chlorination necessitated deprotonation followed by chlorination, which was achieved efficiently with dichloramine-T. The protocol afforded chloride **13j** in good yield and high selectivity for the  $\beta$ -chloride shown, with hydrogen abstraction taking place *anti* to the adjacent C13 methyl group.

The underlying concept of the reductive chlorination appeared transferrable to bromination. Thus, upon treatment of **11a** with *N*-bromosuccinimide (NBS) in place of *t*BuOCl followed by addition of EtSH and warming, it was converted into the expected reductive bromination product **16a** (Table 2). Three other hydrazones were similarly subjected

with a range of trityl hydrazones, and affords chloride products with stereoselectivities that may complement those available through ionic processes. The basic transformation was also successfully demonstrated for the synthesis of alkyl bromides.<sup>[19]</sup> The ability of this method to efficiently construct neopentyl alkyl chlorides from the carbonyl functional group provides a distinct tactic for the preparation of such halide-containing natural products, syntheses of which are of ongoing interest in our laboratories.





## Acknowledgments

Financial support from the National Cancer Institute of the NIH (R01 CA101438) is gratefully acknowledged. Additionally, we generously thank Dr. Antoni Jurkiewicz for his NMR expertise and assistance.

**Keywords:** diastereoselectivity · halogenation · hydrazones · radical reactions · synthetic methods

**How to cite:** *Angew. Chem. Int. Ed.* **2016**, *55*, 3077–3080  
*Angew. Chem.* **2016**, *128*, 3129–3132

**Table 2:** Reductive bromination of trityl hydrazones.

$\text{R}-\text{C}(=\text{O})-\text{N}(\text{CPh}_3)_2 \xrightarrow[\text{then EtSH; warm to 40 } ^\circ\text{C}]{\text{NBS (1.1 equiv)}^{[a]}, 4:1 \text{ CyH/DCM, } -20 ^\circ\text{C}}$				
Hydrazone	Bromide	Yield [%] <sup>[b]</sup>	d.r. <sup>[c]</sup>	
<b>11a</b>		65	–	
<b>11d</b>		69	2.5:1	
<b>11e</b>		49	1.1:1	
<b>11h</b>		60	17:1	

[a] NBS solubilized in THF. [b] Yield of isolated products. [c] Diastereomeric ratio (d.r.) determined by <sup>1</sup>H NMR analysis of purified bromides.

to the bromination conditions and gave the anticipated alkyl bromides in good yields. Of note, hydrogen abstraction by the  $\alpha$ -bromocarbonyl radicals gave consistently lower diastereoselectivities than that observed for their  $\alpha$ -chloro congeners, with apparent selectivity reversal in the bromination of *trans*-1-decalone (**16e**). The effect of the different halides on selectivity appears to parallel that observed for other free-radical processes, including halogenations of alkanes. The slightly higher selectivity seen for reductive chlorination versus bromination comports with greater stabilization of the radical accorded by chlorine over bromine, as reflected by C–H bond dissociation energies of simple haloalkanes.<sup>[18]</sup>

In summary, we have disclosed a novel method for the conversion of ketones into the respective saturated alkyl chlorides. The key step involves chlorination of a ketone trityl hydrazone, which upon warming fragments to give an  $\alpha$ -halo-stabilized carbonyl radical that is then reduced by EtSH to furnish the alkyl chloride product. The method is effective

- [1] a) V. Bhat, V. H. Rawal, *Chem. Commun.* **2011**, 47, 9705–9707; b) V. Bhat, J. A. MacKay, V. H. Rawal, *Tetrahedron* **2011**, 67, 10097–10104.
- [2] Both the groups of Shea and Garg have noted similar challenges in converting an alcohol into the chloride. See: a) L. Cleary, J. Pitzen, J. A. Brailsford, K. J. Shea, *Org. Lett.* **2014**, 16, 4460–4463; b) N. A. Weires, E. D. Styduhar, E. L. Baker, N. K. Garg, *J. Am. Chem. Soc.* **2014**, 136, 14710–14713.
- [3] a) L. A. Gortey, M. Vairamani, C. Djerassi, *J. Org. Chem.* **1985**, 50, 4173–4182; b) K. S. Albone, J. Macmillan, A. R. Pitt, C. L. Willis, *Tetrahedron* **1986**, 42, 3203–3214; c) D. C. Snyder, *J. Org. Chem.* **1995**, 60, 2638–2639; d) P. Korakas, S. Chaffee, J. B. Shotwell, P. Duque, J. L. Wood, *Proc. Natl. Acad. Sci. USA* **2004**, 101, 12054–12057; e) W. Pluempunapat, W. Chavasiri, *Tetrahedron Lett.* **2006**, 47, 6821–6823.
- [4] a) G. W. Gribble, *Acc. Chem. Res.* **1998**, 31, 141–152; b) G. W. Gribble, *Heterocycles* **2012**, 84, 157–207; c) acutumine: S. M. King, S. B. Herzon, *Alkaloids Chem. Biol.* **2014**, 73, 161–222; d) chlorosulpholipids: D. K. Bedke, C. D. Vanderwal, *Nat. Prod. Rep.* **2011**, 28, 15–25; e) Rhodometaceae: B.-G. Wang, J. B. Gloer, N.-Y. Ji, J.-C. Zhao, *Chem. Rev.* **2013**, 113, 3632–3685.
- [5] Synthesis of alkyl chlorides from vinyl halides: a) K. Iwasaki, K. K. Wan, A. Oppedisano, S. W. M. Crossley, R. A. Shenvi, *J. Am. Chem. Soc.* **2014**, 136, 1300–1303; b) S. M. King, X. Ma, S. B. Herzon, *J. Am. Chem. Soc.* **2014**, 136, 6884–6887; c) J. C. Lo, J. Gui, Y. Yabe, C.-M. Pan, P. S. Baran, *Nature* **2014**, 516, 343–348.
- [6] For general reviews of hydrazone chemistry: a) B. A. Rogen, A. El-Awa, “Hydrazine,” *e-EROS Encyclopedia of Reagents for Organic Synthesis* **2014**, pp. 1–7; b) A. R. Chamberlin, J. E. Sheppeck, B. Goess, C. Lee, “*p*-Toluenesulfonylhydrazide,” *e-EROS Encyclopedia of Reagents for Organic Synthesis* **2007**, pp. 1–11; c) B. Robinson, *Chem. Rev.* **1963**, 63, 373–401; d) B. Robinson, *Chem. Rev.* **1969**, 69, 227–250.
- [7] Baldwin has employed lithiated trityl hydrazones as umpoled synthons: a) J. E. Baldwin, J. C. Bottaro, J. N. Kolhe, R. M. Adlington, *J. Chem. Soc. Chem. Commun.* **1984**, 22–23; b) J. E. Baldwin, R. M. Adlington, J. C. Bottaro, A. U. Jain, J. N. Kolhe, M. W. D. Perry, I. M. Newington, *J. Chem. Soc. Chem. Commun.* **1984**, 1095–1096; c) J. E. Baldwin, R. M. Adlington, I. M.

- Newington, *J. Chem. Soc. Chem. Commun.* **1986**, 176–178; d) J. E. Baldwin, R. M. Adlington, J. C. Bottaro, J. N. Kolhe, I. M. Newington, M. W. D. Perry, *Tetrahedron* **1986**, 42, 4235–4246.
- [8] Trityl hydrazides have been shown to provide entry to acyl radicals: S. Bath, N. M. Laso, H. Lopez-Ruiz, B. Quiclet-Sire, S. Z. Zard, *Chem. Commun.* **2003**, 204–205.
- [9] a) M. W. Moon, *J. Org. Chem.* **1972**, 37, 383–385; b) M. W. Moon, *J. Org. Chem.* **1972**, 37, 386–390; c) J. M. Wyman, S. Jochum, M. Brewer, *Synth. Commun.* **2008**, 38, 3623–3630.
- [10] Synthesis of vinyl halides from hydrazone precursors: a) D. H. R. Barton, R. E. O'Brien, S. Sternhell, *J. Chem. Soc.* **1962**, 470–476; b) H. Mori, S. Wada, K. Tsuneda, *Chem. Pharm. Bull.* **1963**, 11, 684–685; c) H. Mori, K. Tsuneda, *Chem. Pharm. Bull.* **1963**, 11, 1413–1417. From *N*-silylhydrazones: d) M. E. Furrow, A. G. Myers, *J. Am. Chem. Soc.* **2004**, 126, 5436–5445.
- [11] Reductive chlorinations: a) M. Brewer, *Tetrahedron Lett.* **2006**, 47, 7731–7733; b) D. A. Mundal, J. J. Lee, R. J. Thomson, *J. Am. Chem. Soc.* **2008**, 130, 1148–1149.
- [12] See the Supporting Information.
- [13] a) Among the by-products were triphenylmethane and ethyl trityl sulfide; b) for a recent review regarding thiyl radicals, including the use of thiols as H-atom donors, see: F. Dénès, M. Pichowicz, G. Povie, P. Renaud, *Chem. Rev.* **2014**, 114, 2587–2693.
- [14] See the Supporting Information for selected <sup>1</sup>H NMR spectra (Figure S1).
- [15] Studies by Malament and co-workers regarding stereospecific 1,4-additions of Cl<sub>2</sub> gas to ketazines have shown that thermolysis (25 °C, 30 days) or photolysis (10 h) of the resulting α,α'-dichloroazoalkanes in the presence of thiophenol produces significant quantities of the radical coupling product relative to alkyl chloride: a) D. S. Malament, J. M. McBride, *J. Am. Chem. Soc.* **1970**, 92, 4586–4593; b) L. Nissim, D. S. Malament, *Isr. J. Chem.* **1974**, 12, 925–936; c) N. Levi, D. S. Malament, *J. Chem. Soc. Perkin Trans. 2* **1976**, 1249–1256.
- [16] Selected references regarding homoallyl–homoallyl rearrangements: a) M. Castaing, M. Pereyre, M. Ratier, P. M. Blum, A. G. Davies, *J. Chem. Soc. Perkin Trans. 2* **1979**, 287–292; b) A. L. J. Beckwith, D. M. O'Shea, *Tetrahedron Lett.* **1986**, 27, 4525–4528; c) G. Stork, R. Mook, *Tetrahedron Lett.* **1986**, 27, 4529–4532; d) M. Toyota, T. Wada, K. Fukumoto, M. Ihara, *J. Am. Chem. Soc.* **1998**, 120, 4916–4925; e) M. Toyota, M. Yokota, M. Ihara, *Tetrahedron Lett.* **1999**, 40, 1551–1554. As an undesired side reaction: f) M. Y. Chu-Moyer, S. J. Danishefsky, G. K. Schulte, *J. Am. Chem. Soc.* **1994**, 116, 11213–11228.
- [17] The NOESY spectrum of **13i** suggests that the stereochemistry of these chlorides has been misassigned in the following reference: J. C. Rees, D. Whittaker, *Org. Magn. Reson.* **1981**, 15, 363–369.
- [18] a) S. Furuyama, D. M. Golden, S. W. Benson, *J. Am. Chem. Soc.* **1969**, 91, 7564–7569; b) E. Tschuikow-Roux, S. Paddison, *Int. J. Chem. Kinet.* **1987**, 19, 15–24; c) E. Tschuikow-Roux, D. R. Salomon, *J. Phys. Chem.* **1987**, 91, 699–702; d) K. Miyokawa, E. Tschuikow-Roux, *J. Phys. Chem.* **1990**, 94, 715–717.
- [19] In preliminary studies, we have found that lithiation of **11a** and subsequent treatment with NFSI generated the corresponding fluoride in 33 % yield (NMR, unoptimized).

Received: November 24, 2015

Published online: January 28, 2016