



Auf diesen Seiten weisen wir auf wichtige aktuelle Beiträge in unseren Schwesterzeitschriften hin. Wenn Sie die Seiten online lesen, dann können Sie

die Artikel mit einem Klick direkt aufrufen, ansonsten sind sie durch Eingabe der DOIs über Wiley Online Library leicht online zugänglich.




### Heterogeneous Catalysis

J. E. Schmidt, C.-Y. Chen, S. K. Brand, S. I. Zones, M. E. Davis\*

Facile Synthesis, Characterization, and Catalytic Behavior of a Large-Pore Zeolite with the IWV Framework

**Microporous materials:** Synthesis of aluminosilicate IWV under industrially relevant conditions has been demonstrated, and the material can be produced in both fluoride and hydroxide media across a wide composition range. The zeolite demonstrates catalytic activity in the hydroisomerization of *n*-hexane (see figure).

Easily produced aluminosilicate IWV made in hydroxide media demonstrates catalytic activity



Chem. Eur. J.

DOI: 10.1002/chem.201504717




### Ionogels

X. Wang, Q. Yang, Y. Cao, J. Zhou, H. Hao, Y. Liang, J. Hao\*

Ionogels of a Sugar Surfactant in Ionic Liquids

**Sweetened gels:** Green and environmentally friendly ionogels are obtained through the self-assembly of a sugar surfactant in imidazolium-based ionic liquids (see figure). Microstructures from ribbons to lamellar structures are induced by temperature. The relationship between the rheological and tribological properties are described.



Chem. Asian J.

DOI: 10.1002/asia.201501198




### Protein-Protein Interactions

V. Azzarito, P. Rowell, A. Barnard, T. A. Edwards, A. Macdonald, S. L. Warriner,\* A. J. Wilson\*


Probing Protein Surfaces: QSAR Analysis with Helix Mimetics

**Rationalising behaviour:** Using a helix mimetic based on an oligoamide scaffold, we have exploited a modular synthesis to access compounds that can readily be used to understand the noncovalent determinants of hDM2 recognition by cell-active p53/hDM2 inhibitors.



ChemBioChem

DOI: 10.1002/cbic.201500504



ChemPhysChem


DOI: 10.1002/cphc.201500686

K. Mathwig, Q. Chi, S. G. Lemay, L. Rassaei\*

Handling and Sensing of Single Enzyme Molecules: From Fluorescence Detection towards Nanoscale Electrical Measurements

**Hard to handle:** Some of the major breakthroughs that have led to significant advances in classical single-molecule enzyme studies are first summarized. In addition, several recent developments in micro- and nanodevices are highlighted, which allow the study and handling of few or even single enzyme molecules in a confined reaction space

## Single Molecules



ChemMedChem



DOI: 10.1002/cmdc.201500490

M. Petrera, T. Wein, L. Allmendinger, M. Sindelar, J. Pabel, G. Höfner, K. T. Wanner\*

Development of Highly Potent GAT1 Inhibitors: Synthesis of Nipecotic Acid Derivatives by Suzuki–Miyaura Cross-Coupling Reactions

**Exploring the gap:** Guided by molecular modeling studies, a series of nipecotic acid derivatives with different 2-biphenyl moieties on an *N*-butenyl linker were synthesized as potential GAT1 inhibitors. A 2',4'-dichlorobiphenyl-2-yl derivative was found to be highly potent in binding and uptake assays and to display high subtype selectivity for GAT1.

## Medicinal Chemistry



ChemSusChem



DOI: 10.1002/cssc.201501584

Z. He, J. Liu, S. Y. Khoo, T. T. Y. Tan\*

Electropolymerization of Uniform Polyaniline Nanorod Arrays on Conducting Oxides as Counter Electrodes in Dye-Sensitized Solar Cells

**PANI for your thoughts:** A rapid and template-free method to grow highly ordered polyaniline (PANI) nanorod arrays on conducting oxide substrates is reported. The substrate, FTO, is modified with a thin layer of reduced graphene oxide, before growth of the PANI nanorods by in situ, low-potential electropolymerization. When employed as electrodes in dye-sensitized solar cells, the PANI arrays offer high electrocatalytic activity, chemical stability, and enhanced performance compared to platinum-based electrodes.

## Solar Cells



Holes transfer at {023} facet/electrolyte interface

ChemCatChem


DOI: 10.1002/cctc.201501162


B. Wang, S. Shen,\* L. Guo

Surface Reconstruction of Facet-Functionalized SrTiO<sub>3</sub> Nanocrystals for Photocatalytic Hydrogen Evolution

**Facet-rich coating:** Surface-reconstructed SrTiO<sub>3</sub> nanocrystals with disordered overlayers are created by a thermal hydrogenation process. The nanocrystals enclosed with {023} and {001} facets in an appropriate ratio show the highest photocatalytic activity for hydrogen evolution. MOR = methanol oxidation reaction; V = vacancy.

## Electrocatalysis






### Density Functional Calculations


H.-Y. T. Chen, S. Livraghi, E. Giamello, G. Pacchioni\*

Mechanism of the Cyclo-Oligomerisation of  $C_2H_2$  on Anatase  $TiO_2$  (101) and (001) Surfaces and Their Reduction: An Electron Paramagnetic Resonance and Density Functional Theory Study

**On the surface of things:** Acetylene oligomerisation occurs on the surface of anatase  $TiO_2$  leading to the formation of polycyclic aromatic hydrocarbons and a reduced titania surface. The mechanism of the reaction has been elucidated by EPR measurements and DFT calculations (see figure).



ChemPlusChem  
DOI: 10.1002/cplu.201500383




### Water Splitting


T. Li, X. Wang, W. Yuan,\* C. M. Li

Unique Co-Catalytic Behavior of Protic Ionic Liquids as Multifunctional Electrolytes for Water Splitting

**DEAFening:** For the first time, a protic ionic liquid, diethylammonium format, is used as a multifunctional electrolyte in a water-splitting cell, demonstrating a unique role as co-catalyst.



ChemElectroChem  
DOI: 10.1002/celc.201500458




### Hydrogen-Storage Materials

Z. Yang, W. Xiong, J. Wang, Y. Zhu, Y. Xia\*

A Systematic Study on the Preparation and Hydrogen Storage of Zeolite 13X-Templated Microporous Carbons

A systematic study on CVD-based strategies for the nanocasting of porous carbon materials with zeolite 13X as template is presented. The resulting carbon materials exhibit hydrogen-uptake capacities of up to 6.3 wt.-%. Linear relationships between the uptake capacity and the total surface area, the micropore volume, and the micropore surface area were found.



Eur. J. Inorg. Chem.  
DOI: 10.1002/ejic.201501180




### Natural Product Synthesis

D. A. Heredia, E. L. Larghi,\* T. S. Kaufman\*


A Straightforward Synthesis of 5-Methylaaptamine from Eugenol, Employing a  $6\pi$ -Electrocyclization Reaction of a 1-Azatriene

5-Methylaaptamine, an analogue of naturally occurring aaptamine, was synthesized from eugenol by using an  $A \rightarrow AC \rightarrow AC-B$  ring-construction strategy. The synthetic approach involved the preparation of a 2,3-dihydro-1H-quinolin-4-one system through an aza-Michael/sulfonamidation/Friedel-Crafts cyclization sequence. A  $6\pi$ -electrocyclization reaction was used to obtain the final B ring.



Eur. J. Org. Chem.  
DOI: 10.1002/ejoc.201501566

## Green chemistry



ChemistryOpen


DOI: 10.1002/open.201500159

## Green Synthesis

S. N. Maddila, S. Maddila, W. E. van Zyl, S. B. Jonnalagadda\*

Ceria–Vanadia/Silica-Catalyzed Cascade for C–C and C–O Bond Activation: Green One-Pot Synthesis of 2-Amino-3-cyano-4H-pyrans

**One-pot wonder:** We designed a ceria–vanadia/silica (Ce–V/SiO<sub>2</sub>) heterogeneous catalyst and used it for the green synthesis of 2-amino-3-cyano-4H-pyran derivatives, typically used in pharmaceuticals. The efficient reaction was a multicomponent one-pot condensation of 5,5-dimethylcyclohexane-1,3-dione, aromatic aldehyde, and malononitrile in ethanol, with yields up to 95%. The catalyst is economical, recyclable, and reusable for over five runs while preserving its high activity.



Asian J. Org. Chem.



DOI: 10.1002/ajoc.201500514

## Amide Synthesis

Y.-F. Guo, T.-L. Ren, B.-H. Xu,\* Y.-F. Wang, S.-J. Zhang\*

Cobalt-Catalyzed Dehydrogenative Coupling of Alcohols/Aldehydes and Amines: An Important Role for Imine Hydration

**A good hyd-ing:** A protocol for Co<sup>II</sup>/I<sup>−</sup>/tert-butyl hydroperoxide-catalyzed oxidative cross-coupling of alcohols or aldehydes with amines has been developed. tBuO•/tBuOO• rather than IO<sup>−</sup>/IO<sub>2</sub><sup>−</sup>/IO<sub>3</sub><sup>−</sup> is the active hydrogen-abstraction species in this catalytic process. Investigation into the reaction mechanism also revealed the potential transiency of a hemiaminal and the important role of its regeneration by imine hydration.



ChemNanoMat


DOI: 10.1002/cnma.201500181

## Biomaterialization

Y. Maeda, Z. Wei, Y. Ikezoe, E. Tam, H. Matsui\*

Biomimetic Crystallization of MnFe<sub>2</sub>O<sub>4</sub> Mediated by Peptide-Catalyzed Esterification at Low Temperature

**A catalytic peptide,** discovered through hydrogel-based phage display, could generate MnFe<sub>2</sub>O<sub>4</sub> nanocrystals exhibiting superparamagnetism at 4 °C in a methanol–benzyl alcohol mixture.



ChemViews magazine

DOI: 10.1002/chemv.201500106

## Biomass Production

F. Derwenskus, C. Holdmann

Microalgae – Underestimated All-Rounders

Microalgae have substantially higher biomass productivity than terrestrial plants. These diverse species have potential for a whole range of applications, including synthesizing food ingredients and recycling inorganic waste. The optimization of algae-based processes is a very promising area of research.

