

## Diazo Compounds

Deutsche Ausgabe: DOI: 10.1002/ange.201605337  
Internationale Ausgabe: DOI: 10.1002/anie.201605337Selective Synthesis of Six Products from a Single Indolyl  $\alpha$ -Diazocarbonyl Precursor

Michael J. James, Peter O'Brien, Richard J. K. Taylor,\* and William P. Unsworth\*

**Abstract:** Indolyl  $\alpha$ -diazocarbonyls can be selectively cyclized to give six distinct products through the careful choice of catalyst and reaction conditions. A range of catalysts were used, including complexes of  $Rh^{II}$ ,  $Pd^{II}$ , and  $Cu^{II}$ , as well as  $SiO_2$ , to promote diazo decomposition and subsequent cyclization/rearrangement through a range of mechanistic pathways.

The ability to access structurally diverse compounds is the cornerstone of lead generation in the pharmaceutical and agrochemical industries.<sup>[1]</sup> In most cases, such compounds are generated using organic synthesis, and over the years, a number of reliable and predictable methods have emerged.<sup>[1,2]</sup> The importance of such methods cannot be over-stated, but nonetheless, there is also value in the examination of reaction systems which react less predictably.<sup>[3]</sup> Reactive precursors known to participate in a wide range of synthetic transformations can significantly streamline the synthesis of diverse compounds by allowing multiple products to be generated from a single precursor, provided their reactivity can be controlled.

With this in mind, we initiated the research described herein, focusing on the reactions of indolyl  $\alpha$ -diazocarbonyl compounds.<sup>[4]</sup> The utility of diazo precursors in diversity-orientated synthesis was elegantly demonstrated by Warriner, Nelson and co-worker in 2014,<sup>[5]</sup> who exploited the unpredictable reactivity of  $\alpha$ -diazocarbonyls to generate product mixtures for bioassays. In our research we have taken an alternative approach, using a different reaction system, and focused on controlling the “unpredictable” nature of diazo-carbonyl reactivity by catalyst variation. The ability to access several distinct products from a common precursor is synthetically important, and such research can also lead to advances in the study of catalysis and mechanism. With this as motivation, we challenged ourselves to uncover a reaction system capable of delivering as many product scaffolds as possible from a single precursor by varying the catalyst and reaction conditions.<sup>[5]</sup> Most reported methods of this type allow the selective synthesis of two distinct products,<sup>[6]</sup> with protocols able to deliver three or more products being much more rare.<sup>[7]</sup> However, herein we report the catalyst-selective synthesis of six structurally distinct cyclic scaffolds from

a single  $\alpha$ -diazocarbonyl, of the form **1**, by a series of mild rhodium(II)-, palladium(II)-, copper(II)-, and  $SiO_2$ -catalyzed processes, as discovered through a mix of careful reaction design and serendipity (Figure 1).




Figure 1. Catalyst-selective synthesis: six scaffolds from one precursor.


Our studies began with the three-step synthesis of the  $\alpha$ -diazocarbonyl **1a** from the commercially available acid **2a** (Table 1),<sup>[8]</sup> which was then treated with a range of potential catalysts (10 mol%) in  $CH_2Cl_2$  at room temperature for 16 hours. Selected results are given in Table 1 (for full details, see the Supporting Information).

A number of catalysts able to promote diazo decomposition and cyclization were uncovered. Five identifiable products were observed in total, with mechanistically related products grouped to aid the subsequent discussion: the spirocyclic indolenine **3a** and  $\alpha,\beta$ -dicarbonyl **4a** (group A), C2 annulated indole **5a** and carbazole **6a** (group B), and isomeric indole **7a** (group C). As expected, many of the catalysts afforded complex mixtures of products, as exemplified by the reactions of the rhodium(II)- and copper(II)-based catalysts (Table 1, entries 1–4). However, more promising catalysts were also found and they enabled the selective synthesis of group A products **3a** and **4a** [ $[Rh_2O_4]$ ], group B redox isomers **5a** and **6a** [ $Pd(MeCN)_4(BF_4)_2$  or  $Cu(OTf)_2$ ], and the rearrangement product **7a** ( $SiO_2$ ), and these catalysts were therefore selected for further optimization.

To the best of our knowledge, the  $[Rh_2O_4]$ -catalyzed procedure<sup>[9]</sup> to form **3a** represents the first reported synthesis of a spirocyclic indolenine<sup>[10]</sup> from a diazocarbonyl precursor, although C3 functionalization of indoles using diazocarbonyl compounds has been reported,<sup>[11,12]</sup> so this outcome was not wholly unexpected.<sup>[13]</sup> However, the formation of the oxidized

[\*] M. J. James, Prof. P. O'Brien, Prof. R. J. K. Taylor, Dr. W. P. Unsworth  
Department of Chemistry, University of York  
York, YO10 5DD (UK)  
E-mail: richard.taylor@york.ac.uk  
william.unsworth@york.ac.uk


Supporting information and the ORCID identification number(s) for the author(s) of this article can be found under <http://dx.doi.org/10.1002/anie.201605337>.

**Table 1:** Initial catalyst screening.


Entry	Catalyst <sup>[a]</sup>	Product composition [%] <sup>[b]</sup>					
		1a	3a	4a	5a	6a	7a
1	mix	[Rh <sub>2</sub> (OAc) <sub>4</sub> ]	–	–	50	20	15
2		[Rh <sub>2</sub> esp <sub>2</sub> ]	–	10	55	15	–
3		Cu(MeCN) <sub>4</sub> OTf	–	20	–	60	20
4		Cu(MeCN) <sub>4</sub> PF <sub>6</sub>	15	65	20	–	–
5	A	[Rh <sub>2</sub> oct <sub>4</sub> ]	–	95	5	–	–
6	B	Pd(MeCN) <sub>4</sub> (BF <sub>4</sub> ) <sub>2</sub>	–	–	–	95	5
7		Cu(OTf) <sub>2</sub>	–	–	–	70	30
8	C	SiO <sub>2</sub> (1 g/g)	–	–	–	–	> 90

[a] Reactions performed with 0.05 mmol of **2a** and 10 mol % catalyst in CH<sub>2</sub>Cl<sub>2</sub> (0.1 M) under argon at RT for 16 h. [b] Calculated using the <sup>1</sup>H NMR spectrum of the unpurified reaction mixture. esp =  $\alpha,\alpha,\alpha',\alpha'$ -tetramethyl-1,3-benzenedipropionate, oct = octanoate, *p*-ABSA = *p*-acetamidobenzenesulfonyl azide, T3P = propane phosphonic acid anhydride, Tf = trifluoromethanesulfonyl.

product **4a** was much more surprising, with the structure of this product confirmed by X-ray crystallography.<sup>[14]</sup> It was found that the selective synthesis of either product could be achieved, with the reaction outcome being dependent on the presence of air in the reaction. By switching the reaction solvent from CH<sub>2</sub>Cl<sub>2</sub> to chloroform, reducing catalyst loading to 5 mol %, and performing the reaction under oxygen-free conditions, **3a** was isolated in 92 % yield (Scheme 1). Furthermore, carrying out the same reaction in a flask open to air was sufficient to completely switch the selectivity to efficiently furnish **4a**.


**Scheme 1.** Selective synthesis of **3a** and **4a**.

We propose that both reactions start with the formation of the rhodium carbenoid **A**,<sup>[15]</sup> which then reacts with the nucleophilic indole to form the spirocycle **B**, before undergoing protodemetalation to furnish **3a** (Scheme 2). Then, in

**Scheme 2.** Proposed mechanism for the formation of **3a** and **4a**.


the presence of oxygen, we propose that **3a** forms the intermediate endoperoxide **D**,<sup>[8a]</sup> possibly by a radical rebound process (**3a** → **C** → **D**),<sup>[16]</sup> before fragmenting as shown to afford the product **4a**.<sup>[17]</sup> Additional evidence for this mechanism (including an X-ray structure for a related endoperoxide) can be found in the Supporting Information.

While **4a** could be isolated in good yield, it was found to be relatively short-lived as it degraded during silica gel chromatography and upon storage, but pleasingly, we were able to exploit its high reactivity to deliver two new oxindole scaffolds, **8a** and **9a** (Scheme 3). Thus, two highly diastereoselective intramolecular aldol-type reactions were developed using either Brønsted-acidic or Brønsted-basic conditions.

**Scheme 3.** Stereoselective formation of **8a** and **9a**.

Both reactions were performed in one pot, thus requiring only a solvent switch to THF and the addition of an excess of either TFA or *t*BuOK. Under acidic conditions **4a** was selectively converted into the *syn*-diastereoisomer **8a** in 99 % yield, and we propose this to be the result of hydrogen bonding between the oxindole and  $\alpha,\beta$ -dicarbonyl moieties (**E**). Conversely, under basic conditions, the *anti*-diastereoisomer **9a** was formed, and we propose that it results from a reactive conformation of the form **F**, in which the destabilizing steric interactions are lower than those in **E**, and the carbonyl dipoles are opposed. Both product structures were confirmed by X-ray analysis.<sup>[14]</sup>


Next, the palladium(II)- and copper(II)-catalyzed reactions were optimized, thus allowing selective formation of the C2-annulated product **5a** and carbazole **6a** using Pd(MeCN)<sub>4</sub>(BF<sub>4</sub>)<sub>2</sub> (5 mol %) and Cu(OTf)<sub>2</sub> (20 mol %), respectively (Scheme 4).<sup>[18,19]</sup> A key difference between these reactions is



**Scheme 4.** Selective formation of **5a** and **6a**. TFA = trifluoroacetic acid, THF = tetrahydrofuran.

that it is necessary to perform the carbazole-forming reaction under oxygen at 50 °C. It is difficult to unambiguously determine whether these reactions proceed by direct nucleophilic attack from the indole C2 or by initial C3 attack followed by a 1,2-migration. Based on related precedent,<sup>[20]</sup> and the observation that **3a** can be converted into a mixture of **5a** and **6a** upon reaction with Cu(OTf)<sub>2</sub>, the latter appears more likely.

The silica-promoted C2-annulation reaction required minimal deviation from the initial screen. The compound **7a** was prepared in good yield by reacting **1a** with an equivalent weight of SiO<sub>2</sub> in CH<sub>2</sub>Cl<sub>2</sub> (Scheme 5). The reaction likely



**Scheme 5.** Synthesis of **7a**.

proceeds by a Wolff rearrangement, induced by the mildly acidic silica,<sup>[21]</sup> and trapping by the nucleophilic indole (either by direct C2 attack or by an initial C3 attack followed by a 1,2-migration). To the best of our knowledge, only one other example of a C2-annulation reaction of this type has been reported.<sup>[22]</sup>

Finally, the scope of all six procedures was tested on five diazocarbonyl substrates (**1a–e**), thus delivering 30 discrete products in total (Scheme 6). The spirocycles **3a–e** were each formed in good yield, with variable diastereomeric ratios, and is likely due to epimerization of the  $\alpha$ -keto stereocenter during chromatography. The other five procedures were all well tolerated by the same precursor set. The spirocyclic oxindoles **8a–e** and **9a–e**, as well C2 annulation products **5a–e**, **6a–e**, and **7a–e** were formed in generally good yields, and is pleasing given that no additional optimization was performed for any of these reactions.<sup>[23]</sup>

In summary, we report a novel catalyst-controlled approach to form six structurally diverse products from a single  $\alpha$ -diazocarbonyl precursor. While other catalyst-selective synthesis systems are known,<sup>[5–8]</sup> we know of no other capable of delivering the level of scaffold diversity by simply varying the catalyst and reaction conditions. Given the importance and diversity of the compound classes accessible, the methods are expected to be of much synthetic interest,<sup>[24,25]</sup> while the novel reactivity and mechanistic information uncovered is likely to be useful to researchers studying catalysis. These discoveries (some of which were serendipitous) were made as a result of challenging the methodology in terms of the number of products which could be selectively formed. Much as natural product synthesis has long been used to inspire the invention of new synthetic processes,<sup>[26]</sup> we believe that the same principles apply in catalyst-selective synthesis.

## Acknowledgments


We wish to thank the University of York (M.J.J. and W.P.U.) and the Leverhulme Trust (for an Early Career Fellowship, ECF-2015-013, W.P.U.) for financial support and Dr. A. C. Whitwood for X-ray crystallography.

**Keywords:** cyclizations · diazo compounds · indoles · spirocycles · synthetic methods

**How to cite:** *Angew. Chem. Int. Ed.* **2016**, *55*, 9671–9675  
*Angew. Chem.* **2016**, *128*, 9823–9827

- [1] S. D. Roughley, A. M. Jordan, *J. Med. Chem.* **2011**, *54*, 3451.
- [2] T. W. J. Cooper, I. B. Campbell, S. J. F. Macdonald, *Angew. Chem. Int. Ed.* **2010**, *49*, 8082; *Angew. Chem.* **2010**, *122*, 8258.
- [3] G. Karageorgis, S. Warriner, A. Nelson, *Nat. Chem.* **2014**, *6*, 872.
- [4] a) H. M. L. Davies, S. J. Hedley, *Chem. Soc. Rev.* **2007**, *36*, 1109; b) A. Ford, H. Miel, A. Ring, C. N. Slattery, A. R. Maguire, M. A. McKervy, *Chem. Rev.* **2015**, *115*, 9981.
- [5] J. Mahatthananchai, A. M. Dumas, J. W. Bode, *Angew. Chem. Int. Ed.* **2012**, *51*, 10954; *Angew. Chem.* **2012**, *124*, 11114.
- [6] See Ref. [5], and: a) J. D. Dooley, S. Reddy Chidipudi, H. W. Lam, *J. Am. Chem. Soc.* **2013**, *135*, 10829; b) M. J. James, R. E. Clubley, K. Y. Palate, T. J. Procter, A. C. Wyton, P. O'Brien, R. J. K. Taylor, W. P. Unsworth, *Org. Lett.* **2015**, *17*, 4372; c) Q.-Q. Cheng, J. Yedoyan, H. Arman, M. P. Doyle, *J. Am. Chem. Soc.* **2016**, *138*, 44.
- [7] For exceptions, see Ref. [5], and: a) G. Zhan, M.-L. Shi, Q. He, W.-J. Lin, Q. Ouyang, W. Du, Y.-C. Chen, *Angew. Chem. Int. Ed.* **2016**, *55*, 2147; *Angew. Chem.* **2016**, *128*, 2187; b) J. T. R. Liddon,





**Scheme 6.** Catalyst-selective synthesis of six products from a single indolyl  $\alpha$ -diazocarbonyl precursor (yields are those of products isolated after column chromatography).

- M. J. James, A. K. Clarke, P. O'Brien, R. J. K. Taylor, W. P. Unsworth, *Chem. Eur. J.* **2016**, 22, 8777.
- [8] a) H. Yang, J. Feng, Y. Tang, *Chem. Commun.* **2013**, 49, 6442; b) S. Muthusamy, M. Sivaguru, *Org. Lett.* **2014**, 16, 4248.
- [9] We are unable to account for the difference in product ratio of the three rhodium(II)-catalyzed reactions in Table 1 (entries 1, 2, and 5). Subtle differences in the solubility, coordination chemistry, and redox potential of the catalysts/intermediates can all influence the reaction outcome. It is noteworthy that all of these reactions had a distinct dark red/burgundy color, which is often indicative of Rh<sup>II</sup> dimer formation.
- [10] For a recent review, see: M. J. James, P. O'Brien, R. J. K. Taylor, W. P. Unsworth, *Chem. Eur. J.* **2016**, 22, 2856.
- [11] Using rhodium(II) catalysts: a) Y. Lian, H. M. L. Davies, *J. Am. Chem. Soc.* **2010**, 132, 440; b) T. Goto, Y. Natori, K. Takeda, H. Nambu, S. Hashimoto, *Tetrahedron: Asymmetry* **2011**, 22, 907; c) A. DeAngelis, V. W. Shurtleff, O. Dmitrenko, J. M. Fox, *J. Am. Chem. Soc.* **2011**, 133, 1650.
- [12] Using other metals: a) M. Delgado-Rebollo, A. Prieto, P. J. Pérez, *ChemCatChem* **2014**, 6, 2047; b) X. Gao, B. Wu, W.-X. Huang, M.-W. Chen, Y.-G. Zhou, *Angew. Chem. Int. Ed.* **2015**, 54, 11956; *Angew. Chem.* **2015**, 127, 12124; c) M. Li, X. Guo, W. Jin, Q. Zheng, S. Liu, W. Hu, *Chem. Commun.* **2016**, 52, 2736.
- [13] For related dearomatization reactions, see: a) C.-X. Zhuo, C. Zheng, S.-L. You, *Acc. Chem. Res.* **2014**, 47, 2558; b) M. J. James, J. D. Cuthbertson, P. O'Brien, R. J. K. Taylor, W. P. Unsworth, *Angew. Chem. Int. Ed.* **2015**, 54, 7640; *Angew. Chem.* **2015**, 127, 7750; c) S. J. Chambers, G. Coulthard, W. P. Unsworth, P. O'Brien, R. J. K. Taylor, *Chem. Eur. J.* **2016**, 22, 6496.
- [14] CCDC 1481924 (**4a**), 1481925 (**5a**), 1481926 (**6a**), 1481927 (**8a**), 1481928 (**9a**), and 1481930 (**7a**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre.

- [15] a) H. M. L. Davies, R. E. J. Beckwith, *Chem. Rev.* **2003**, *103*, 2861; b) H. M. L. Davies, D. Morton, *Chem. Soc. Rev.* **2011**, *40*, 1857.
- [16] a) B. Ensing, F. Buda, M. C. M. Gribnau, E. J. Baerends, *J. Am. Chem. Soc.* **2004**, *126*, 4355; b) R. Curci, L. D'Accolti, C. Fusco, *Acc. Chem. Res.* **2006**, *39*, 1; c) C. Yuan, Y. Liang, T. Hernandez, A. Berriochoa, K. N. Houk, D. Siegel, *Nature* **2013**, *499*, 192.
- [17] Purified **3a** slowly converts into **4a** when stored in solution open to air, thus confirming that the rhodium catalyst is not essential for oxidation.
- [18] Two other C2 annulation reactions of indole derivatives by reaction with a diazocarbonyls have been reported: a) E. Cuevas-Yañez, J. M. Muchowski, R. Cruz-Almanza, *Tetrahedron* **2004**, *60*, 1505; b) C. S. Shanahan, P. Truong, S. M. Mason, J. S. Leszczynski, M. P. Doyle, *Org. Lett.* **2013**, *15*, 3642.
- [19] For related carbazole syntheses, see: K. S. Rathore, B. S. Lad, H. Chennamsetti, S. Katukojvala, *Chem. Commun.* **2016**, *52*, 5812.
- [20] a) Q.-F. Wu, C. Zheng, S.-L. You, *Angew. Chem. Int. Ed.* **2012**, *51*, 1680; *Angew. Chem.* **2012**, *124*, 1712; b) V. A. Peshkov, O. P. Pereshivko, E. V. Van der Eycken, *Adv. Synth. Catal.* **2012**, *354*, 2841; c) C. Zheng, Q.-F. Wu, S.-L. You, *J. Org. Chem.* **2013**, *78*, 4357.
- [21] No reaction took place when Et<sub>3</sub>N was included as an additive.
- [22] M. Salim, A. Capretta, *Tetrahedron* **2000**, *56*, 8063.
- [23] For all entries a single product was isolated and we observed no appreciable amounts of any of the other products in these reactions. In the lower-yielding examples, the mass balance was largely made up of intractable, polar material, which we were unable to identify.
- [24] For other approaches to scaffold diversity, see: a) H. Mizoguchi, H. Oikawa, H. Oguri, *Nat. Chem.* **2014**, *6*, 57; b) W. Liu, V. Khedkar, B. Baskar, M. Schürmann, K. Kumar, *Angew. Chem. Int. Ed.* **2011**, *50*, 6900; *Angew. Chem.* **2011**, *123*, 7032; c) D. J. Foley, R. G. Doveston, I. Churcher, A. Nelson, S. P. Marsden, *Chem. Commun.* **2015**, *51*, 11174.
- [25] Efforts to extend these methods to other heteroarenes are ongoing.
- [26] a) P. S. Baran, T. J. Maimone, J. M. Richter, *Nature* **2007**, *446*, 404; b) K. C. Nicolaou, C. R. H. Hale, *Natl. Sci. Rev.* **2014**, *1*, 233.

Received: June 3, 2016

Published online: July 20, 2016