




Nickel-Catalyzed Cross-Coupling of Redox-Active Esters with Boronic Acids

Jie Wang, Tian Qin, Tie-Gen Chen, Laurin Wimmer, Jacob T. Edwards, Josep Cornella, Benjamin Vokits, Scott A. Shaw, and Phil S. Baran*


Abstract: A transformation analogous in simplicity and functional group tolerance to the venerable Suzuki cross-coupling between alkyl-carboxylic acids and boronic acids is described. This Ni-catalyzed reaction relies upon the activation of alkyl carboxylic acids as their redox-active ester derivatives, specifically *N*-hydroxy-tetrachlorophthalimide (TCNHPI), and proceeds in a practical and scalable fashion. The inexpensive nature of the reaction components ($\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$ —\$9.5 mol^{-1} , Et_3N) coupled to the virtually unlimited commercial catalog of available starting materials bodes well for its rapid adoption.

Alkyl carboxylic acids and boronic acids are amongst the most widely available building blocks for modern organic synthesis (Figure 1A). Recent surveys and informatic approaches to mining chemical reaction space within drug discovery explain this phenomenon: Amide bond formation and Suzuki couplings are the most often employed reactions.^[1,2] A reaction that could therefore marry these ubiquitous and convenient building blocks in one diversifying step would likely be of great utility in all areas of chemical science. Earlier this year we reported that common activated esters (such as HOAt, HOBt, and NHPI) normally employed in amide-bond formation could also engage in Negishi-type couplings with aryl- and alkylzinc reagents under nickel-based catalysis.^[3,4] The fact that all of these activated esters not only prime the carboxyl for nucleophilic attack but also accept an electron from a low-valent metal in an SET-based thermal process was previously unrecognized.^[5,6] Whereas the former process allows for ester and amide formation, the latter enables a simple thermal decarboxylative radical formation with immediate capture by that transition metal (Ni).^[3–5,7] Herein, a reaction analogous to the venerable Suzuki coupling for the cross-coupling of activated alkyl carboxylic acids with aryl and vinyl boronic acids is reported exhibiting a broad substrate scope, scalability, functional group tolerance, and inherent accessibility.^[8]

A. New coupling: boronic acids and alkyl acids as "off the shelf" coupling partners.



B. Invention: Ni-catalyzed cross-coupling of redox active esters with arylboronic acids^e



entry	deviation from above	yield (%) ^b	entry	deviation from above	yield (%) ^b
1	1,4-dioxane	20 ^c	8	L2	64
2	THF:DMF (10:1)	10	9	L3	58
3	DIPEA	68	10	L4	82
4	Bu_3N	43	11	NHPI ester	<5
5	K_2PO_4	<5	12	w/o L1	<5
6	Cs_2CO_3	<5	13	w/o $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$	<5
7	40 mol% L1	44	14	w/o Et_3N	40

^a 0.1 mmol. ^b Yield determined by GC using dodecane as internal standard. ^c Isolated yield.

Figure 1. A) Boronic acids and carboxylic acids as desired coupling partners in a new cross-coupling reaction. B) Optimization of a Ni-catalyzed cross-coupling between boronic acids and redox-active esters.

Although conceptually simple, the execution of this coupling relied on extensive experimentation and some empirical observations that are at this time poorly understood.

Figure 1B presents an abbreviated picture of the optimization conducted on ester **1** and boronic acid **2**, culminating in 78% isolated yield of **3**. Solvent and concentration played a crucial role in this chemistry, with a mixture of 1,4-dioxane and *N,N*-dimethylformamide (DMF, 0.023 M) emerging as the optimal.^[9] For example, exclusion of DMF diminished the yield (entry 1) and when 1,4-dioxane was substituted for THF this seemingly trivial change essentially shut down the reaction (entry 2). The choice of base was also unusually important with excess triethylamine furnishing the best results (entries 3–6).

Next, numerous ligands were screened and it was found that a 1:1 ratio of Ni/L performed the best with **L1** and **L4**


[*] Dr. J. Wang, Dr. T. Qin, Dr. T.-G. Chen, Dr. L. Wimmer, J. T. Edwards, Dr. J. Cornella, Prof. P. S. Baran
The Scripps Research Institute (TSRI)
North Torrey Pines Road, La Jolla, CA 92037 (USA)
E-mail: pbaran@scripps.edu

B. Vokits, Dr. S. A. Shaw
Discovery Chemistry, Bristol-Myers Squibb
350 Carter Road, Hopewell, NJ 08540 (USA)

Supporting information and the ORCID identification number(s) for the author(s) of this article can be found under <http://dx.doi.org/10.1002/anie.201605463>.

providing the most reliable outcomes (entries 7–10). The choice of ester was also critical as the more electron-rich redox-active NHPI derivative was an incompetent coupling partner under these conditions (entry 11). TCNHPI, an activating agent used for the first time in synthesis as a catalyst for hydrogen atom transfer,^[10] proved to be ideal for this coupling (Commercially available from Sigma-Aldrich, catalog # ALD00564). Control experiments pointed to the essential roles of the catalyst and ligand (entries 12–13). Curiously, even in the absence of base the reaction proceeds to some extent, possibly due to the basic nature of the ligands employed (entry 14). It is worth mentioning that a preformed Ni-ligand complex in DMF added to a 1,4-dioxane solution of boronic acid, base, and activated ester gave the most robust results.^[9]


Over thirty examples of this cross-coupling are depicted in Scheme 1. The reaction appears to be general for both coupling partners and largely undeterred by a wide variety of functional groups such as aryl halides (**3–13**, **15**, **17**, **19**, **20**, **23**, **26**, **28**, **35**, **41**), common protecting groups (**4**, **11**, **13–16**, **21–38**), esters (**12**, **15**, **16**), nitriles (**25**, **39**), ketones (**17**, **22**), carbamates (**15–16**), and even medicinally relevant heterocycles (**14**, **16**, **18**, **28–34**). Both primary and secondary alkyl carboxylic acids with or without a radical stabilizing groups are viable (**11**, **18–19**) and *N*-Boc protected amino acid side chains show no erosion of optical activity (**15–16**). Both electron-rich (**24**, **27**, **34**, **40**) and electron-poor (**22–23**, **25–26**, **28–33**, **39**) aryl boronic acids can be employed as well as vinyl boronic acids (**35–36**).



An important feature of this new cross-coupling hinges on the simplicity of its setup and the development of a new one-pot protocol. Hence, the transformation could be conducted without the need for isolation of the activated esters, as exemplified by 10 representative substrates in Scheme 1 (**3–5**, **8**, **21–25**, **29**). A simple activation of the carboxylic acid by DCC, followed by the Suzuki protocol, afforded comparable yields as starting from the isolated redox-active ester. This rapid access coupled with the experimental ease of the union might have a tangible impact on the speed with which many organic compounds can be made. For example, the preparation of *p*-bromophenyl substituted species such as **5** has previously required as many as five steps to procure.^[11] Most routes to compounds such as **3–10** involve organometallic additions to ketones followed by cationic or radical-based reductions.^[12] The simple ester **12** has previously required more than five steps to prepare.^[13] The unnatural amino acids **15** and **16** have required multistep pathways involving homologation of aspartic acid rather than a direct coupling with glutamic acid as in the present work.^[14] A direct comparison to reported substrates for Ni-catalyzed protocols using alkyl bromides and arylboronic species resulted similar yields of the cross-coupling product when the redox-active ester is utilized (**39–40**).^[8a,b] Albeit the apparent similarities of both protocols, we hypothesized that a different mechanism could be operating under our optimized conditions. To test the hypothesis, an alkyl bromide was embedded in a pending chain of a linear redox-active ester and subjected to the optimized conditions. Interestingly, complete chemoselectivity favoring coupling of the redox-active ester was obtained for substrate **41**, which highlights the orthogonal compatibility with other protocols in the area of Suzuki arylations. It is worth mentioning that the catalyst loading was standardized to 20 mol% across the examples in Scheme 1 for optimal yields. However, we demonstrated that comparable yields could also be obtained when the catalyst loading was lowered to 10% (**4**, **21–23**, **29**).

The experimental ease of this transformation cannot be overemphasized as the demanding environment of medicinal chemistry prioritizes those reactions that require minimal experimental precaution. Thus, as depicted in Scheme 2A, a reaction of **1** was conducted using wet solvents without any precaution to exclude air or moisture (open-flask) to produce adduct **3** in 65% isolated yield. The scalability was also evaluated using **42** and it was found to proceed smoothly to deliver **3** on a gram-scale (Scheme 2B). In accord with our previous findings,^[3,4] the reaction of redox-active esters with in situ generated low valent Ni species generates radical intermediates as evidenced by the cyclopropane-opened product **44** derived from **43** (Scheme 2C).

A postulated mechanism for this transformation is depicted in Scheme 2D. By analogy to prior mechanistic investigations of Ni-catalyzed cross-coupling reactions with alkyl halides,^[7] and our previous studies using organozinc reagents,^[3,4] a related pathway is postulated for the coupling of redox-active esters with boronic acids. Initially, a Ni^I complex I undergoes transmetalation with an arylboronic acid aided by Et₃N and H₂O delivering aryl-Ni^I complex II.^[15] Reduction of the redox-ester by complex II affords species III



(TCNHPI). Numerous examples have been presented to demonstrate the broad scope of this transformation and it has already found use in the context of drug development.^[17] It is likely that many other branches of chemistry will benefit from this cross-coupling as well. Studies to expand the scope of coupling to alkyl boronic acids, develop asymmetric variants, and deeply probe the mechanism of this fascinating transformation are underway and will be reported in due course.

Acknowledgements

Financial support for this work was provided by NIH (grant number GM-097444), Bristol-Myers Squibb, the Shanghai Institute of Organic Chemistry, Zhejiang Medicine Co., and Pharmaron (postdoctoral fellowship for J.W.), Austrian Science Fund (FWF, Schrödinger Fellowship to L. W.), Department of Defense (NDSEG fellowship to J. T. E.) and Catalan Government (Beatriu de Pinos Postdoctoral Fellowship to J.C.). We thank David E. Hill for help in HPLC measurements. We also thank Ke Chen, Michael Schmidt, and Martin D. Eastgate for helpful discussions. We are grateful to D.-H. Huang and L. Pasternack (The Scripps Research Institute) for assistance with nuclear magnetic resonance (NMR) spectroscopy. We also thank A. L. Rheingold, C. E. Moore, and M. A. Galella for X-ray crystallographic analysis.^[18]

Keywords: decarboxylation · homogeneous catalysis · nickel catalysts · redox-active esters · Suzuki cross-coupling

How to cite: *Angew. Chem. Int. Ed.* **2016**, 55, 9676–9679
Angew. Chem. **2016**, 128, 9828–9831

- [1] N. Schneider, D. M. Lowe, R. A. Sayle, M. A. Tarselli, G. A. Landrum, *J. Med. Chem.* **2016**, 59, 4385.
- [2] D. G. Brown, J. Boström, *J. Med. Chem.* **2016**, 59, 4443.
- [3] J. Cornella, J. T. Edwards, S. Kawamura, J. Wang, C.-M. Pan, R. Gianatassio, M. Schmidt, M. D. Eastgate, P. S. Baran, *J. Am. Chem. Soc.* **2016**, 138, 2174.
- [4] T. Qin, J. Cornella, C. Li, L. Malins, J. T. Edwards, S. Kawamura, B. D. Maxwell, M. D. Eastgate, P. S. Baran, *Science* **2016**, 352, 801.
- [5] Shortly after ref. [3] was published, D. J. Weix reported that NHPI esters could undergo reductive cross-coupling with aryl iodides using Zn^0 as the reducing agent: K. M. M. Huihui, J. A. Caputo, Z. Melchor, A. M. Olivares, A. M. Spiewak, K. A. Johnson, T. A. DiBenedetto, S. Kim, L. K. G. Ackerman, D. J. Weix, *J. Am. Chem. Soc.* **2016**, 138, 5016.
- [6] For the use of NHPI esters in photochemically initiated SET processes, see: a) M. J. Schnermann, L. E. Overman, *Angew. Chem. Int. Ed.* **2012**, 51, 9576; *Angew. Chem.* **2012**, 124, 9714; b) G. Pratsch, G. L. Lackner, L. E. Overman, *J. Org. Chem.* **2015**, 80, 6025; c) K. Okada, K. Okamoto, M. Oda, *J. Am. Chem. Soc.* **1988**, 110, 8736; d) K. Okada, K. Okamoto, N. Morita, K. Okubo, M. Oda, *J. Am. Chem. Soc.* **1991**, 113, 9401.
- [7] As mentioned in ref. [3], redox-active esters are analogous to alkyl halides. Thus, a mechanistic parallel can be directly drawn to pioneering work in the area of Ni-catalyzed single-electron transfer cross-coupling of those species, see: a) T. J. Anderson, G. D. Jones, D. A. Vicić, *J. Am. Chem. Soc.* **2004**, 126, 8100; b) G. D. Jones, J. L. Martin, C. McFarland, O. R. Allen, R. E. Hall, A. D. Haley, R. J. Brandon, T. Konovalova, P. J. Desrochers, P. Pulay, D. A. Vicić, *J. Am. Chem. Soc.* **2006**, 128, 13175; c) N. D. Schley, G. C. Fu, *J. Am. Chem. Soc.* **2014**, 136, 16588; d) V. B. Phapale, E. Buñuel, M. García-Iglesias, D. J. Cárdenas, *Angew. Chem. Int. Ed.* **2007**, 46, 8790–8795; *Angew. Chem.* **2007**, 119, 8946–8951; e) O. Gutierrez, J. C. Tellis, D. N. Primer, G. A. Molander, *J. Am. Chem. Soc.* **2015**, 137, 4896; f) A. N. Vedernikov, *ChemCatChem* **2014**, 6, 2490; g) B. Zheng, F. Tang, J. Luo, J. W. Schultz, N. P. Rath, L. M. Mirica, *J. Am. Chem. Soc.* **2014**, 136, 6499; h) J. Breitenfeld, J. Ruiz, M. D. Wodrich, X. Hu, *J. Am. Chem. Soc.* **2013**, 135, 12004; i) S. Biswas, D. J. Weix, *J. Am. Chem. Soc.* **2013**, 135, 16192.
- [8] For selected examples of Ni-catalyzed alkyl–aryl Suzuki–Miyaura cross-couplings, see: a) J. Zhou, G. C. Fu, *J. Am. Chem. Soc.* **2004**, 126, 1340; b) F. González-Bobes, G. C. Fu, *J. Am. Chem. Soc.* **2006**, 128, 5360; c) P. Lundin, G. C. Fu, *J. Am. Chem. Soc.* **2010**, 132, 11027; d) A. Wilsily, F. Tramutola, N. A. Owston, G. C. Fu, *J. Am. Chem. Soc.* **2012**, 134, 5794; e) S. Zultanski, G. C. Fu, *J. Am. Chem. Soc.* **2013**, 135, 624; f) R. Jana, T. P. Pathak, M. S. Sigman, *Chem. Rev.* **2011**, 111, 1417.
- [9] See Supporting Information for details.
- [10] E. J. Horn, B. R. Rosen, Y. Chen, J. Tang, K. Chen, M. D. Eastgate, P. S. Baran, *Nature* **2016**, 533, 77.
- [11] a) K. Ueda, H. Umihara, S. Yokoshima, T. Fukuyama, *Org. Lett.* **2015**, 17, 3191; For an older route involving Friedel–Crafts of benzene with cyclohexene followed by bromination, see: b) B. B. Corson, V. N. Ipatieff, *Org. Synth.* **1939**, 19, 36.
- [12] a) I. Papanastasiou, S. Riganas, G. B. Foscolos, A. Tsotinis, S. A. Akhtar, M. A. Khan, K. M. Rahman, D. E. Thurston, *Lett. Org. Chem.* **2015**, 12, 319; b) D. M. Allwood, D. C. Blakemore, A. D. Brown, S. V. Ley, *J. Org. Chem.* **2014**, 79, 328.
- [13] M. Zaidlewicz, A. Wolan, *J. Organomet. Chem.* **2002**, 657, 129.
- [14] R. F. W. Jackson, J. L. Frase, N. Wishart, B. Porter, M. J. Wythes, *J. Chem. Soc. Perkin Trans. 1* **1998**, 1903.
- [15] Recently, the formation of hydroxy-bridged species between a boronate and the metal center have disclosed as key step for transmetalation in Suzuki cross-couplings with Pd: A. A. Thomas, S. E. Denmark, *Science* **2016**, 352, 329.
- [16] The capture of similar radicals was also invoked during the decarboxylative coupling of α -heteroatom stabilized acids with aryl halides: Z. Zuo, D. T. Ahneman, L. Chu, J. A. Terrett, A. G. Doyle, D. W. C. MacMillan, *Science* **2014**, 345, 437.
- [17] This reaction is currently being enlisted at Bristol-Myers Squibb in several ongoing programs, including one involving the co-authors (B.V. and S.A.S.).
- [18] X-ray information for compound **17** can be obtained free of charge from The Cambridge Crystallographic Data center with number CCDC 1477360.

Received: June 5, 2016

Published online: July 6, 2016