



Auf diesen Seiten weisen wir auf wichtige aktuelle Beiträge in unseren Schwesterzeitschriften hin. Wenn Sie die Seiten online lesen, dann können Sie

die Artikel mit einem Klick direkt aufrufen, ansonsten sind sie durch Eingabe der DOIs über Wiley Online Library leicht online zugänglich.




Ionic Liquids

Y. Mieno, Y. Kohno, S. Saita, H. Ohno*

Zwitterion/Brønsted Acid Mixtures Showing Controlled Lower Critical Solution Temperature-Type Phase Changes with Water

Just a phase! Ammonium cation-based zwitterions with adequately designed hydrophobicity were mixed with aqueous Brønsted acid solutions and were shown to reversibly undergo thermoresponsive phase changes between a homogeneous mixture and liquid–liquid phase separation with water by a lower critical solution temperature-type phase transition (see figure).



Chem. Eur. J.
DOI: 10.1002/chem.201600973




Substituent Effects

G. He, C. Yu, Y. Li, J. Hu, Z. Liu, D. Zhang, Q. Guo,* A. Xia*

Excitation Energy Transfer in *meta*-Substituted Phenylacetylene Multibranch Chromophores

Branch effects: A comprehensive investigation into the spectral properties and excitation energy transfer in di- or tribranched dithienylidiotopyrrolopyrrole molecules with *meta*-substituted benzene as a central core, by means of steady-state and transient measurements and quantum chemical calculations, is reported (see figure).



Chem. Asian J.
DOI: 10.1002/asia.201600326




Fluorescent Probes


T. Oura, K. Murata, T. Morita, A. Nezu, M. Arisawa, S. Shuto,* A. Tanimura*

Highly Sensitive Measurement of Inositol 1,4,5-Trisphosphate by Using a New Fluorescent Ligand and Ligand Binding Domain Combination

The binding of fluorescent ligand (FL) to a sensor consisting of fluorescent protein (CFP) and the ligand binding domain of the IP₃R (LBD) causes FRET. Excitation of CFP leads to transfer of the excitation energy to FL. IP₃ competes with FL for binding to the LBD, thereby resulting in a decrease in the FRET signal.



ChemBioChem
DOI: 10.1002/cbic.201600096



ChemPhysChem



DOI: 10.1002/cphc.201600174

Rotational Motion

B. Neupane, F. Chen, Y. Wei, N. Fang,* F. S. Ligler,* G. Wang*

Nanosecond Time-Resolution Study of Gold Nanorod Rotation at the Liquid–Solid Interface

In a spin: A nanosecond time-resolution study yields new insights into the rotational motion of anchored rod-like nanoparticles at the liquid–solid interface. The continuous rotation observed under low temporal resolution is found to be comprised of numerous fast, intermittent transitions between a limited number of weakly immobilized states.



ChemMedChem



DOI: 10.1002/cmdc.201500383

Multitarget Drugs

O. Benek, O. Soukup, M. Pasdiorova, L. Hroch, V. Sepsova, P. Jost, M. Hrabanova, D. Jun, K. Kuca, D. Zala, R. R. Ramsay, J. Marco-Contelles,* K. Musilek*

Design, Synthesis and in vitro Evaluation of Indolotacrine Analogues as Multitarget-Directed Ligands for the Treatment of Alzheimer's Disease

MAO meets ChE! By using a multitarget-directed ligand approach, a series of novel compounds were designed to act simultaneously as cholinesterase (ChE) and monoamine oxidase (MAO) inhibitors. The most promising compound, indolotacrine **9b**, was found to be a potent inhibitor of both cholinesterases and MAO A and a moderately potent inhibitor of MAO B.



ChemSusChem



DOI: 10.1002/cssc.201600462

Water Splitting

L. Wang, N. T. Nguyen, P. Schmuki*

A Facile Surface Passivation of Hematite Photoanodes with Iron Titanate Cocatalyst for Enhanced Water Splitting

Shields up! A thin Fe₂TiO₅ layer was decorated on nanostructured hematite photoanodes by a facile water-based solution method. The Fe₂O₃/Fe₂TiO₅ heterostructure exhibits an obvious enhancement in solar water splitting. The enhancement in photocurrent can be attributed to the effect of Fe₂TiO₅ overlayers passivating surface states and thus reducing surface electron–hole recombination.



ChemCatChem


DOI: 10.1002/cctc.201600330


Hydrogenation

F. J. L. Heutz, C. Erken, M. J. B. Aguilá, L. Lefort, P. C. J. Kamer*

Heterogeneous Hydrogenation of Esters under Mild Conditions using Solid-Supported Phosphorus–Ruthenium Catalysts

Jekyll and Hyde: the first heterogeneous catalyst capable of hydrogenating esters under very mild, typically homogeneous, conditions (25 °C, 5.0 MPa) is described. Additionally, easy catalyst recovery and recycling is achieved. The system is based on phosphorus ligands covalently attached to a polymeric support, which is readily obtained by a solid-phase synthetic protocol.






Metal–Organic Frameworks


J.-P. Zheng, S. Ou, M. Zhao, C.-D. Wu*

A Highly Sensitive Luminescent Dye@MOF Composite for Probing Different Volatile Organic Compounds

Metal–organic framework sensor: We report an interesting layered coordination network based on the derivative of the natural amino acid L-phenylalanine and zinc ions, which can embed Rhodamine B dye to result in a luminescent sensor of Rho@MOF composite, which can detect different volatile organic compounds with clearly different and unique luminescence readouts by tuning the ligand-to-dye energy transfer (see figure).



ChemPlusChem
DOI: 10.1002/cplu.201600057




Metal–Organic Frameworks


X. An, D. Shin, J. Jeong, J. Lee*

Metal-Derived Mesoporous Structure of a Carbon Nanofiber Electrocatalyst for Improved Oxygen Evolution Reaction in Alkaline Water Electrolysis

Cheaper catalysts: The lowest overpotential value of the oxygen evolution reaction with cost-effective carbon nanofiber (CNF) electrocatalysts is reported. This can be attributed to the homogeneous placement of active species in the metal-derived mesoporous structure.



ChemElectroChem
DOI: 10.1002/celec.201600072




MOFs and Sensors


W. Yang, G. Chang, H. Wang,* T.-L. Hu, Z. Yao, K. Alfooty, S. Xiang, B. Chen

A Three-Dimensional Tetraphenylethene-Based Metal–Organic Framework for Selective Gas Separation and Luminescence Sensing of Metal Ions

A three-dimensional multifunctional metal–organic framework (MOF), UTSA-86, made up of cadmium ions and a tetraphenylethene-based tetracarboxylate ligand has been synthesized and characterized by single-crystal X-ray diffraction analysis. This MOF exhibits permanent porosity, selective gas uptake, and luminescence sensing of metal ions.



Eur. J. Inorg. Chem.
DOI: 10.1002/ejic.201600201




Natural Product Synthesis


M. Karak, J. A. M. Acosta, L. C. A. Barbosa,* J. Boukouvalas

Late-Stage Bromination Enables the Synthesis of Rubrolides B, I, K, and O

The first total synthesis of the marine natural metabolites rubrolides B, I, K and O is described; this approach employs a site-selective Suzuki coupling, vinylogous aldol condensation and a late-stage regioselective bromination.



Eur. J. Org. Chem.
DOI: 10.1002/ejoc.201600473



ChemistryOpen


DOI: 10.1002/open.201600009

Catalysis

S. H. Mir, B. Ochiai*

Development of Hierarchical Polymer@Pd Nanowire-Network: Synthesis and Application as Highly Active Recyclable Catalyst and Printable Conductive Ink

Facile but versatile: The facile one-pot preparation and directed growth of a multifunctional Pd nanowire-network on interconnected hollow polymer nanospheres is reported. The Pd nanowire-network served as a highly active recyclable catalyst for Mizoroki–Heck and Suzuki–Miyaura couplings. Inkjet printing of the nanosphere ink produced a highly conductive pattern without annealing.



Asian J. Org. Chem.



DOI: 10.1002/ajoc.201600173

Heterogeneous Catalysis

A. Singh, G. Ramanathan*

Rational Design of Heterogeneous Silver Catalysts by Exploitation of Counteranion-Induced Coordination Geometry

Green machine: The coordination complexes of a green fluorescent protein (GFP) chromophore analogue with a variety of silver salts have been synthesized by varying the counteranion. The counteranion influences the geometry and coordination number of the silver ion in these complexes. Complexes that have a free coordination site have been exploited as heterogeneous catalysts for cyclization reactions under “green” conditions.



ChemNanoMat

DOI: 10.1002/cnma.201600078

Drug Delivery

S. Y. Tan, C. Y. Ang, A. Mahmood, Q. Qu, P. Li, R. Zou,* Y. Zhao*

Doxorubicin-Loaded Metal–Organic Gels for pH and Glutathione Dual-Responsive Release

Metal–organic gels are capable of storing anticancer drugs within the pores, and then releasing them under acidic pH or in the presence of glutathione. In vitro experiments show a low toxicity of metal–organic gels and confirm the triggered drug release.



ChemViews magazine

DOI: 10.1002/chemv.201600053

Analytical Chemistry

D. Bradley

Jackson Pollock's “Alchemy” Analyzed

Analyzing the chemical makeup of paintings without damaging them can be difficult. Researchers from Italy have used non-invasive spectroscopy methods to reveal the technique and paint palette of famous artist Jackson Pollock – without needing to scrape off or cut out samples.

