

# A Single Fluorescent Probe to Visualize Hydrogen Sulfide and Hydrogen Polysulfides with Different Fluorescence Signals

Wei Chen, Armando Pacheco, Yoko Takano, Jacob J. Day, Kenjiro Hanaoka, and Ming Xian\*

**Abstract:** Hydrogen sulfide ( $H_2S$ ) and hydrogen polysulfides ( $H_2S_n$ ,  $n > 1$ ) are endogenous regulators of many physiological processes. In order to better understand the symbiotic relationship and cellular cross-talk between  $H_2S$  and  $H_2S_n$ , it is highly desirable to develop single fluorescent probes which enable dual-channel discrimination between  $H_2S$  and  $H_2S_n$ . Herein, we report the rational design, synthesis, and evaluation of the first dual-detection fluorescent probe **DDP-1** that can visualize  $H_2S$  and  $H_2S_n$  with different fluorescence signals. The probe showed high selectivity and sensitivity to  $H_2S$  and  $H_2S_n$  in aqueous media and in cells.

Due to their vital physiological functions, reactive sulfur species (RSS) form an important and ever-increasing research field.<sup>[1]</sup> Among RSS,  $H_2S$  is perhaps most attractive as it has been characterized as a crucial gaseous transmitter.<sup>[2]</sup> While the research on  $H_2S$  is still actively ongoing, a new hot topic about RSS has recently emerged which focuses on the chemical biology of hydrogen polysulfides ( $H_2S_n$ ,  $n > 1$ ).<sup>[3]</sup> Endogenous  $H_2S_n$  may be generated from  $H_2S$  upon reacting with reactive oxygen species (ROS) like  $ClO^-$ .<sup>[4]</sup> Cystathionine  $\gamma$ -lyase (CSE) and cystathionine- $\beta$ -synthase (CBS), the enzymes responsible for  $H_2S$  biosynthesis, were found to also produce persulfides (RSSH), which could further be converted to  $H_2S_n$ .<sup>[5]</sup> Very recently 3-mercaptopyruvate sulfurtransferase (3MST) was identified to be an important enzyme in brain for  $H_2S_n$  generation.<sup>[6]</sup> The significance of  $H_2S_n$  in redox biology has only been recognized recently. Evidences suggest that  $H_2S_n$  might be the actual signaling molecules that activate ion channels, transcription factors, and tumor suppressors with higher potency than  $H_2S$ .<sup>[3,7]</sup> One example is protein S-sulfhydration,<sup>[5,8]</sup> which was previously thought to be resulted from  $H_2S$ . Recent studies found that  $H_2S_n$  were much more effective in causing S-sulfhydration than  $H_2S$ .<sup>[3,7]</sup>

Given the importance of  $H_2S$  and  $H_2S_n$  in redox biology, convenient and accurate detection methods for these species are invaluable research tools. In the past five years, the detection of  $H_2S$  has received wide attention and a large number of fluorescent probes for  $H_2S$  have been developed.<sup>[9]</sup>

It should be noted that in all of those studies the selectivity of the probes for  $H_2S$  versus  $H_2S_n$  was not considered, mainly because the significance of  $H_2S_n$  in biological samples was not recognized. On the other hand, the detection of  $H_2S_n$  is much less studied, due to very limited understanding of the chemical reactivity of  $H_2S_n$ .<sup>[10]</sup> Very recently our laboratory and several others have developed a few fluorescent probes for  $H_2S_n$ .<sup>[11]</sup> In these works the selectivity of probes for  $H_2S_n$  versus  $H_2S$  was verified and  $H_2S$  did not trigger any fluorescent signals for these probes.

With the increasing knowledge available for sensing  $H_2S$  and  $H_2S_n$ , it is now possible to develop fluorescent probes for dual-channel differentiation of  $H_2S$  and  $H_2S_n$ . Such probes will be very useful for understanding the mutual relationship and cellular cross-talk between  $H_2S$  and  $H_2S_n$ . While simply mixing two specific probes might be able to detect both analytes with distinct fluorescence signals, this strategy suffers from limitations such as 1) potential interference between two probes, 2) larger invasive effects, and 3) possible different localization and metabolisms of the probes.<sup>[12]</sup> Therefore, it is highly desirable (but also challenging) to develop a single fluorescent probe which enable visualization of  $H_2S$  and  $H_2S_n$  using different emission channels. Such dual-detection probes have not been reported so far. Herein, we report the rational design, synthesis, and evaluation of a single fluorescent probe that can differentiate  $H_2S$  and  $H_2S_n$  in aqueous media and in cells.


As shown in Scheme 1, we expected the dual-detection probe would contain two separated and quenched fluorophores A and B. Ideally each pseudo-fluorophore should only react with one species ( $H_2S$  or  $H_2S_n$ ) to cause the corresponding fluorescence “turn-on”. As such,  $H_2S$  and  $H_2S_n$  can be determined by different fluorescence emission wavelengths. Even if  $H_2S$  and  $H_2S_n$  are present together, the ratio of the two emission intensity might be able to determine the ratio of the two sulfur species.

To achieve this design, the key is to construct two specific reactive sites on the probe for  $H_2S$  and  $H_2S_n$ . In our studies of  $H_2S_n$  probes, we found phenyl 2-(benzoylthio)benzoate-based

[\*] Dr. W. Chen, A. Pacheco, J. J. Day, Prof. Dr. M. Xian  
Department of Chemistry, Washington State University  
Pullman, WA 99164 (USA)  
E-mail: mxian@wsu.edu

Y. Takano, Prof. Dr. K. Hanaoka  
Graduate School of Pharmaceutical Sciences, The University of Tokyo  
Tokyo 113-0033 (Japan)


Supporting information and the ORCID identification number(s) for the author(s) of this article can be found under <http://dx.doi.org/10.1002/anie.201604892>.



**Scheme 1.** The design of dual detection probes for  $H_2S$  and  $H_2S_n$ .


probes exhibited high sensitivity and selectivity for  $\text{H}_2\text{S}_n$ .<sup>[11b]</sup> Importantly,  $\text{H}_2\text{S}$  did not show any response to this type of probes. Therefore, phenyl 2-(benzoylthio)benzoate should be a suitable choice as  $\text{H}_2\text{S}_n$  reactive site. On the other hand, the selection of the  $\text{H}_2\text{S}$  reaction site was more difficult as previously reported  $\text{H}_2\text{S}$  probes did not verify their selectivity for  $\text{H}_2\text{S}_n$ . So far three types of reactions have been used in the development of  $\text{H}_2\text{S}$  probes: a)  $\text{H}_2\text{S}$ -mediated reductions (mostly using azides), b)  $\text{H}_2\text{S}$ -mediated nucleophilic reactions, and c) metal-sulfide formations.<sup>[9]</sup> As  $\text{H}_2\text{S}_n$  are expected to be stronger nucleophiles than  $\text{H}_2\text{S}$ , nucleophilic reactions or metal-sulfide formations are likely to occur with  $\text{H}_2\text{S}_n$ , leading to poor selectivity.  $\text{H}_2\text{S}$ -mediated azide reduction might be useful.  $\text{H}_2\text{S}_n$  are oxidation products of  $\text{H}_2\text{S}$  and might have weaker reducing ability (at least for some azides). Therefore, azide-based fluorophores might be able to differentiate  $\text{H}_2\text{S}$  from  $\text{H}_2\text{S}_n$ .

With this consideration in mind, we first compared the fluorescence responses of several azide-based fluorophores to  $\text{H}_2\text{S}$  and  $\text{H}_2\text{S}_n$ .  $\text{Na}_2\text{S}$  and  $\text{Na}_2\text{S}_2$  were used in buffers as  $\text{H}_2\text{S}$  and  $\text{H}_2\text{S}_n$  equivalents, respectively. One example using 4-methyl-7-azidocoumarin (C7-Az) is shown in Figure 1 and Figure S1 (see the Supporting Information). As expected, the treatment of C7-Az with  $\text{H}_2\text{S}_n$  led to much decreased fluorescence signals as compared to  $\text{H}_2\text{S}$  treatment. These results indicated this azide moiety has higher reactivity to  $\text{H}_2\text{S}$  than  $\text{H}_2\text{S}_n$ , and could be used in dual-detection.



**Figure 1.** Time-dependent fluorescence intensity changes of C7-Az (10  $\mu\text{M}$ ) with 50  $\mu\text{M}$   $\text{Na}_2\text{S}$  (■) or 50  $\mu\text{M}$   $\text{Na}_2\text{S}_2$  (●).

Next we proposed a dual-detection probe **DDP-1** (Scheme 2). In this design, four factors were considered: (1) coumarin and rhodol were selected as two fluorophores due to their excellent solubility, high quantum yields, and well-separated maximum emission wavelengths (ca. 445 nm for coumarin and ca. 542 nm for rhodol). As such, dual-color imaging of  $\text{H}_2\text{S}$  and  $\text{H}_2\text{S}_n$  from different emission channels was possible. (2) A rigid piperazine linker was used to bridge the two fluorophores and provided an advantage for Förster resonance energy transfer (FRET) in the coumarin-rhodol scaffold, which preventing the  $\pi$ - $\pi$  stacking between dyes.<sup>[13]</sup> (3) Azidation of coumarin and phenyl 2-(benzoylthio)benzoate-protected rhodol should effectively quench the fluorescence of the probe via the intramolecular charge transfer (ICT) effect and the intramolecular spirocyclization, respectively. The probe should bear very low background fluores-



**Scheme 2.** Possible reactions of **DDP-1** with  $\text{H}_2\text{S}$  and  $\text{H}_2\text{S}_n$ .


cence that is favorable for high sensitivity. (4) The azide and phenyl 2-(benzoylthio)benzoate moieties provide selective reaction sites for  $\text{H}_2\text{S}$  and  $\text{H}_2\text{S}_n$ .

The fluorescence turn-on mechanism of **DDP-1** is also proposed in Scheme 2. When the probe is treated with  $\text{H}_2\text{S}_n$ , phenyl 2-(benzoylthio)benzoate should be preferably reacted to release the fluorescence of rhodol (to form **1**). Even if the azide group of **1** is partially reduced by  $\text{H}_2\text{S}_n$  to form a small amount of **3**, it should not affect the fluorescence emission channel due to FRET between the two fluorophores (Figure S2). Overall the reaction with  $\text{H}_2\text{S}_n$  should only produce green fluorescence of rhodol. In contrast, the reaction between **DDP-1** and  $\text{H}_2\text{S}$  is more complicated. Previous results have demonstrated that  $\text{H}_2\text{S}$  cannot turn on phenyl 2-(benzoylthio)benzoate-based fluorophore.<sup>[11b]</sup> Therefore we expected  $\text{H}_2\text{S}$  would preferably react with the azide moiety to produce **2** and release blue fluorescence of coumarin. It should be noted that recent studies demonstrated that the reaction of  $\text{H}_2\text{S}$  with azides led to the formation of  $\text{H}_2\text{S}_n$ .<sup>[14]</sup> Therefore, **3** would also be formed in this process, which should exhibit green fluorescence of rhodol because of FRET. However, less than 0.5 equivalent of  $\text{H}_2\text{S}_n$  are generated from the reaction of 1 equivalent of  $\text{H}_2\text{S}$  and azide. Moreover, the reaction with phenyl 2-(benzoylthio)benzoate consumes at least 2 equivalent of  $\text{H}_2\text{S}_n$ . Therefore, only a small amount of **3** would be produced during this process. Overall the reaction between **DDP-1** and  $\text{H}_2\text{S}$  should produce the emission signals of both coumarin (major) and rhodol (minor). Taken together, it is anticipated that probe **DDP-1** could detect  $\text{H}_2\text{S}$  and  $\text{H}_2\text{S}_n$  from distinct emission channels.

The proposed probe was then prepared and characterized (see Supporting Information). We first tested the probe's fluorescence response to  $\text{H}_2\text{S}$  and  $\text{H}_2\text{S}_n$  in PBS buffers under the excitation wavelength of coumarin ( $\lambda_{\text{ex}} = 360$  nm). As shown in Figure S3, **DDP-1** alone showed almost no fluorescence. Upon reacting with  $\text{Na}_2\text{S}$  or  $\text{Na}_2\text{S}_2$ , **DDP-1** gave appreciable fluorescence enhancements in one hour. As expected, different fluorescence emission behaviors were observed for  $\text{Na}_2\text{S}$  and  $\text{Na}_2\text{S}_2$ .  $\text{Na}_2\text{S}$  led to two distinct emissions at 452 nm and 542 nm (blue–green fluorescence). While  $\text{Na}_2\text{S}_2$  induced only one strong emission at 542 nm (green fluorescence). Therefore,  $\text{H}_2\text{S}$  could be easily identified by the simultaneous appearance of the two well-

separated emissions, whereas  $\text{H}_2\text{S}_n$  could be characterized by the single emission at 542 nm. In addition, these distinguished fluorescence color changes might be favorable for the simple detection of  $\text{H}_2\text{S}$  and  $\text{H}_2\text{S}_n$  by the naked eyes.

Next we studied the sensitivity of **DDP-1** for  $\text{H}_2\text{S}$  and  $\text{H}_2\text{S}_n$  using varied concentrations of  $\text{Na}_2\text{S}$  or  $\text{Na}_2\text{S}_2$  (0 to 150  $\mu\text{M}$ ). As shown in Figure 2, the increase in fluorescence




**Figure 2.** Fluorescence spectra of **DDP-1** (10  $\mu\text{M}$ ) under various concentrations of a)  $\text{H}_2\text{S}$  (0, 1, 5, 10, 20, 40, 75, 100, 120, 150  $\mu\text{M}$  for curves 1–10, respectively); b)  $\text{H}_2\text{S}_n$  (0, 1, 5, 10, 20, 45, 70, 100, 120, 150  $\mu\text{M}$  for curves 1–10, respectively).


intensity ( $\lambda_{\text{ex}} = 360 \text{ nm}$ ) with the gradual increase of  $\text{Na}_2\text{S}$  or  $\text{Na}_2\text{S}_2$  concentrations was observed. A good linear relation was obtained (Figure S4). For  $\text{H}_2\text{S}$ , the fluorescence intensity at 452 nm increased linearly with  $\text{Na}_2\text{S}$  concentration from 0 to 20  $\mu\text{M}$ . The fluorescence intensity at 542 nm increased linearly with  $\text{Na}_2\text{S}$  in the concentration range of 0 to 40  $\mu\text{M}$ . The detection limits ( $S/N = 3$ ) were 100 nM and 150 nM for  $\text{H}_2\text{S}$ , corresponding emission at 452 and 542 nm. The fluorescence intensity at 542 nm increased linearly with  $\text{Na}_2\text{S}_2$  concentrations from 0 to 20  $\mu\text{M}$ . The detection limit was calculated to be 24 nM. These results indicated that **DDP-1** displayed much higher sensitivity to  $\text{H}_2\text{S}_n$  than  $\text{H}_2\text{S}$ . The effects of pH in these reactions were also studied. **DDP-1** was found to work effectively at neutral to basic pH (7–10) (Figure S5).

We then wondered if the probe **DDP-1** could give meaningful responses when  $\text{H}_2\text{S}$  and  $\text{H}_2\text{S}_n$  co-exist. To this end, we tested fluorescence changes of varying  $\text{Na}_2\text{S}_2/\text{Na}_2\text{S}$  mixture solutions, while the total sulfur concentration was fixed in samples (200  $\mu\text{M}$ ). As  $\text{H}_2\text{S}$  is a much more stable species than  $\text{H}_2\text{S}_n$  and the concentration of  $\text{H}_2\text{S}$  is likely to be higher than  $\text{H}_2\text{S}_n$  in biological systems, we varied  $[\text{H}_2\text{S}_2]/[\text{H}_2\text{S}]$  ratios from 0 to 1. The fluorescence signals of these solutions were measured by the probe. As shown in Figure 3, following the increases of  $[\text{H}_2\text{S}_2]/[\text{H}_2\text{S}]$  ratios, the emission at 452 nm decreased with a concurrent increase at 542 nm. The  $F_{542\text{nm}}/F_{452\text{nm}}$  ratios increased linearly with  $[\text{H}_2\text{S}_2]/[\text{H}_2\text{S}]$  ratios in the range of 0 to 0.176 (Figure S6). These results indicated the probe could be used for the ratiometric detection of relative  $\text{H}_2\text{S}_n$  and  $\text{H}_2\text{S}$  concentrations when they coexist.

To verify the specificity of **DDP-1** for  $\text{H}_2\text{S}$  and  $\text{H}_2\text{S}_n$ , its responses to a series of biologically relevant RSS (GSH, Cys, Hcy, GSSG,  $\text{SO}_3^{2-}$ ,  $\text{S}_2\text{O}_3^{2-}$ ,  $\text{CH}_3\text{SSSCH}_3$ , and  $\text{S}_8$ ) were tested. As shown in Figure 4, these RSS did not cause any fluorescence increase. Only  $\text{Na}_2\text{S}_2$ ,  $\text{Na}_2\text{S}_3$ , and  $\text{Na}_2\text{S}_4$  triggered significant fluorescence increases. We also examined the



**Figure 3.** Fluorescence spectra ( $\lambda_{\text{ex}} = 360 \text{ nm}$ ) of **DDP-1** (10  $\mu\text{M}$ ) with varying  $\text{Na}_2\text{S}_2/\text{Na}_2\text{S}$  mixture solutions ( $\text{Na}_2\text{S}_2/\text{Na}_2\text{S}$  ratios were 0, 0.01, 0.026, 0.053, 0.081, 0.111, 0.176, 0.333, 0.667, 1 for curves 1–10, respectively).




**Figure 4.** Fluorescence spectra ( $\lambda_{\text{ex}} = 360 \text{ nm}$ ) of **DDP-1** (10  $\mu\text{M}$ ) in the presence of various reactive species. (1) Probe only; (2) 5 mM GSH; (3) 1 mM Cys; (4) 100  $\mu\text{M}$  Hcy; (5) 100  $\mu\text{M}$  GSSG; (6) 100  $\mu\text{M}$   $\text{Na}_2\text{S}_2\text{O}_3$ ; (7) 100  $\mu\text{M}$   $\text{Na}_2\text{SO}_3$ ; (8) 50  $\mu\text{M}$   $\text{CH}_3\text{SSSCH}_3$ ; (9) 50  $\mu\text{M}$   $\text{S}_8$ ; (10) 250  $\mu\text{M}$   $\text{H}_2\text{O}_2$ ; (11) 50  $\mu\text{M}$   $\text{ClO}^-$ ; (12) 50  $\mu\text{M}$   $\text{O}_2^-$ ; (13) 50  $\mu\text{M}$   $\cdot\text{OH}$ ; (14) 50  $\mu\text{M}$   $^1\text{O}_2$ ; (15) 100  $\mu\text{M}$  Alanine; (16) 100  $\mu\text{M}$  Serine; (17) 100  $\mu\text{M}$  Arginine; (18) 100  $\mu\text{M}$  Isoleucine; (19) 100  $\mu\text{M}$  Lysine; (20) 100  $\mu\text{M}$  Ascorbic acid; (21) 100  $\mu\text{M}$   $\text{Na}_2\text{S}$ ; (22) 50  $\mu\text{M}$   $\text{Na}_2\text{S}_2$ ; (23) 50  $\mu\text{M}$   $\text{Na}_2\text{S}_3$ ; (24) 50  $\mu\text{M}$   $\text{Na}_2\text{S}_4$ ; (25) 100  $\mu\text{M}$   $\text{Na}_2\text{S} + 50 \mu\text{M}$   $\text{ClO}^-$ .

responses of **DDP-1** to common ROS, such as  $\text{H}_2\text{O}_2$ ,  $\text{ClO}^-$ , superoxide ( $\text{O}_2^-$ ), hydroxy radical ( $\cdot\text{OH}$ ), and singlet oxygen ( $^1\text{O}_2$ ). No fluorescence increase was detected for these species. Moreover, as  $\text{H}_2\text{S}_n$  could be efficiently generated from  $\text{H}_2\text{S}$  and  $\text{ClO}^-$ , the probe was used to analyze in situ generation of  $\text{H}_2\text{S}_n$ . When  $\text{Na}_2\text{S}$  (100  $\mu\text{M}$ ) and  $\text{ClO}^-$  (50  $\mu\text{M}$ ) co-existed, a very strong fluorescence emission at 542 nm was observed. We also tested the specificity of **DDP-1** under the excitation wavelength of rhodol ( $\lambda_{\text{ex}} = 515 \text{ nm}$ ).  $\text{H}_2\text{S}_n$  induced significant fluorescence increase while  $\text{H}_2\text{S}$  gave only small fluorescent enhancement (Figure S7). Almost no fluorescence changes were detected for other RSS or ROS. The responses of **DDP-1** to representative amino acids and ascorbic acid were also tested, they did not induce any response. These results demonstrated the excellent specificity of **DDP-1** for  $\text{H}_2\text{S}$  and  $\text{H}_2\text{S}_n$ .

Finally we evaluated the ability of **DDP-1** to detect  $\text{H}_2\text{S}$  and  $\text{H}_2\text{S}_n$  in cells. As shown in Figure 5, HeLa cells were first incubated with **DDP-1** (20  $\mu\text{M}$ ) for 30 minutes. Then extracellular probe was washed off. Only very weak fluorescence was observed in the blue and green channels. When cells were treated with  $\text{Na}_2\text{S}$  (100  $\mu\text{M}$ ), an apparent fluorescence enhancement was detected in the blue and green channel. When





**Figure 5.** HeLa cells were incubated with **DDP-1** (20  $\mu\text{M}$ ) for 30 minutes, then washed, and subjected to different treatments. a,e) Controls (no added  $\text{Na}_2\text{S}$ ,  $\text{NaClO}$ , or  $\text{Na}_2\text{S}_2$ ); b,f)  $\text{Na}_2\text{S}$  (100  $\mu\text{M}$ ); c,g) the mixture of  $\text{Na}_2\text{S}$  (100  $\mu\text{M}$ ) and  $\text{NaClO}$  (50  $\mu\text{M}$ ); d,h)  $\text{Na}_2\text{S}_2$  (50  $\mu\text{M}$ ). a–d) Fluorescence image of HeLa cells at the blue channel; e–h) fluorescence image of the corresponding image (a–d) from the green channel.

cells were treated with  $\text{Na}_2\text{S}_2$  or in situ generated  $\text{H}_2\text{S}_n$ , the fluorescence signals in the blue channel decreased with a coincstantaneous increase in the green channel. These results proved that **DDP-1** is cell membrane permeable and could be used for the detection of  $\text{H}_2\text{S}$  and  $\text{H}_2\text{S}_n$  from distinct emission channels in cells. Additionally, the cell viability assay implied that **DDP-1** has low cytotoxicity and good biocompatibility (Figure S8).

In summary, we report in this study the rational design, synthesis, and evaluation of the first single fluorescent probe **DDP-1** that can clearly differentiate  $\text{H}_2\text{S}$  and  $\text{H}_2\text{S}_n$  with different fluorescence signals. This also represents a significant improvement in the development of  $\text{H}_2\text{S}$  probes as previously reported  $\text{H}_2\text{S}$  probes can hardly discriminate  $\text{H}_2\text{S}_n$ . This novel probe is expected to serve as a useful tool in understanding the redox signaling of  $\text{H}_2\text{S}$  and  $\text{H}_2\text{S}_n$ .

## Acknowledgements

This work is supported by NIH (grant number R01HL116571).

**Keywords:** cyclization · fluorescence · fluorescent probes · hydrogen polysulfides · sulfides

**How to cite:** *Angew. Chem. Int. Ed.* **2016**, 55, 9993–9996  
*Angew. Chem.* **2016**, 128, 10147–10150

- [1] a) C. E. Paulsen, K. S. Carroll, *Chem. Rev.* **2013**, 113, 4633; b) Y. H. Seo, K. S. Carroll, *Angew. Chem. Int. Ed.* **2011**, 50, 1342; *Angew. Chem.* **2011**, 123, 1378; c) T. H. Poole, J. A. Reisz, W. Zhao, L. B. Poole, C. M. Furdui, S. B. King, *J. Am. Chem. Soc.* **2014**, 136, 6167; d) J. L. Miljkovic, I. Kenkel, I. Ivanovic-Burmazovic, M. R. Filipovic, *Angew. Chem. Int. Ed.* **2013**, 52, 12061; *Angew. Chem.* **2013**, 125, 12283; e) X. Yang, Y. Guo, M. Strongin, *Angew. Chem. Int. Ed.* **2011**, 50, 10690; *Angew. Chem.* **2011**, 123, 10878; f) D. Zhang, I. Macinkovic, N. O. Devarie-Baez, J. Pan, C. Park, K. S. Carroll, M. R. Filipovic, M. Xian, *Angew. Chem. Int. Ed.* **2014**, 53, 575; *Angew. Chem.* **2014**, 126, 586; g) Y. Zheng, B. Yu, K. Ji, Z. Pan, V. Chittavong, B. Wang,

*Angew. Chem. Int. Ed.* **2016**, 55, 4514; *Angew. Chem.* **2016**, 128, 4590.

- [2] a) J. M. Fukuto, S. J. Carrington, D. J. Tantillo, J. G. Harrison, L. J. Ignarro, B. A. Freeman, A. Chen, D. A. Wink, *Chem. Res. Toxicol.* **2012**, 25, 769; b) R. Wang, *Physiol. Rev.* **2012**, 92, 791.
- [3] a) S. Koike, Y. Ogasawara, N. Shibuya, H. Kimura, K. Ishii, *FEBS Lett.* **2013**, 587, 3548; b) R. Greiner, Z. Pálincás, K. Bäsell, D. Becher, H. Antelmann, P. Nagy, T. P. Dick, *Antioxid. Redox Signaling* **2013**, 19, 1749; c) K. Ono, T. Akaike, T. Sawa, Y. Kumagai, D. Wink, D. J. Tantillo, A. J. Hobbs, P. Nagy, M. Xian, J. Lin, J. M. Fukuto, *Free Radical Biol. Med.* **2014**, 77, 82; d) H. Kimura, *Antioxid. Redox Signaling* **2015**, 22, 362; e) E. C. Estevam, L. Faulstich, S. Griffin, T. Burkholz, C. Jacob, *Curr. Org. Chem.* **2016**, 20, 211.
- [4] P. Nagy, C. C. Winterbourn, *Chem. Res. Toxicol.* **2010**, 23, 1541.
- [5] a) T. Ida, T. Sawa, H. Ihara, Y. Tsuchiya, Y. Watanabe, Y. Kumagai, M. Suematsu, H. Motohashi, S. Fujii, T. Matsunaga, M. Yamamoto, K. Ono, N. O. Devarie-Baez, M. Xian, J. M. Fukuto, T. Akaike, *Proc. Natl. Acad. Sci. USA* **2014**, 111, 7606; b) P. K. Yadav, M. Martinov, V. Vitvitsky, J. Seravalli, R. Wedmann, M. R. Filipovic, R. Banerjee, *J. Am. Chem. Soc.* **2016**, 138, 289.
- [6] Y. Kimura, Y. Toyofuku, S. Koike, N. Shibuya, N. Nagahara, D. Lefer, Y. Ogasawara, H. Kimura, *Sci. Rep.* **2015**, 5, 14774.
- [7] a) T. V. Mishanina, M. Libiad, R. Banerjee, *Nat. Chem. Biol.* **2015**, 11, 457; b) H. Kimura, *Proc. Jpn. Acad. Ser. B* **2015**, 91, 131.
- [8] a) N. Krishnan, C. Fu, D. J. Pappin, N. K. Tonks, *Sci. Signaling* **2011**, 4, ra86; b) N. Sen, B. D. Paul, M. M. Gadalla, A. K. Mustafa, T. Sen, R. Xu, S. Kim, S. H. Snyder, *Mol. Cell* **2012**, 45, 13; c) G. Yang, K. Zhao, Y. Ju, S. Mani, Q. Cao, S. Puukila, N. Khaper, L. Wu, R. Wang, *Antioxid. Redox Signaling* **2013**, 18, 1906.
- [9] Reviews: a) H. Peng, W. Chen, Y. Cheng, L. Hakuna, R. Strongin, B. Wang, *Sensors* **2012**, 12, 15907; b) X. Li, X. Gao, W. Shi, H. Ma, *Chem. Rev.* **2014**, 114, 590; c) V. S. Lin, W. Chen, M. Xian, C. J. Chang, *Chem. Soc. Rev.* **2015**, 44, 4596; d) X. Zhou, S. Lee, Z. Xu, J. Yoon, *Chem. Rev.* **2015**, 115, 7944; e) M. D. Hartle, M. D. Pluth, *Chem. Soc. Rev.* **2016**, DOI: 10.1039/c6cs00212a. f) W. Xuan, C. Sheng, Y. Cao, W. He, W. Wang, *Angew. Chem. Int. Ed.* **2012**, 51, 2282; *Angew. Chem.* **2012**, 124, 2328.
- [10] a) R. Steudel, *Top. Curr. Chem.* **2003**, 231, 99; b) R. Steudel, *Top. Curr. Chem.* **2003**, 231, 127.
- [11] For selected references, see: a) C. Liu, W. Chen, W. Shi, B. Peng, Y. Zhao, H. Ma, M. Xian, *J. Am. Chem. Soc.* **2014**, 136, 7257; b) W. Chen, E. W. Rosser, T. Matsunaga, A. Pacheco, T. Akaike, M. Xian, *Angew. Chem. Int. Ed.* **2015**, 54, 13961; *Angew. Chem.* **2015**, 127, 14167; c) W. Chen, E. W. Rosser, D. Zhang, W. Shi, Y. Li, W. Dong, H. Ma, D. Hu, M. Xian, *Org. Lett.* **2015**, 17, 2776; d) L. Zeng, S. Chen, T. Xia, W. Hu, C. Li, Z. Liu, *Anal. Chem.* **2015**, 87, 3004; e) M. Gao, F. Yu, H. Chen, L. Chen, *Anal. Chem.* **2015**, 87, 3631.
- [12] a) H. Komatsu, T. Miki, D. Citterio, T. Kubota, Y. Shindo, Y. Kitamura, K. Oka, K. Suzuki, *J. Am. Chem. Soc.* **2005**, 127, 10798; b) L. Yuan, W. Lin, Y. Xie, B. Chen, S. Zhu, *J. Am. Chem. Soc.* **2012**, 134, 1305; c) G. C. Van de Bittner, C. R. Bertozzi, C. J. Chang, *J. Am. Chem. Soc.* **2013**, 135, 1783; d) M. D. Hammers, M. D. Pluth, *Anal. Chem.* **2014**, 86, 7135.
- [13] a) L. Yuan, W. Lin, K. Zheng, S. Zhu, *Acc. Chem. Res.* **2013**, 46, 1462; b) W. Xuan, Y. Cao, J. Zhou, W. Wang, *Chem. Commun.* **2013**, 49, 10474.
- [14] H. A. Henthorn, M. D. Pluth, *J. Am. Chem. Soc.* **2015**, 137, 15330.

Received: May 18, 2016

Published online: July 13, 2016