


Cycloaddition Reactions

Deutsche Ausgabe: DOI: 10.1002/ange.201604371
Internationale Ausgabe: DOI: 10.1002/anie.201604371Rhenium(I)-Catalyzed Generation of α,β -Unsaturated Carbene Complex Intermediates from Propargyl Ethers for the Preparation of Cycloheptadiene Derivatives

Hideyuki Sogo and Nobuharu Iwasawa*


Abstract: The rhenium(I)-catalyzed generation of α,β -unsaturated carbene complex intermediates from easily available propargyl ethers was achieved for the concise construction of cycloheptadiene derivatives through the formal [4+3] cycloaddition reaction with siloxydienes.

Catalytic generation of α,β -unsaturated carbene complex intermediates through electrophilic activation of alkynes has attracted much attention as a useful method for the reaction of novel three-carbon reactive species.^[1–4] In these reactions, the nucleophilic addition of appropriate internal nucleophiles onto alkynes having a leaving group at the propargylic position, followed by electron donation from the metal to release the leaving group, generates α -Nu-substituted α,β -unsaturated carbene complex intermediates (Scheme 1).



Scheme 1. Generation of α,β -unsaturated carbene complex intermediates.

We then thought of the possibility of using easily available propargyl ethers as substrate for catalytic generation of α,β -unsaturated carbene complex intermediates with the expectation that hydride transfer from the ether moiety would be promoted by electron donation from the lone pair of the oxygen atom (Scheme 2).^[5–8] For generation of α,β -unsaturated carbene complex intermediates, the reaction necessitates 1,4-hydride transfer onto the π -alkyne complex **A**, which is rather rare in such systems,^[6,9] and we also thought of examining the possibility of 1,5-hydride transfer through vinylidene intermediate **B**, which has several precedents as a stoichiometric reaction for preparation of metathesis catalysts.^[10–13] These hydride transfers would generate the same intermediate **C** containing an oxonium moiety, which would work as an efficient leaving group for carbene generation. Herein, we report our successful realization of this concept for catalytic generation of simple α,β -unsaturated carbene complexes from propargyl ethers and their



Scheme 2. Generation of α,β -unsaturated carbene complex intermediates from propargyl ethers.

utilization as a three-carbon unit for the concise preparation of cycloheptadiene derivatives.

The reaction was examined with various metal catalysts using the benzyl ether of 1,1-dimethyl-2-propyn-1-ol **1a** and 1.5 molar amounts of 4-aryl-2-siloxy-1,3-butadiene **2a** as a carbene trapping reagent expecting the formation of cycloheptadiene **3a** as [4+3]-cycloadduct (Table 1).^[4i,j] It was found that $\text{ReX}(\text{CO})_5$ and PtCl_2 were the active catalyst for this reaction, and the desired product **3a** was obtained by carrying out of the reaction in 1,4-dioxane at 100 °C along with a small amount of its olefinic isomerization product

Table 1: Examination of various reaction conditions.

Entry	Metal cat.	Time [h]	3a + 4a [%] ^[a]	3a : 4a [%] ^[b]
1	PtCl_2	12	33	93:7
2	$\text{ReCl}(\text{CO})_5$	12	62	> 95:5
3	$\text{ReBr}(\text{CO})_5$	4	84	95:5
4	$\text{ReI}(\text{CO})_5$	2	96	> 95:5
5 ^[c]	$\text{ReI}(\text{CO})_5$	4	97	> 95:5
6 ^[c,d]	$\text{ReI}(\text{CO})_5$	4	75	61:39
7 ^[c,d,e]	$\text{ReI}(\text{CO})_5$	4	90	> 95:5

Reaction conditions : **1a** (0.10 mmol), **2a** (0.15 mmol), and Metal cat. (0.01 mmol) in 1,4-dioxane (2.0 mL) at 100 °C. [a] Yield of isolated product. [b] Determined by ^1H NMR spectroscopy. [c] 0.10 M of **1a** and 2.5 mol% of $\text{ReI}(\text{CO})_5$ were used. [d] 1.0 equiv of **2a** was used. [e] Activated 4 Å M.S. were used.

[*] H. Sogo, Prof. Dr. N. Iwasawa

Department of Chemistry, Tokyo Institute of Technology
2-12-1, O-okayama, Meguro-ku, Tokyo 152-8551 (Japan)
E-mail: niwasawa@chem.titech.ac.jp

Supporting information for this article can be found under:
<http://dx.doi.org/10.1002/anie.201604371>.

4a.^[14] Formation of bezaldehyde was noted in these reactions. Among several $\text{ReX}(\text{CO})_5$, $\text{ReI}(\text{CO})_5$ gave the best result (entries 1–4), and even 2.5 mol % of $\text{ReI}(\text{CO})_5$ afforded the cycloheptadiene **3a** in 97% yield (entry 5). Even an equimolar amount of propargyl ether **1a** and siloxydiene **2a** gave the desired product **3a** in good yield by carrying out the reaction in the presence of activated 4 Å M.S. (entries 6 and 7).^[15]

Generality of the reaction with various 2-siloxydienes **2** was examined using propargyl benzyl ether **1a** under the optimized conditions (Table 2). Various 4-aryl substituted 2-

Table 2: Examination of various siloxydienes.

Entry	R ¹	R ²	Yield [%]
1	H		2a 3a 90
2	H	Ph	2b 3b 86
3	H	<i>p</i> -MeOC ₆ H ₄	2c 3c 88
4	H	<i>p</i> -BrC ₆ H ₄	2d 3d 86
5	H	2,5-(MeO) ₂ C ₆ H ₃	2e 3e 92
6	H	2-furyl	2f 3f 63
7	H		2g 3g 88
8	H		2h 3h 82
9	H	<i>n</i> Pr	2i 3i 77
10	H	<i>i</i> Pr	2j 3j 85
11	H		2k 3k 93
12 ^[a]	C ₂ H ₄ OBOM ^[b]	H	2l 3l 72
13 ^[c]			2m 3m 91

[a] The reaction was carried out for 12 h. [b] BOM = benzyloxymethyl. [c] The reaction was carried out for 8 h. [d] That is, R¹, R² = (CH₂)₄.

siloxy-1,3-dienes such as electron-donating and electron-withdrawing phenyl, furyl, and indolyl groups underwent the reaction smoothly to give the products in high yield (entries 1–7). 4-Alkenyl and 4-alkyl-2-siloxy-1,3-dienes also gave the corresponding cycloheptadienes **3** in good yield (entries 8–11). Even 3-alkyl- and 3,4-dialkyl-substituted 2-siloxy-1,3-dienes were also employable without problem (entries 12 and 13). All these reactions proceeded by heating just an equimolar mixture of **1a** and **2** to give synthetically useful cycloheptadiene derivatives **3** in good yield.

The result of the examination of the generality of the propargyl ethers is shown in Table 3. Various ethers such as alkyl and THP ethers were also employable to give the product **3a** in good yield (entries 1 and 2). A propargyl ether containing cyclohexane moiety gave a spirocyclic product in good yield (entry 3). Functional groups such as carbamate,


Table 3: Examination of various propargyl ethers.

Entry	Propargyl ethers	Yield [%]
1 ^[a]		1b 3a 90
2 ^[b]		1c 3a 59
3		5a 6a 80
4 ^[b]		5b 6b 65
5 ^[c]		5c 6c 87
6 ^[b]		5d 6d 69
7 ^[c]		5e 6e 75
8 ^[d]		5f 6f (cis/trans 79:21) ^[e]
9 ^[d,f]		5g 6g 69 (cis/trans 77:23) ^[e]
		5g 6g 58 ^[g]


[a] For 6 h. [b] For 12 h. [c] For 8 h. [d] 5 mol % Re catalyst was used. [e] The ratio was determined by ¹H NMR spectroscopy. [f] 1.5 equiv of **2a** was used. [g] Obtained as a mixture with **7g**. The yield was determined by ¹H NMR spectroscopy.

acetal, and silyl ether were compatible under the reaction conditions, and desired products were obtained without problem (entries 4–6). Secondary propargyl ether gave the corresponding product with reasonable diastereoselectivity (entries 7 and 8). More importantly, a simple propargyl ether itself gave the corresponding product in reasonable yield (entry 9). In this case, formation of methylene cyclohexane derivative **7g** in 17% yield was also noted.^[16]

To obtain information on the mechanism of the reaction, several control experiments were carried out. First, the reaction of **5c** with silyl enol ether **8** was examined under the standard reaction conditions, and cyclopropane **9** was obtained in good yield, confirming generation of the α,β-unsaturated carbene complex intermediate (Scheme 3). Next, a deuterium labeling experiment was carried out (Scheme 4). When propargyl ether [D]-**1a** with the hydrogen atom of the terminal alkyne deuterated was employed as substrate, the deuterium was introduced onto C-5 position of the product [D]-**3a**. This result clearly showed that the reaction proceeded through the vinylidene intermediate in which terminal deuterium transferred to the internal position. Based on



Scheme 3: Cyclopropanation of silyl enol ether.



Scheme 4. Deuterium labeling experiments.

these results, the total mechanism of the reaction was shown in Scheme 4. Thus, one or two of the carbonyl ligands of $\text{ReX}(\text{CO})_5$ are liberated by heating and vinylidene intermediate **D** is generated by the reaction with propargyl ether.^[17] Then 1,5-hydride transfer of the benzylic hydrogen onto the vinylidene carbon followed by elimination of benzaldehyde generates α,β -unsaturated carbene complex intermediate **E**.^[18] This intermediate reacts with siloxydiene to give divinylcyclopropane **F**, which undergoes Cope rearrangement to give the product.^[19]

In conclusion, a very concise and practical method for generation of simple α,β -unsaturated carbene complex intermediates is realized by the reaction of easily available propargyl ethers with a catalytic amount of $\text{ReI}(\text{CO})_5$. Further intermolecular addition reaction with siloxydienes efficiently gives synthetically useful silyloxycycloheptadienes in high yields.

Experimental Section

General procedure: 4 Å M.S. (150 mg) were placed in a test tube and was activated by heating at 230 °C for 8 h under reduced pressure (ca. 0.30 mm Hg). The test tube was cooled to room temperature under reduced pressure. $\text{ReI}(\text{CO})_5$ (1.7 mg, 3.8 μmol , 2.5 mol%), propargyl ether (0.15 mmol, 1.0 equiv.), and siloxydiene (0.15 mmol, 1.0 equiv.) were dissolved in 1,4-dioxane (1.5 mL) under argon atmosphere and the resulting mixture was transferred to the test tube. The test tube was closed and the mixture was stirred at 100 °C for 4 h. The reaction was quenched by addition of TMEDA (0.2 mL). After filtration to remove the 4 Å M.S., the filtrate was concentrated under reduced pressure in vacuo. The crude product was purified by preparative TLC to give the [4+3] addition product.

Acknowledgements

This research was supported by a Grant-in-Aid for Scientific Research on Innovative Areas “Precisely Designed Catalysts

with Customized Scaffolding” (No. 15H05800) from the Ministry of Education, Culture, Sports, Science, and Technology of Japan. H.S. thanks JSPS for a fellowship.


Keywords: cycloaddition reactions · propargyl ether · rhenium · unsaturated carbene complex · vinylidene

How to cite: *Angew. Chem. Int. Ed.* **2016**, 55, 10057–10060

Angew. Chem. **2016**, 128, 10211–10214

- [1] For recent reviews of generation of α,β -unsaturated carbene complexes by transition-metal-catalyzed 1,2-acyloxy migration, see: a) K. Miki, S. Uemura, K. Ohe, *Chem. Lett.* **2005**, 34, 1068–1073; b) N. Marion, G. Lemi re, A. Correa, C. Costabile, R. S. Ram n, X. Moreau, P. Fr mont, R. Dahmane, A. Hours, D. Lesage, J.-C. Tabet, J.-P. Goddard, V. Gandon, L. Cavallo, L. Fensterbank, M. Malacria, S. P. Nolan, *Chem. Eur. J.* **2009**, 15, 3243–3260; c) X.-Z. Shu, D. Shu, C. M. Schienebeck, W. Tang, *Chem. Soc. Rev.* **2012**, 41, 7698–7711; d) L. Fensterbank, M. Malacria, *Acc. Chem. Res.* **2014**, 47, 953–965.
- [2] For other examples for generation of α,β -unsaturated carbene complexes based on electrophilic activation of alkynes using metal catalysts, see: a) L. Peng, X. Zhang, S. Zhang, J. Wang, *J. Org. Chem.* **2007**, 72, 1192–1197; b) G. Zhang, L. Zhang, *J. Am. Chem. Soc.* **2008**, 130, 12598–12599; c) R. Sanz, D. Miguel, M. Gohain, P. Garc a-Garc a, M. A. Fern ndez-Rodr guez, A. Gonz lez-P rez, O. Nieto-Faza,  . R. de Lera, F. Rodr guez, *Chem. Eur. J.* **2010**, 16, 9818–9828; d) K. Saito, H. Sogou, T. Suga, H. Kusama, N. Iwasawa, *J. Am. Chem. Soc.* **2011**, 133, 689–691; e) P. A. Allegretti, E. M. Ferreira, *Org. Lett.* **2011**, 13, 5924–5927; f) P. A. Allegretti, E. M. Ferreira, *Chem. Sci.* **2013**, 4, 1053–1058; g) Y. Kwon, I. Kim, S. Kim, *Org. Lett.* **2014**, 16, 4936–4939; h) X. Li, H. Li, W. Song, P.-S. Tseng, L. Liu, I. A. Guzei, W. Tang, *Angew. Chem. Int. Ed.* **2015**, 54, 12905–12908; *Angew. Chem.* **2015**, 127, 13097–13100.
- [3] For representative examples of cyclopropanation reaction by using unsaturated carbene complexes, see Ref. [1] and the following: a) D. Qian, J. Zhang, *Chem. Soc. Rev.* **2015**, 44, 677–698; b) V. Rautenstrauch, *J. Org. Chem.* **1984**, 49, 950–952; c) E. Mainetti, V. Mour  s, L. Fensterbank, M. Malacria, J. Marco-Contelles, *Angew. Chem. Int. Ed.* **2002**, 41, 2132–2135; *Angew. Chem.* **2002**, 114, 2236–2239; d) Y.-W. Sun, X.-Y. Tang, M. Shi, *Chem. Commun.* **2015**, 51, 13937–13940.
- [4] For representative examples of other cycloaddition reactions of unsaturated carbene complexes generated from alkyne substrates using electrophilic metal catalysts, see: a) N. D. Shapiro, F. D. Toste, *J. Am. Chem. Soc.* **2008**, 130, 9244–9245; b) N. D. Shapiro, Y. Shi, F. D. Toste, *J. Am. Chem. Soc.* **2009**, 131, 11654–11655; c) H. Zheng, J. Zheng, B. Yu, Q. Chen, X. Wang, Y. He, Z. Yang, X. She, *J. Am. Chem. Soc.* **2010**, 132, 1788–1789; d) V. V. Pagar, A. M. Jadhav, R.-S. Liu, *J. Am. Chem. Soc.* **2011**, 133, 20728–20731; e) D. Garayalde, K. Kr ger, C. Nevado, *Angew. Chem. Int. Ed.* **2011**, 50, 911–915; *Angew. Chem.* **2011**, 123, 941–945; f) R. C. Conyers, B. W. Gung, *Chem. Eur. J.* **2013**, 19, 654–664; g) N. Iqbal, C. A. Sperger, A. Fiksdahl, *Eur. J. Org. Chem.* **2013**, 907–914; h) C. Zhao, X. Xie, S. Duan, H. Li, R. Fang, X. She, *Angew. Chem. Int. Ed.* **2014**, 53, 10789–10793; *Angew. Chem.* **2014**, 126, 10965–10969; i) M. E. Muratore, A. Homs, C. Obradors, A. M. Echavarren, *Chem. Asian J.* **2014**, 9, 3066–3082; j) D. Shu, W. Song, X. Li, W. Tang, *Angew. Chem. Int. Ed.* **2013**, 52, 3237–3240; *Angew. Chem.* **2013**, 125, 3319–3322; k) H. Kusama, H. Sogo, K. Saito, T. Suga, N. Iwasawa, *Synlett* **2013**, 24, 1364–1370; l) W. Yang, T. Wang, Y. Yu, S. Shi, T. Zhang, A. S. K. Hashmi, *Adv. Synth. Catal.* **2013**, 355, 1523–1528.

- [5] For recent reviews on hydride transfer in organic synthesis, see: a) B. Peng, N. Maulide, *Chem. Eur. J.* **2013**, *19*, 13274–13287; b) M. C. Haibach, D. Seidel, *Angew. Chem. Int. Ed.* **2014**, *53*, 5010–5036; *Angew. Chem.* **2014**, *126*, 5110–5137; c) L. Wang, J. Xiao, *Adv. Synth. Catal.* **2014**, *356*, 1137–1171.
- [6] Gagosz et al. reported gold-catalyzed 1,5-hydride transfer of propargyl ethers to give allenes: B. Bolte, Y. Odabachian, F. Gagosz, *J. Am. Chem. Soc.* **2010**, *132*, 7294–7296.
- [7] For representative examples of reactions utilizing hydride transfer onto π -alkyne complexes of other substrates, see: a) D. Shikanai, H. Murase, T. Hata, H. Urabe, *J. Am. Chem. Soc.* **2009**, *131*, 3166–3167; b) I. D. Jurberg, Y. Odabachian, F. Gagosz, *J. Am. Chem. Soc.* **2010**, *132*, 3543–3552; c) J. Kuang, S. Ma, *J. Am. Chem. Soc.* **2010**, *132*, 1786–1787; d) S. Bhunia, S. Ghorpade, D. B. Huple, R.-S. Liu, *Angew. Chem. Int. Ed.* **2012**, *51*, 2939–2942; *Angew. Chem.* **2012**, *124*, 2993–2996; e) T. Sugiishi, H. Nakamura, *J. Am. Chem. Soc.* **2012**, *134*, 2504–2507; f) X. Wu, S.-S. Chen, Y. Hu, L.-Z. Gong, *Org. Lett.* **2014**, *16*, 3820–3823, and references therein.
- [8] For examples of reactions utilizing alkoxy migration of propargyl ethers, see: a) E. Jiménez-Núñez, M. Raducan, T. Lauterbach, K. Molawi, C. R. Solorio, A. M. Echavarren, *Angew. Chem. Int. Ed.* **2009**, *48*, 6152–6155; *Angew. Chem.* **2009**, *121*, 6268–6271; b) J. Carreras, M. Livendahl, P. R. McGonigal, A. M. Echavarren, *Angew. Chem. Int. Ed.* **2014**, *53*, 4896–4899; *Angew. Chem.* **2014**, *126*, 4996–4999.
- [9] For a rare example of 1,4-hydride transfer onto electrophilically activated alkynes, see: S. Yang, Z. Li, X. Jian, C. He, *Angew. Chem. Int. Ed.* **2009**, *48*, 3999–4001; *Angew. Chem.* **2009**, *121*, 4059–4061.
- [10] For reviews of vinylidene complexes, see: a) M. I. Bruce, *Chem. Rev.* **1991**, *91*, 197–257; b) C. Bruneau, P. H. Dixneuf, *Acc. Chem. Res.* **1999**, *32*, 311–323; c) “Preparation and stoichiometric reactivity of mononuclear metal vinylidene complexes”: M. I. Bruce in *Metal Vinylidenes and Allenylidenes in Catalysis* (Eds.: C. Bruneau, P.-H. Dixneuf), Wiley-VCH, Weinheim, **2008**; d) B. M. Trost, A. McClory, *Chem. Asian J.* **2008**, *3*, 164–194; e) J. M. Lynam, *Chem. Eur. J.* **2010**, *16*, 8238–8247.
- [11] To our knowledge, there is no report on catalytic reactions incorporating this 1,5-hydride transfer of vinylidene intermediates derived from propargyl ethers. a) J. Ipaktschi, J. Mohsseni-Ala, A. Dülmer, C. Loschen, G. Frenking, *Organometallics* **2005**, *24*, 977–989; b) R. Castarlenas, M. Eckert, P. H. Dixneuf, *Angew. Chem. Int. Ed.* **2005**, *44*, 2576–2579; *Angew. Chem.* **2005**, *117*, 2632–2635; c) T. Bolaño, R. Castarlenas, M. A. Esteruelas, E. Oñate, *J. Am. Chem. Soc.* **2007**, *129*, 8850–8859; d) C.-P. Chung, C.-C. Chen, Y.-C. Lin, Y.-H. Liu, Y. Wang, *J. Am. Chem. Soc.* **2009**, *131*, 18366–18375; e) I. Hyder, M. Jiménez-Tenorio, M. C. Puerta, P. Valerga, *Organometallics* **2011**, *30*, 726–737.
- [12] For representative examples of reactions utilizing hydride transfer onto vinylidene complexes derived from other substrates, see: a) G. B. Bajracharya, N. K. Pahadi, I. D. Gridnev, Y. Yamamoto, *J. Org. Chem.* **2006**, *71*, 6204–6210; b) A. Odedra, S. Datta, R.-S. Liu, *J. Org. Chem.* **2007**, *72*, 3289–3292; c) M. Tobisu, H. Nakai, N. Chatani, *J. Org. Chem.* **2009**, *74*, 5471–5475; for an example of hydride transfer onto carbene complexes, see: d) F. Cambeiro, S. López, J. A. Varela, C. Saá, *Angew. Chem. Int. Ed.* **2012**, *51*, 723–727; *Angew. Chem.* **2012**, *124*, 747–751.
- [13] For preparation of dienes from propargyl ethers utilizing metal-catalyzed hydride transfer, see: K.-L. Yeh, B. Liu, C.-Y. Lo, H.-L. Huang, R.-S. Liu, *J. Am. Chem. Soc.* **2002**, *124*, 6510–6511.
- [14] Cycloheptadiene **3a** easily isomerized to isomer **4a** by acid catalysis.
- [15] Activation of 4 Å M.S. was carried out by heating at 230 °C for 8 h under reduced pressure (ca. 0.30 mm Hg⁻¹).
- [16] Methylene cyclohexane derivative **7g** was produced via formal [4+2] cycloaddition reaction of unsaturated carbene complex intermediate with siloxydiene. For a similar reaction, see: W. D. Wulff, W. E. Bauta, R. W. Kaesler, P. J. Lankford, R. A. Miller, C. K. Murray, D. C. Yang, *J. Am. Chem. Soc.* **1990**, *112*, 3642–3659.



- [17] For recent examples of rhenium vinylidene complexes, see: a) Y. Kuninobu, K. Takai, *Chem. Rev.* **2011**, *111*, 1938–1953; b) D. R. Senn, A. Wong, A. T. Patton, M. Marsi, C. E. Strouse, J. A. Gladysz, *J. Am. Chem. Soc.* **1988**, *110*, 6096–6109; c) C. Kelley, L. A. Mercado, M. R. Terry, N. Lugan, G. L. Geoffroy, Z. Xu, A. L. Rheingold, *Angew. Chem. Int. Ed. Engl.* **1992**, *31*, 1053–1055; *Angew. Chem.* **1992**, *104*, 1066–1068; d) C. Bianchini, N. Mantovani, A. Marchi, L. Marvelli, D. Masi, M. Peruzzini, R. Rossi, A. Romerosa, *Organometallics* **1999**, *18*, 4501–4508; e) D. A. Valyaev, O. V. Semeikin, M. G. Peterleitner, Y. A. Borisov, V. N. Khrustalev, A. M. Mazhuga, E. V. Kremer, N. A. Ustynyuk, *J. Organomet. Chem.* **2004**, *689*, 3837–3846; f) J. García-Álvarez, J. Díez, J. Gimeno, C. M. Seifried, C. Vidal, *Inorg. Chem.* **2013**, *52*, 5428–5437; g) D. Xia, Y. Wang, Z. Du, Q.-Y. Zheng, C. Wang, *Org. Lett.* **2012**, *14*, 588–591; h) S. Hori, M. Murai, K. Takai, *J. Am. Chem. Soc.* **2015**, *137*, 1452–1457.
- [18] Generation of rhenium vinylidene complexes by the reaction of alkynylrhenium species with imines, followed by hydride transfer to give allylamine derivatives, was reported: Y. Fukumoto, M. Daijo, N. Chatani, *J. Am. Chem. Soc.* **2012**, *134*, 8762–8765.
- [19] For review of divinylcyclopropane rearrangement, see: a) P. Tang, Y. Qin, *Synthesis* **2012**, *44*, 2969–2984; b) S. Krüger, T. Gaich, *Beilstein J. Org. Chem.* **2014**, *10*, 163–193; c) A. C. Jones, J. A. May, R. Sarpong, B. M. Stoltz, *Angew. Chem. Int. Ed.* **2014**, *53*, 2556–2591; *Angew. Chem.* **2014**, *126*, 2590–2628; divinylcyclopropane rearrangement could proceed with analogous *trans* divinylcyclopropane by heating at 90 °C; see: d) W. D. Wulff, D. C. Yang, C. K. Murray, *J. Am. Chem. Soc.* **1988**, *110*, 2653–2655.

Received: May 5, 2016

Published online: July 8, 2016