

J. J. Schneider, U. Denninger, J. Hagen, C. Krüger, D. Bläser, R. Boese, *Chem. Ber.* **1997**, *130*, 1433.

[16] J. Weiss, T. Priermeier, R. A. Fischer, *Inorg. Chem.* **1996**, *35*, 71.

[17] J. Weiss, A. Frank, E. Herdtweck, S. Nlate, M. Mattner, R. A. Fischer, *Chem. Ber.* **1996**, *129*, 297.


[18] R. A. Fischer, M. Kleine, O. Lehmann, M. Stuke, *Chem. Mat.* **1995**, *7*, 1863.

Selective Activation of Ar–Cl and Ar–C Bonds with Iron(II) Complexes**

Géraldine Poignant, Sourisak Sinbandhit, Loïc Toupet, and Véronique Guerschais*

Selective bond-activation reactions mediated by transition metals are currently the focus of synthetic and mechanistic interest. These fundamental processes are particularly important in carbene and arene chemistry.^[1, 2] In this context, the electrophilic arylcarbene chelate complexes **1** (X = Cl) and **2** (X = OMe) are good candidates for promoting new reactions within the coordination sphere, since the chelating group X can be activated by the Lewis acidic organo-iron fragment^[3, 4] or dissociate to provide a vacant coordination site.^[5, 6] Here we report on the reactivity of **1** and **2**^[5] towards alkoxides. The outcome of the reaction depends on the nature of the chelate group; selective activation of Ar–Cl and Ar–C bonds was achieved.


Complex **1** reacts cleanly with alkoxides RONA to give the unexpected neutral chelate complexes **3a** (R = Me) and **3b** (R = Et), which were isolated as stable black crystals in 83–89% yield from pentane (Scheme 1). Both contain two additional alkoxy groups. The ¹H NMR spectrum (C₆D₆, 25 °C) of **3a** exhibits three broad signals for the methoxy substituents, one of which corresponds to the coordinated OMe group, while the other two methoxy groups are diastereotopic. The coalescence of these signals on warming (T_C (300 MHz) = 42 °C) to give a singlet at δ = 2.90, indicates free exchange on the NMR time scale. For **3b**, the ¹H and ¹³C NMR spectra each exhibit two well-resolved pairs of signals



Scheme 1. a) RONA (3 equiv)/ROH, THF, –80 °C → room temperature. TfO = trifluoromethanesulfonate.


for the methyl and methylene groups, which indicates the presence of two different ethoxy groups. Moreover, in the ¹H NMR spectrum (25 °C) one of the methylene groups gives rise to an AB system at δ = 3.00 and 2.78 (²J_{HH} = 10 Hz) when the corresponding methyl signal at δ = 0.98 is selectively ¹H-decoupled.

As expected the OMe ligand is labile, and treatment of **3a** with PMe₃ quantitatively yields orange crystals of **4**. In this



case, the three OMe groups are magnetically equivalent in the ¹H (δ = 3.29) and ¹³C (δ = 49.9) NMR spectra (25 °C). An X-ray structure analysis confirms the proposed structure (Figure 1).^[7] The FeC≡O unit deviates from the expected linear geometry (Fe–C–O 171.0(3)°). The bond angles at C_{ipso} of the aryl group are quite different owing to the presence of the bulky tris(methoxy)methyl substituent.

The formation of complexes **3a–b** involves cleavage of the Ar–Cl bond. This cleavage is promoted by coordination of the chlorine atom, as shown by the following experiment. Treatment of the nonchelated complex **5**⁺OTf[–]^[5] under the same conditions led to formation of the acetal complexes **6a** (R = Me) and **6b** (R = Et) by addition of RO[–] to the electrophilic carbene carbon atom. Although stable in the



solid state as a yellow powder, these species undergo thermal decomposition in solution at 0 °C. Addition of HBF₄ · OEt₂ to a crude solution of **6a** led quantitatively to the carbene

[*] Dr. V. Guerschais, G. Poignant
UMR 6509 CNRS-Université de Rennes 1
"Organométalliques et Catalyse: Chimie et Electrochimie Moléculaires"
Beaulieu, Université de Rennes 1
F-35042 Rennes Cedex (France)
Fax: (+33) 299-281646
E-mail: guerschais@univ-rennes1.fr

Dr. S. Sinbandhit
Centre Régional de Mesures Physiques de l'Ouest (C.R.M.P.O.)
Beaulieu, Université de Rennes 1
F-35042 Rennes Cedex (France)
Dr. L. Toupet
UMR 6626 Groupe Matière Condensée et Matériaux
Beaulieu, Université de Rennes 1
F-35042 Rennes Cedex (France)

[**] This work was supported by the CNRS and HCM program (contract number 940501). We thank the "Région Bretagne" for a grant to G.P. and Dr. P. Guénot for mass spectrometric measurements.





Figure 1. Crystal structure of **4**. Selected bond lengths [Å] and angles [°]: C17–C18 1.519(6), O2–C18 1.382(4), O3–C18 1.411(4), O4–C18 1.413(5), Fe–C12 2.033(3), Fe–P 2.187(1), Fe–C11 1.699(4), O2–C19 1.426(6), O3–C20 1.420(4), O4–C21 1.421(5); Fe–C11–O1 171.0(3), C11–Fe–C12 99.6, P–Fe–C11 85.9(1), P–Fe–C12 89.2(1), Fe–C12–C17 133.3(3), Fe–C12–C13 112.6(2).

complex $5^+BF_4^-$, which still contains the chlorine atom, as confirmed by mass spectrometry.

Activation of an Ar–F bond by an iron complex with formation of an Ar–Fe bond has been previously postulated, but no spectroscopic evidence for an Fe–F interaction was obtained.^[8] Halohydrocarbon complexes $[M(\eta^1-XR)]$ (X = halogen; R = alkyl, aryl) undergo nucleophilic attack at the C_α atom.^[3] However, similar activation is not observed for the chlorobenzene complex **7** on addition of Lewis bases.^[9] In our



case, the intermediate formation of an Ar–OMe bond by nucleophilic substitution^[10] was ruled out by the investigation of the anisyl derivative **2**. The formation of the Fe–aryl bond and the subsequent rearrangement processes are not yet understood.

In contrast, the reaction of **2** with EtONa takes a completely different course: the new carbene–aryl complex **8** was isolated as a yellow solid in 80% yield (Scheme 1). The presence of the carbene ligand is indicated by the downfield signal at $\delta = 263.8$ in the ^{13}C NMR spectrum. The crystal structure (Figure 2) clearly shows that the geometry of the carbene ligand is distorted. There are two different Fe–C–O bond angles due to the presence of the anisyl ligand.^[7]

We assume that the ethoxide ion initially adds to the carbene center and that the intermediate complex **9** then





Figure 2. Crystal structure of **8**. Selected bond lengths [Å] and angles [°]: Fe–C17 1.990(6), Fe–C12 1.881(5), O2–C12 1.305(7), O3–C12 1.335(7), O4–C23 1.423(7); Fe–C11–O1 173.2(6), O2–C12–O3 108.8(4), Fe–C12–O2 133.4(4), Fe–C12–O3 117.8(4).



rearranges by Ar– C_α bond cleavage, that is, α -aryl elimination. Examples of hydride, alkyl, and aryl migration to a carbene ligand are known,^[11, 12] the reverse reaction is probably induced by the presence of the labile ligand^[13] and/or the instability of the acetal complex (see above). Exchange of the alkyl group of the carbene ligand occurs in the presence of an excess of EtO[−]/EtOH, a behavior already observed for vinyl ether complexes.^[14]

This study illustrates the dichotomy of reactivity of iron complexes: exclusive Ar–Cl bond cleavage is promoted by the Lewis acid–base interaction, whereas the presence of a labile chelate ligand activates the Ar–C bond.

Experimental Section

A solution of **1**, **2**, or **5** (1 mmol) in THF was treated at $-80^\circ C$ with a freshly prepared solution of RONA (3 equiv) in ROH (R = Me, Et). After stirring for 1 h, the solution was evaporated to dryness. The products were extracted with pentane and crystallized.

3a (89% yield, dark brown crystals): 1H NMR (300 MHz, C_6D_6): $\delta = 7.84$ (dd, $^3J(H,H) = 7.5$, $^4J(H,H) = 0.8$ Hz, 1H, Ar), 7.35 (td, $^3J(H,H) = 7.4$, $^4J(H,H) = 1.6$ Hz, 1H, Ar), 7.10 (td, $^3J(H,H) = 7.4$, $^4J(H,H) = 1$ Hz, 1H, Ar), 6.99 (dd, $^3J(H,H) = 7.5$, $^4J(H,H) = 1.5$ Hz, 1H, Ar), 3.02 (brs, 3H, OMe), 2.88 (brs, 3H, OMe), 2.73 (brs, 3H, OMe), 1.42 (s, 15H, C_5Me_5); $^{13}C\{^1H\}$ NMR (50.3 MHz, $CDCl_3$): $\delta = 221.7$ (CO), 174.6 (FeAr), 140.2 (Ar), 136.5 (Ar_C), 127.9 (Ar), 126.8 ($C(OMe)_3$), 123.9 (Ar), 121.9 (Ar), 89.6 (C_5Me_5), 56.3 (OMe), 52.7 (OMe), 52.5 (OMe), 10.8 (C_5Me_5); IR (pentane): $\tilde{\nu} = 1917$ cm^{-1} (CO); C,H analysis calcd for $C_{21}H_{28}O_4Fe$: C 63.01, H 7.05; found: C 62.78, H 7.12.

3b (83% yield, dark brown crystals): 1H NMR (300 MHz, $CDCl_3$): $\delta = 7.57$ (d, $^3J(H,H) = 7.5$ Hz, 1H, Ar), 7.17 (t, $^3J(H,H) = 6.7$ Hz, 1H, Ar), 6.96 (t, $^3J(H,H) = 7$ Hz, 1H, Ar), 6.84 (d, $^3J(H,H) = 7$ Hz, 1H, Ar), 3.31 (q, $^3J(H,H) = 7$ Hz, 2H, OCH_2CH_3), 3.15 (s, 3H, OMe), 3.00 (m, 1H, OCH_2CH_3), 2.79 (m, 1H, OCH_2CH_3), 1.60 (s, 15H, C_5Me_5), 1.14 (t,

$^3J(\text{H,H}) = 7 \text{ Hz}$, 3H , OCH_2CH_3), $0.98 \text{ (t, } ^3J(\text{H,H}) = 7 \text{ Hz, } 3\text{H, OCH}_2\text{CH}_3)$; $^{13}\text{C}\{^1\text{H}\}$ NMR (75.47 MHz, CDCl_3): $\delta = 221.4 \text{ (CO)}$, 173.5 (FeAr) , 139.7 (Ar) , 137.4 (Ar) , 127.4 (Ar) , $125.8 \text{ (C(OMe)(OEt))}_2$, 123.4 (Ar) , 121.5 (Ar) , $89.2 \text{ (C}_5\text{Me}_5)$, $60.2 \text{ (OCH}_2\text{CH}_3)$, $59.8 \text{ (OCH}_2\text{CH}_3)$, 55.8 (OMe) , $15.3 \text{ (OCH}_2\text{CH}_3)$, $14.8 \text{ (OCH}_2\text{CH}_3)$, $10.1 \text{ (C}_5\text{Me}_5)$; IR (pentane): $\tilde{\nu} = 1914 \text{ cm}^{-1}$ (CO); C,H analysis calcd for $\text{C}_{23}\text{H}_{32}\text{O}_4\text{Fe}$: C 64.49, H 7.53; found: C 64.82, H 7.47; HRMS (70 eV): m/z : 428.1641 [M^+], calcd for $\text{C}_{23}\text{H}_{32}\text{O}_4\text{Fe}$: 428.1649.

4 (65% yield): ^1H NMR (200 MHz, CDCl_3): $\delta = 7.84 \text{ (d, } ^3J(\text{H,H}) = 7.6 \text{ Hz, } 1\text{H, Ar)}$, $7.38 \text{ (dd } ^3J(\text{H,H}) = 7.8, ^4J(\text{H,H}) = 1.7 \text{ Hz, } 1\text{H, Ar)}$, $6.85 \text{ (td, } ^3J(\text{H,H}) = 7.3, ^4J(\text{H,H}) = 1.4 \text{ Hz, } 1\text{H, Ar)}$, $6.69 \text{ (td, } ^3J(\text{H,H}) = 7.3, ^4J(\text{H,H}) = 1.8 \text{ Hz, } 1\text{H, Ar)}$, $3.29 \text{ (s, } 9\text{H, OMe)}$, $1.65 \text{ (s, } 15\text{H, C}_5\text{Me}_5)$, $1.12 \text{ (d, } ^2J(\text{P,H}) = 8.4 \text{ Hz, } 9\text{H, PMe}_3)$; $^{31}\text{P}\{^1\text{H}\}$ NMR (81 MHz, CDCl_3): $\delta = 36.51 \text{ (s, PMe}_3)$; $^{13}\text{C}\{^1\text{H}\}$ NMR (75.47 MHz, CDCl_3): $\delta = 222.6 \text{ (d, } ^2J(\text{P,C}) = 38 \text{ Hz, CO)}$, $170.4 \text{ (d, } ^2J(\text{P,C}) = 26 \text{ Hz, FeAr)}$, $146.1 \text{ (d, } ^2J(\text{P,C}) = 12 \text{ Hz, Ar)}$, 145.8 (Ar) , 129.2 (Ar) , 122.6 (Ar) , 119.3 (Ar) , 117.7 (C(OMe))_3 , $92.1 \text{ (C}_5\text{Me}_5)$, 49.9 (OMe) , $17.6 \text{ (d, } ^1J(\text{P,C}) = 26 \text{ Hz, PMe}_3)$, $10.2 \text{ (C}_5\text{Me}_5)$; IR (CH_2Cl_2): $\tilde{\nu} = 1898 \text{ cm}^{-1}$ (CO); C,H analysis calcd for $\text{C}_{24}\text{H}_{37}\text{O}_4\text{FeP}$: C 60.51, H 7.83; found: C 60.78, H 7.75.

6b (80% yield, isolated at -40°C , yellow powder): ^1H NMR (300 MHz, CDCl_3 , -30°C): $\delta = 7.58, 7.47, 7.17, 6.98 \text{ (Ar)}$, $3.55, 3.40 \text{ (brm, OCH}_2\text{CH}_3)$, 3.31 (s, OMe) , $3.25, 3.05 \text{ (brm, OCH}_2\text{CH}_3)$, 2.98 (s, OMe) , $1.73, 1.64 \text{ (s, C}_5\text{Me}_5)$, $1.35, 1.17 \text{ (brm, OCH}_2\text{CH}_3)$; $^{13}\text{C}\{^1\text{H}\}$ NMR (75.47 MHz, CDCl_3 , -30°C): $\delta = 219.1, 218.8, 218.2 \text{ (CO)}$, $149.6, 147.9 \text{ (Ar)}$, $131.3, 130.9 \text{ (Ar)}$, $129.8, 129.2 \text{ (Ar)}$, $128.0, 126.9, 126.5, 125.9, 125.6 \text{ (Ar)}$, $116.5, 114.9 \text{ (C}_a)$, $97.9, 96.6 \text{ (C}_5\text{Me}_5)$, $58.9, 57.1 \text{ (OCH}_2\text{CH}_3)$, $51.5, 50.8 \text{ (OMe)}$, $15.5, 15.0 \text{ (OCH}_2\text{CH}_3)$, $10.0, 9.8 \text{ (C}_5\text{Me}_5)$; IR (pentane): $\tilde{\nu} = 1997 \text{ (CO)}$, 1947 cm^{-1} (CO). Two isomers were observed at -30°C owing to hindered $\text{C}_a\text{--Ar}$ rotation at low temperature. No coalescence was observed up to the decomposition temperature of 0°C .

8 (80% yield, yellow crystals): ^1H NMR (300 MHz, C_6D_6): $\delta = 7.75 \text{ (d, } ^3J(\text{H,H}) = 6.6 \text{ Hz, } 1\text{H, Ar)}$, $7.11 \text{ (t, } ^3J(\text{H,H}) = 7.5 \text{ Hz, } 1\text{H, Ar)}$, $6.96 \text{ (t, } ^3J(\text{H,H}) = 7 \text{ Hz, } 1\text{H, Ar)}$, $6.59 \text{ (d, } ^3J(\text{H,H}) = 6.6 \text{ Hz, } 1\text{H, Ar)}$, $4.05 \text{ (brm, } 2\text{H, OCH}_2\text{CH}_3)$, $3.92 \text{ (brm, } 2\text{H, OCH}_2\text{CH}_3)$, $3.48 \text{ (s, } 3\text{H, OMe)}$, $1.61 \text{ (s, } 15\text{H, C}_5\text{Me}_5)$, $0.92 \text{ (brm, } 6\text{H, OCH}_2\text{CH}_3)$; $^{13}\text{C}\{^1\text{H}\}$ NMR (75.47 MHz, CD_2Cl_2): $\delta = 263.8 \text{ (=C}_a)$, 226.6 (CO) , 166.7 (FeAr) , 154.3 (ArOMe) , 143.9 (Ar) , 122.5 (Ar) , 119.8 (Ar) , 108.2 (Ar) , $95.5 \text{ (C}_5\text{Me}_5)$, $66.7 \text{ (brs, OCH}_2\text{CH}_3)$, 55.1 (OMe) , $14.6 \text{ (OCH}_2\text{CH}_3)$, $9.7 \text{ (C}_5\text{Me}_5)$; IR (pentane): $\tilde{\nu} = 1936 \text{ cm}^{-1}$ (CO); HRMS (70 eV): m/z : 383.1314 [$M^+ - \text{OEt}$], calcd for $\text{C}_{21}\text{H}_{27}\text{O}_3\text{Fe}$: 383.1310; 355.1355 [$M^+ - \text{OEt} - \text{CO}$], calcd for $\text{C}_{20}\text{H}_{27}\text{O}_2\text{Fe}$: 355.1361.

Received: October 21, 1997 [Z 11021 IE]

German version: *Angew. Chem.* **1998**, *110*, 1009–1012

Keywords: carbene complexes • C–C activation • C–Cl activation • iron

- [1] V. V. Grushin, H. Alper, *Chem. Rev.* **1994**, *94*, 1047–1062, and references therein.
- [2] Selective bond activation controlled by the choice of metal complex was recently described: M. E. van der Boom, S.-Y. Liou, Y. Ben-David, A. Vigalok, D. Milstein, *Angew. Chem.* **1997**, *109*, 636–637; *Angew. Chem. Int. Ed. Engl.* **1997**, *36*, 625–626; M. Gandelman, A. Vigalok, L. J. W. Shimon, D. Milstein, *Organometallics* **1997**, *16*, 3981–3986.
- [3] R. J. Kulawiec, R. H. Crabtree, *Coord. Chem. Rev.* **1990**, *99*, 89–115, and references therein.
- [4] M. D. Butts, B. L. Scott, G. J. Kubas, *J. Am. Chem. Soc.* **1996**, *118*, 11831–11843.
- [5] G. Poignant, S. Nlate, V. Guerchais, A. J. Edwards, P. R. Raithby, *Organometallics* **1997**, *16*, 124–132.
- [6] S. Nlate, P. Guénot, S. Sinbandhit, L. Toupet, C. Lapinte, V. Guerchais, *Angew. Chem.* **1994**, *106*, 2294–2296; *Angew. Chem. Int. Ed. Engl.* **1994**, *33*, 2218–2219.
- [7] Crystal structure of **4**: Enraf-Nonius CAD4 diffractometer, MoK_α radiation, $\mu = 6.91 \text{ cm}^{-1}$, $F(000) = 508$, $T = 294 \text{ K}$, triclinic, space group $P\bar{1}$, $a = 9.601(5)$, $b = 10.711(5)$, $c = 13.600(6) \text{ Å}$, $\alpha = 75.55(4)$, $\beta = 84.46(4)$, $\gamma = 67.88(3)^\circ$, $V = 1246(1) \text{ Å}^3$, $Z = 2$, $\rho = 1.270 \text{ g cm}^{-3}$. Of 4402 reflections, 3524 with $I > 2\sigma(I)$ were observed ($\omega/2\theta = 1$, hkl :

$011, -12.12, -16.16$). Lorentzian and polarization corrections (DEFLT 1990), $R = 0.046$, $R_w = 0.047$, $w = 1/\sigma(F_o)^2 = [\sigma^2(I) + (0.04 F_o^2)^2]^{-1/2}$, $S_w = 0.890$ (residual $\Delta\rho < 0.44 \text{ e Å}^{-3}$). Crystal structure of **8**: Enraf-Nonius CAD4 diffractometer, MoK_α radiation, $\mu = 7.086 \text{ cm}^{-1}$, $F(000) = 456$, $T = 294 \text{ K}$, triclinic, space group $P\bar{1}$, $a = 8.773(6)$, $b = 9.185(9)$, $c = 14.572(9) \text{ Å}$, $\alpha = 99.64(5)$, $\beta = 89.89(2)$, $\gamma = 108.34(5)^\circ$, $V = 1097(1) \text{ Å}^3$, $Z = 2$, $\rho = 1.296 \text{ g cm}^{-3}$. Of 4133 reflections, 2364 observed with $I > 4\sigma(I)$ ($\omega/2\theta = 1$, hkl : $010, -10.10, -17.17$), Lorentzian and polarization corrections (DEFLT 1990), $R = 0.056$, $R_w = 0.051$, $w = 1/\sigma(F_o)^2 = [\sigma^2(I) + (0.04 F_o^2)^2]^{-1/2}$, $S_w = 1.13$ (residual $\Delta\rho < 0.46 \text{ e Å}^{-3}$). Crystallographic data (excluding structure factors) for the structures reported in this paper have been deposited with the Cambridge Crystallographic Data Centre as supplementary publication no. CCDC-100931. Copies of the data can be obtained free of charge on application to CCDC, 12 Union Road, Cambridge CB2 1EZ, UK (fax: (+44) 1223-336-033; e-mail: deposit@ccdc.cam.ac.uk).

- [8] M. L. H. Green, J. Haggitt, C. P. Mehnert, *J. Chem. Soc. Chem. Commun.* **1995**, 1853–1854; A. N. Chernega, A. J. Graham, M. L. H. Green, J. Haggitt, J. Lloyd, C. P. Mehnert, N. Metzler, J. Souter, *J. Chem. Soc. Dalton Trans.* **1997**, 2293–2303.
- [9] T.-S. Peng, C. H. Winter, J. A. Gladysz, *Inorg. Chem.* **1994**, *33*, 2534–2542.
- [10] For a review, see L. Balas, D. Jurrhy, L. Latxague, S. Grelier, Y. Morel, M. Hamdani, N. Ardoin, D. Astruc, *Bull. Soc. Chim. Fr.* **1990**, *127*, 401–426.
- [11] T. Braun, O. Gevert, H. Werner, *J. Am. Chem. Soc.* **1995**, *117*, 7291–7292.
- [12] J. Yang, W. M. Jones, J. K. Dixon, N. T. Allison, *J. Am. Chem. Soc.* **1995**, *117*, 9776–9777.
- [13] Y. Strentom, W. M. Jones, *Inorg. Chem.* **1986**, *5*, 178–180; J. Yang, J. Yin, K. A. Abboud, J. W. Jones, *Organometallics* **1994**, *13*, 971–978, and references therein.
- [14] A. Davison, D. L. Reger, *J. Am. Chem. Soc.* **1972**, *94*, 9237–9238. M. Rosenblum, *J. Organomet. Chem.* **1986**, *300*, 191–218.

Synthetic Studies on Ciguatoxin: A Convergent Strategy for Construction of the F–M Ring Framework**

Masayuki Inoue, Makoto Sasaki,* and Kazuo Tachibana*

Ciguatoxin (CTX1B, **1**) and its congeners, naturally occurring polycyclic ethers found in marine unicellular algae, are the principal toxins associated with ciguatera fish poisoning.^[1, 2] These potent neurotoxins reportedly bind to the same sites on voltage-sensitive sodium channels (VSSC) as brevetoxins, another class of structurally related marine toxins.^[3] An important structural characteristic is the fact that the hexahydrooxonin ring (the F ring) in **1** and its congeners

[*] Dr. M. Sasaki, Prof. Dr. K. Tachibana, M. Inoue
Department of Chemistry, School of Science
The University of Tokyo
Hongo, Bunkyo-ku, Tokyo 113-0033 (Japan)
Fax: (+81) 3-5800-6898
E-mail: msasaki@chem.s.u-tokyo.ac.jp
ktachi@chem.s.u-tokyo.ac.jp

[**] This work was supported by the Japanese Ministry of Education, Science, Sports, and Culture (grant-in-aid no. 08245103) and the Japan Society for the Promotion of Science for Young Scientists through a fellowship for M.I. We thank Dr. Keiichi Konoki and Professor Michio Murata from our Department for binding assays and valuable discussions and advice.