


The $\text{Na}^+ - \text{K}^+$ transport system of the cell, the $\text{Na}^+ - \text{K}^+$ pump, consists of a carrier part located in the interior of the membrane and the ATP-binding part on the cytoplasmic side of the membrane. The carrier exists in two conformations: E_1 is the high-affinity Na^+ form, and E_2 is the high-affinity K^+ form.



The carrier part has two gates, energy barriers, which can open or close access to the channel side and to the cytoplasmic side. For each ATP that is hydrolyzed to ADP and inorganic phosphate P_i , three Na^+ ions are pumped out of and two K^+ ions into the cell; 70–85% of the free energy of the ATP hydrolysis is used.

The Identification of the Sodium–Potassium Pump (Nobel Lecture)**

Jens C. Skou

Looking for the answer

*You hunt it,
you catch it,
you fool yourself;
the answer,
is always,
a step ahead.*

J. C. S.

Introduction

The cell membrane separates the cell from the surrounding medium. In 1925 Gorter and Grendel^[1] extracted the lipids from red blood cells, spread them in a monomolecular layer on a water phase, and measured the compressed area. It was about twice the surface area of the extracted red blood cells. They suggested that the cell membrane is a bilayer of lipids with the charged head groups of the phospholipids facing the water phase on the two sides of the membrane, and with the hydrocarbon chains meeting in the middle of the membrane. The thickness of the membrane is about 40 Å, much too small to be seen under a microscope.

The surface tension of invertebrate eggs and other cells is, however, much lower than for a water–lipid interphase. To explain this, Danielli and Davson in 1935^[2] suggested that there is an adsorbed layer of proteins on each side of the lipid bilayer. They also introduced a layer of nonoriented lipids in the middle of the bilayer.

In the cytoplasm there are proteins for which the cell membrane is impermeable. At the cell pH level they carry negative charges, neutralized by potassium, which in the cell is at a concentration of about 150 milligram equivalents per liter (mEq L^{-1}), while outside the cell the K^+ concentration is 4 mEq L^{-1} . The difference in K^+ concentration was explained as being due to a Donnan effect of the proteins, and the fact that the membrane is permeable to K^+ . The Donnan effect gives an osmotic pressure which is higher inside the cell than outside; since the membrane is permeable to water, water will

flow in, and as the lipid bilayer cannot resist a hydrostatic pressure the cell will swell and finally burst.

The higher osmotic pressure inside the cell is opposed by a high concentration of sodium outside (140 mEq L^{-1}), while the concentration inside is low (about $10\text{--}20 \text{ mEq L}^{-1}$). It is a problem to explain how this difference in the Na^+ concentration on the two sides of the membrane can be maintained. There are two possible explanations:

One is that the membrane is impermeable to Na^+ , that it is an equilibrium situation. This view was advocated by Conway, an Irish biochemist from Dublin. Boyle and Conway published in 1941 a paper in which they showed that for muscle fibers soaked in solutions with varying K^+ concentrations, the calculated intracellular K^+ concentrations based on this assumption agreed with the measured.^[3] There were, however, two problems. One was that this agreement did not hold at the normal physiological concentrations of K^+ , only at higher concentrations. The other was the requirement of Na^+ impermeability. But how then to explain that there is sodium in the cell, even if it is at a low concentration. Conway gave no answers to these problems.

The other possibility is that the membrane is permeable to Na^+ , and that there are secretory, energy-dependent processes in the cell which compensate for the steady influx of Na^+ , a steady-state distribution. This view was advocated by R. B. Dean in a paper also published in 1941 entitled “Theories of Electrolyte Equilibrium in Muscle”.^[4] Referring to investigations by L. A. Heppel (1939, 1940), by L. A. Heppel and C. L. A. Schmidt (1938), and by H. B. Steinbach (1940)^[4] on muscle fibers, which had shown that the muscle membrane, contrary to the view held by Conway, is permeable to Na^+ , Dean concluded: “the muscle can actively move potassium and sodium against concentration gradients [...] this requires work. Therefore there must be some sort of a pump possibly located in the fiber membrane, which can pump out sodium or, what is equivalent, pump in potassium.”

In the following decade, helped by the introduction of radioactive isotopes of Na^+ and K^+ ,^[5] it was shown not only from experiments on muscle fibers, but on red blood cells, on nerves, and on frog skin that the membrane is permeable to Na^+ as well as to K^+ (for references see an extensive review by Ussing^[6]). An energy-dependent efflux of Na^+ is therefore necessary.

However, Conway strongly defended his view about the impermeability to sodium, and only reluctantly gave way. He

[*] Prof. Dr. J. C. Skou
Department of Biophysics
University of Århus
Ole Worms Allé 185, DK-8000 Aarhus C (Denmark)
Fax: (+45) 8612-9599

[**] Copyright© The Nobel Foundation 1998. We thank the Nobel Foundation, Stockholm, for permission to print this lecture.

admitted that there may be a certain permeability for Na^+ and thereby a need for a pump, but maintained that low permeability for Na^+ is the main explanation of the concentration gradient. Conway's concern was that it is a waste of energy to have a membrane permeable to Na^+ and then spend energy to pump Na^+ out. Krogh^[7] had in a Croonian lecture in 1946 entitled "The active and passive exchange of inorganic ions through the surfaces of living cells and through living membranes generally" criticized Conway's view about impermeability to sodium. Krogh concluded: "The power of active transport of ions is of a common occurrence both in the vegetable and the animal kingdom and is possibly a general characteristic of the protoplasmic surface membrane". Conway replied the same year in a paper in *Nature*:^[8] "Krogh [...] considers the apparent impermeability to sodium as due to an active extrusion, sodium ions entering the (muscle) fibers as fast, if indeed not faster, than potassium. The following may then be considered: the minimal energy required for extrusion of sodium ions from the normal frog's sartorius if sodium enters as fast as potassium". A calculation showed that the energy requirement was about twice the resting metabolism of the muscle. Ussing^[9] came to the same result, but the explained what the apparent problem was. Only a part of the measured Na^+ flux is due to active transport of Na^+ , the other part is due to an exchange across the membrane of Na^+ from the one side for Na^+ from the other side, a $\text{Na}-\text{Na}$ exchange which is energetically neutral and which gives no net flux of Na^+ across the membrane. Taking this into account, the energy available is more than sufficient. However, Conway's concern was relevant in the sense that with the knowledge at that time about membrane function, an active transport seemed energetically an expensive way to solve the osmotic problem of the cell.

In the 1940s and first half of the 1950s the concept of active transport developed.^[6] It was defined as a transport against an electrochemical gradient.^[10] It was shown that the active efflux of sodium was coupled to an influx of potassium, a pump, and that the substrate for the transport was energy-rich phosphate esters.^[11]

But what was the nature of the pump? With the information available in the beginning of the 1950s it was possible to foresee that the pump is a membrane-bound protein with enzymatic activity, which has ATP as substrate and is activated by Na^+ on the cytoplasmic side and by K^+ on the


extracellular side. But nobody apparently thought that way. A reason may be that the membrane according to the model by Danielli and Davson is a bilayer of lipids with no room for proteins inside the bilayer spanning the bilayer. On the contrary, it was assumed that protein in the bilayer would destabilize the cell membrane. It was assumed that the protein was in the interphase between the lipids and the water on the two sides of the membrane.

The Way to the Sodium Pump^[12, 13]

My scientific interest was the mechanism of action of local anaesthetics. I held a position at the Institute of Physiology at the University of Århus, and was using this problem as subject for a thesis. In 1953 I had finished a series of experiments on the problem, which was published in 1954 as a book in Danish, and accepted by the faculty to be defended for the medical doctors degree.^[14] It was also published as six papers in English.^[15]

I had received my medical degree in the summer of 1944, and started my internship at a hospital in Hjørring in the northern part of the country, six months in the medical ward followed by six months in the surgical ward. I became interested in surgery, and after my internship I continued for another year in the surgical ward. We had no anaesthetists and, to avoid the unpleasant ether narcosis, we used whenever possible spinal and local anaesthesia. From the teaching of pharmacology I knew the Meyer–Overton theory,^[16, 17] which states that there is a correlation between solubility of general anaesthetics in lipids and the anaesthetic potency. General anaesthetics are nonpolar substances, while local anaesthetics are weak bases, which at the physiological pH exist as a mixture of charged and uncharged molecules. I wondered which of the two components is the anaesthetic component, and whether a correlation, similar to that for the general anaesthetics, existed for the local anaesthetics. I decided to use this problem as a subject for a thesis. After the two years at the hospital in Hjørring, I took up a position for a year at the Orthopaedic Hospital in Århus, and after this in 1947 I applied for and received a position at the Institute of Medical Physiology at Århus University.

Århus University was young, founded 19 years earlier. There were Institutes for Anatomy, Biochemistry, and Phys-



Jens C. Skou was born in 1918 in Lemvig, Denmark. He received his medical degree at Copenhagen University in 1944. He got his doctor of medical sciences degree in 1954 at Århus University, where he became Professor of Physiology as well as Chairman of the Institute of Physiology in 1963. In 1977 he was appointed Professor of Biophysics at the same university. He is a member of the Danish Academy of Sciences, and has received numerous awards, among them the 1997 Nobel Prize in Chemistry.

iology as a beginning of a Medical Faculty, which was not completed until 1957. These were the only biological Institutes at the Campus, and the scientific biology milieu was poor, with little or no contact with the outside scientific world. We were three young doctors besides the Professor, Søren L. Ørskov, in the department, each working on our thesis. None of us had any scientific background, but Ørskov was very helpful and patient and let us take the necessary time. We had an intake of 140 medical students a year so the teaching load was heavy. After having passed physiology the students had to continue at the Medical Faculty in Copenhagen for their medical degree.

I used the intact sciatic nerve of frog legs as a test object for measuring the blocking potency of five different local anaesthetics, which are weak bases, and of butanol as a representative of a nonpolar blocking agent. After removing the sheath around the nerve to get easier access to the single nerve fibers, I measured the blocking concentration as a function of time, and from this the minimum blocking concentration at infinite time of exposure could be determined. The concentrations necessary varied from the weakest to the strongest of the local anaesthetics by a factor of 1:920 (with butanol included, 1:13 500). The order of anaesthetic potency and solubility in lipids were the same, but the quantitative correlation was poor, that is, local anaesthetics did not follow the Meyer–Overton rule for general anaesthetics.

I was looking for another test object. As the cell membrane is a bilayer of lipids, I decided to use a monomolecular layer of lipids on a Krebs–Ringer water phase as a model for a water–cell membrane interphase. The inspiration came from reading about Langmuir's work on monomolecular layers of lipids on a water phase in "The Physics and Chemistry of Surfaces" by N. K. Adam, and that Schulmann had applied capillary active drugs in the waterphase beneath the monolayer and observed that they penetrated up into the monolayer.^[18]

In a Langmuir trough the area of the monolayer can be measured as a function of the pressure that the monolayer exerts on a floating barrier which separates the monolayer from the pure water phase without the monolayer. My first experiments were with a monolayer of stearic acid. At a given area per molecule, which also means at a given surface pressure, application of the local anaesthetics to the water phase gave an increase in pressure, indicating that the local anaesthetics penetrated up into the monolayer, and the pressure increased with the concentration. There was a certain correlation between anaesthetic potency and pressure increase, but quantitatively not clear enough. However, with a monolayer of lipids extracted from the sciatic frog nerves there was a reasonably good correlation. The order of the concentrations necessary to increase the pressure followed the order of anaesthetic potency. And the minimum blocking concentration of the five local anaesthetics, which as mentioned varied by a factor of 1:920, gave a pressure increase in the monolayer at a certain area which was of the same order; they varied by a factor of 1:3.2. Also the effect of a change in pH on the local anaesthetic potency correlated reasonably well to the effect of pH on the pressure increase at a given area.

The rising phase of the nerve impulse, the depolarization, is due to a transient increase in permeability of the membrane for Na^+ .^[19] The molecular basis for this was unknown, but it seemed unlikely that it was connected to the lipids. I assumed that the permeability increase was on proteins in the membrane. The monolayer results suggested to me that the effect of the penetration of local anaesthetics into the lipid part of the nerve membrane was a blocking of the conformational change in proteins in the membrane, which gave the increase in permeability to Na^+ .

To test this I wanted to see if pressure in a monolayer could influence the enzymatic activity of a protein in the monolayer, and take this as indication of an effect of pressure on conformation. And if there was an effect, I then wanted to form a monolayer of a mixture of lipids and the enzyme, and test if penetration of local anaesthetics into the monolayer had an effect on the enzymatic activity. For this I needed an enzyme with high activity, and which was related to membrane function. A candidate was acetylcholinesterase, which was then being prepared from electric eel by Professor Nachmansohn at Columbia University in New York. It had the further advantage that it involved a visit New York.

The Professor of Physiology in Copenhagen, Einar Lunds-gaard, was a close friend of David Nachmansohn and introduced me. I had planned to spend August in New York, take a break at the end of August and beginning of September to attend the 19th International Congress of Physiology in Montreal, and then return to New York in September (and not, as mentioned in ref. [12, 13], in July or August). This would fit with my teaching schedule. Nachmansohn would not be in New York until September, as he spent the summer at the Marine Biological Station in Woods Hole; he therefore suggested I should join him there in August. In September he would return to New York. I agreed, although I did not know what to do in Woods Hole; there was no access to electric eel.

Scientists interested in the function of the nervous system came from all over the world to Woods Hole during the summer to use the giant axons from squids as test objects. Coming from a young University with a poor scientific milieu, this was like coming to another planet. The place was bubbling with scientific activity. I realized that science is a serious affair and not just a temporary hobby for young doctors writing a thesis in order to qualify for a clinical career. And also that it is competitive. I listened to lectures, met people whose names I knew from the textbooks and from the literature, and spent time in the laboratories looking on and learning from the experiments.

In between I did some reading, and in a paper written by Nachmansohn it was mentioned that B. Libet^[20] in 1948 had shown that there is an ATP hydrolyzing enzyme in the sheath part of the giant axon from squid: an ATPase. As ATP is the energy source in cells I wondered what the function could be of an ATPase in the membrane of a nerve. As it is situated in the membrane, I assumed that it was a lipoprotein, and this was what I needed for the monolayer experiments. I decided to look for the enzyme when I came home.

I prepared acetylcholinesterase at Columbia University in September from the electric eel. Back in Århus I continued the monolayer experiments. I had no access to giant axons in

Århus, but decided to look for the putative nerve membrane ATPase in crab nerves because the crab nerve, like the giant axon, has no myelin sheath. In 1954 I arranged with a fisherman south of Århus to send me some crabs and started to isolate the sciatic nerve from the legs. The nerves were homogenized, and the membrane pieces isolated by a differential centrifugation.

The experiments showed that the membrane fractions had a low Mg^{2+} -activated ATP hydrolyzing enzyme activity. Addition of Na^+ besides Mg^{2+} gave a slight increase in activity. K^+ had no effect in the presence of Mg^{2+} (Figure 1). However, the activity varied from experiment to experiment.




Figure 1. Effect of Na^+ and K^+ in the presence of Mg^{2+} on the hydrolysis of ATP by membrane fractions isolated from nerves of the shore crab *Carcinus maenas*. The test solution contained 30 mM histidine HCl as buffer (pH 7.2), 3 mM ATP, 6 mM Mg^{2+} , Na^+ (0 \blacktriangle , 3 \triangle , 10 \square , 20 \bullet , 40 mM \circ), and K^+ in concentrations shown. The activity a is given as the amount [mg] of inorganic phosphate (P) hydrolyzed from ATP in 30 min at 36 °C (reprinted with permission from ref. [27]).

Calcium was excluded as the reason for the variations. After having spent November and December trying to find a solution, I gave up and went on my Christmas holidays. I resumed the experiments in June of the following year, but still unable to get reproducible results, went on summer holidays. Returning in August, I made a Na^+ salt and a K^+ salt of ATP and found to my surprise that the activity with the K^+ salt was higher than with the Na^+ salt. This could not be due to a difference in ATP, but to an effect of K^+ which differed from that of Na^+ . But why in this experiment and not in the previous experiments, where K^+ had no effects in the presence of Mg^{2+} ? The answer was that in the experiments with the K^+ salt of ATP there was Na^+ in the medium. In other words the enzyme needed a combined effect of Na^+ and K^+ for activation. I then started a systematic investigation of the combined effect of the two cations.

As seen from Figure 1, K^+ has two effects in the presence of Na^+ . It activates with increasing concentration of Na^+ . The K^+ affinity for the activating effect is high. At higher concentrations the activating effect of K^+ decreases, and the apparent affinity for K^+ for this effect decreases with an increase in the Na^+ concentration. With 3 mM Na^+ it is seen that K^+ not only inhibits its own activation, but also the small activation due to Na^+ . The results suggest that there are two sites on the enzyme, one where Na^+ is necessary for activation and

another where K^+ activates when Na^+ is bound to the former. When present in higher concentrations K^+ competes with Na^+ at the Na^+ site, and by displacing Na^+ from the site decreases the activity.

I now understood the reason for the varying results. With little effect of Na^+ on the activity, and of no effect of K^+ , I had not bothered whether or not there was Na^+ or K^+ in the medium. I got ATP as an insoluble barium salt, which was converted into a soluble Na^+ or K^+ salt, sometimes the one sometimes the other, and in between instead of homogenizing the nerves in glucose I used a 0.58 M KCl solution. It never occurred to me that there could be a combined effect of Na^+ and of K^+ .

The problem then was, what was the physiological function of the enzyme? I was interested in the effect of local anaesthetics on nerve conduction, and my first reaction was that this was the Na^+ channel which opens for the influx of Na^+ leading to the nerve impulse. I soon rejected the idea because the opening of the channel is dependent on voltage, and not on ATP. The other possibility was that it was part of or the sodium pump. As I had little knowledge about the active transport of Na^+ and of K^+ , I started to look into the literature to see what the substrate was for the active transport in nerves. I had limited access to the literature, so there were few papers I read about active transport. The closest I could come was that A. L. Hodgkin and R. D. Keynes^[21] had shown that poisoning giant axons with dinitrophenol, cyanide, or azide decreased the active transport of sodium, suggesting that high-energy phosphate esters are the substrate. And as ATP is a high-energy phosphate ester, I thought it likely that it could be the substrate.

There were two papers I did not read: One was by G. Gardos^[22] from Budapest published in 1954 in a Hungarian journal, which I did not have access to and which was not cited in the paper by Hodgkin and Keynes. In this Gardos showed that ATP supported the active uptake of K^+ in red blood cells. The other was a paper published in 1956 by Hodgkin and Keynes,^[23] in which they reported that injecting ATP in a giant axon poisoned with cyanide gave no dramatic recovery of the active extrusion of sodium. Fortunately, I did not see this last paper until I had sent my paper on the crab nerve enzyme for publication. In 1957 the experiment was repeated by Caldwell and Keynes,^[24] with the result that ATP was the substrate for the active transport.

In a discussion after a paper by R. D. Keynes on „Electrolytes and Nerve Activity“^[25] at an international symposium on Neurochemistry in Århus in 1956, I showed the results with the crab nerve ATPase.^[26] The same year I wrote the paper, and suggested, from the characteristics of the effect of the cations and the fact that ATP was a substrate, that the enzyme was involved in the active transport of Na^+ across the cell membrane. I considered putting the word pump in the title, but found it too provocative, so it became “The Influence of Some Cations on an Adenosine Triphosphatase from Peripheral Nerves”. No wonder that few people noticed that this enzyme had to do with active transport of Na^+ . It was published in 1957.^[27]

With my little knowledge about active transport, I was unaware of the importance of the observation on the crab

nerves. Parallel with the crab nerve experiments I continued the experiments on the monolayer with the acetylcholinesterase. In 1958 I presented a paper at the 4th International Congress of Biochemistry in Vienna on “The Influence of the Degree of Unfolding and the Orientation of the Side Chains on the Activity of a Surface Spread Enzyme”.

There was one important experiment I had not done. I realized this when I met Robert Post at the conference. I knew Robert from Woods Hole. We had spend time in the same laboratory, and I had driven with him and his wife Elizabeth in their car from Woods Hole to the International Congress for Physiology in Montreal. He told me that he had since worked with active transport of Na^+ and K^+ in red blood cells, and had shown that the stoichiometry between the Na^+ transport out of and of K^+ into the cell was 3:2.^[28]

I told him about the $\text{Na}^+ + \text{K}^+$ -activated crab nerve enzyme, and that it seemed to be part of or the sodium pump. His reaction suggested to me that this was more important than surface spread enzymes. “Is it inhibited by ouabain?” he asked. “What is ouabain?” was my reply. He then told me that Schatzmann in Switzerland in 1954 had shown that cardiac glycosides, of which ouabain is the most water-soluble, specifically inhibits the active transport in red blood cells.^[29] When Robert Post came to Århus after the conference I had the answer. The enzyme was inhibited by ouabain, even if the sensitivity of the crab nerve enzyme is much lower than the sensitivity of the transport in red blood cells. It convinced Robert that the enzyme had to do with active transport. I had learned that red blood cells were a classical test object for experiments on active transport, and had started to look for the enzyme in these cells. Robert asked if he could go on with these experiments when he returned to U.S. I had no experience with this test object, and as he had the experience, I agreed and continued the experiments with the crab nerve enzyme^[30] and looked for the enzyme in other tissues.^[31]

In 1959, after I had given my first paper on the crab nerve enzyme at the 21th International Congress for Physiology in Buenos Aires, Professor Hodgkin, who was the great name in neurophysiology and came from the famous Cambridge University, invited me for lunch to hear more about the enzyme. His interest suggested to me that the observation was of a certain importance.

Looking back, it was a very simple experiment to identify the pump. Just break the membrane and thus gain access to the Na^+ site on the inside and the K^+ site on the outside, add some ATP, and test for the combined effect of Na^+ and K^+ . It ought to have been done by someone who worked in the transport field and knew about active transport. I felt like an intruder in a field that was not mine.

There was, however, much luck involved. First, that Nachmansohn had invited me to Woods Hole, where I learned about the giant axons and read about the observation by Libet. Next that from the monolayer experiments I became interested in membrane proteins, especially lipoproteins, and therefore took notice of Libet’s observation. Finally, that I chose crab nerves as a test object. I learned later that after homogenization of most other tissues in order to break the membrane, the membrane pieces form vesicles, which must be opened by treatment with detergent in order to get access to

both sides of the membrane and thereby to see the combined effect of Na^+ and K^+ . This is not the case with the crab nerve membranes.

From then on the Na^+/K^+ -ATPase took me away from the monolayer experiments. I never did the planned experiments on the effect of the local anaesthetics on a monolayer of a mixture of lipids and proteins; and I never used the Na^+/K^+ -ATPase for monolayer experiments. First of all because it was not until 1980 that it became possible to extract the enzyme from a membrane in a pure stable watersoluble form, and secondly, because that was only possible with the use of detergent.^[32] A problem would have been the detergent, but I had lost interest in monolayers.

In 1960 Robert Post published the paper on the red blood cell experiments “Membrane adenosine triphosphatase as a participant in the active transport of sodium and potassium in the human erythrocytes”.^[33] In this he convincingly showed that in the red blood cells there is a $\text{Na}^+ + \text{K}^+$ -activated ATPase also, and that the effect of the cations on the activity correlated with the effect on transport. Robert Post was known in the transport field, and his paper had a better title than mine, so it attracted more attention.

In the following few years many papers were published which showed that the enzyme could be found in many different tissues, and evidence was given for its involvement in active transport.^[34] In 1965 so much evidence was at hand, that in a review paper I could conclude that the enzyme system fulfilled the following requirements for a system responsible for the active transport across the cell membrane:

- 1) It is located in the cell membrane.
- 2) On the cytoplasmic side, it has a higher affinity for Na^+ than for K^+ .
- 3) It has an affinity for K^+ on the extracellular side which is higher than that for Na^+ .
- 4) It has enzymatic activity and catalyzes ATP hydrolysis.
- 5) The rate of ATP hydrolysis depends on cytoplasmic Na^+ as well as on extracellular K^+ .
- 6) It is found in all cells that have coupled active transport of Na^+ and K^+ .
- 7) The effect of Na^+ and of K^+ on transport in intact cells, and on the activity of the isolated enzyme, correlates quantitatively.
- 8) The enzyme is inhibited by cardiac glycosides, and the inhibitory effect on the active fluxes of the cations correlates with the inhibitory effect on the isolated enzyme system.^[34]

The enzyme was named the Na^+ - and K^+ -activated ATPase, or Na^+/K^+ -ATPase.

The Na^+/K^+ -ATPase as an Energy Transducer

Let us return to Conway’s concern about the waste of energy. Conway was right in the sense that even if the cell membrane is permeable to sodium, and therefore an energy-requiring pump is needed, the ground permeability to sodium of the cell membrane must be low, so that the maintenance of the gradient for sodium does not become energetically too costly for the cell. But what is the meaning of having a membrane permeable to sodium and then spend energy, 10–

60% of the cell metabolism, to keep Na^+ out of the cell and K^+ within the cell?

1) As mentioned above it solves the osmotic problem due to the presence of impermeable protein anions in the cytoplasm.

2) Furthermore, the gradients for Na^+ into and K^+ out of the cell sustained by the pump represent an energy source, which is used for the creation of a membrane potential. The cell membrane is more permeable to K^+ than to Na^+ , which means that K^+ flows out of the cell faster than Na^+ into the cell. This leads to a diffusion potential across the membrane, negative on the inside, which slows the K^+ outflux and increases the Na^+ influx until the potential reaches a value of about -70 mV , at which point the rates of the two fluxes are equal; a steady-state situation. The membrane potential is the basis for the function of all excitable tissue. The nerve impulse is a depolarization of the membrane potential due to a transient increase in permeability to Na^+ , with influx of Na^+ followed by a repolarization due to an outflux of K^+ .^[35] This leads to an increase in intracellular Na^+ and a decrease in K^+ , which subsequently must be compensated for by the pump.

3) The energy from the concentration gradient is used for transport of other substances in and out of the cell (Figure 2). In the cell membrane there are a number of protein molecules which act as co- or countertransporters, or sym- and antiporters.^[36] There are cotransporters which use the gradient for Na^+ into the cell to transport glucose^[37] or amino acids^[38] into cells so that they have a higher concentration inside than outside, and a Na^+/Cl^- , a $\text{Na}^+/\text{K}^+/2\text{Cl}^-$, and a K^+/Cl^- cotransporter, which may be involved in volume regulation.^[39] There is a $3\text{Na}^+/\text{Ca}^{2+}$ countertransporter of importance for regulation of the intracellular Ca^{2+} concentration^[40] and a Na^+/H^+ countertransport for regulation of internal pH. In addition, there are other examples.

4) The energy from the concentration gradient is used for transepithelial transport in intestine, kidney, and secretory glands.




Figure 2. Transport across the cell membrane. For an explanation see text (reproduced with permission from ref. [73]).

The $\text{Na}^+ - \text{K}^+$ pump thus acts as an energy transducer that converts the chemical energy from the hydrolysis of ATP into another form of energy, a concentration gradient, which is used for exchange of substances across the cell membrane. This is named a secondary active transport, while the transport which is directly dependent on the energy from a chemical reaction, the active Na^+/K^+ transport, is named a primary active transport. The Na^+/K^+ -ATPase has thus a key function in the exchange of substances across the cell membrane.

A problem was where to place the Na^+/K^+ -ATPase in the membrane. The transport system, which is a protein, must have access to both sides of the membrane. As mentioned previously there is no room for proteins in the bilayer of the Danielli–Davson membrane model. Neither is there in Robertson's unit membrane model from 1959^[41] which replaced it. The unit membrane was a bilayer of lipids with proteins arranged asymmetrically on the two sides of the membrane, but still with no proteins spanning the membrane. It was not until 1972 that a suitable model was introduced by Singer and Nicolson.^[42] In their fluid mosaic membrane model there are globular proteins embedded with their nonpolar parts in the bilayer, and with the polar parts facing the two sides of the bilayer. They can move laterally in the membrane, but do not flip-flop. They form pathways for the transport of hydrophilic substances across the membrane.

Other mammalian ion-transporting pumps were identified in the following years: a Ca^{2+} -ATPase in sarcoplasmic reticulum in muscle responsible for the transport of Ca^{2+} out of the muscle fiber,^[43] a Ca^{2+} -ATPase isolated from red blood cells which transports Ca^{2+} out of the cell,^[44] and a H^+/K^+ -ATPase in the stomach which transports H^+ out of the cells in exchange for K^+ , producing the stomach acid.^[45] They all have in common with the Na^+/K^+ -ATPase that the reaction with ATP involves a phosphorylation, and they are therefore named P-type ATPases. A number of other P-type ATPases have been identified in bacteria and fungi.^[46]

In the years that followed the identification of the sodium–potassium pump, many scientists from many countries took part in the elucidation of the structure of the system and of the reaction steps in the transport process. This cannot be covered in the present lecture, but for those interested I shall refer to an extensive review by I. Glynn in 1985,^[11] to the proceedings from the international conferences on the Na^+/K^+ -ATPase held every third year of (the latest is from 1996),^[47] to recent reviews,^[48–50] and to the recently published book by J. D. Robinson: “Moving Questions. A History of Membrane Transport and Bioenergetics”.^[46]

There is, however, one question I would like to touch on shortly without going into details, namely this: What is our present view on the way the system transports the cations?

A Model for the Transport Reaction

The model in Figure 3 is based on the Albers–Post scheme, a reaction scheme in which the system reacts consecutively with the cations, and in which the reaction with ATP in the




Figure 3. The active transport reaction. For an explanation see text (reproduced with permission from ref. [73]).

presence of Na^+ leads to a phosphorylation and the following reaction with K^+ to a dephosphorylation.^[51–56]

The transport system consists of a carrier part located in the innermost part of the membrane with the ATP-binding part on the cytoplasmic side of the membrane (Figure 3). The carrier is in series with a narrow channel which spans half to two-thirds of the membrane and opens to the extracellular side of the membrane^[57, 58] (in Figure 3 the channel part is shown only as the opening to the outside). The carrier exists in two major different conformations:^[51–56, 59–65] E_1 , which is the high-affinity sodium form, and E_2 , which is the high-affinity potassium form. Each of the two conformations can exist in a phosphorylated^[51–56] and a nonphosphorylated form,^[59–65] and each of the two major conformations have subconformations marked with primes in Figure 3. The carrier part has two gates, energy barriers, which can open and close access to the channel side and to the cytoplasmic side, respectively.

Referring to the model, with the system in the E_1 conformation with the gate to the cytoplasmic side open but that to the channel side closed, three Na^+ ions are bound, two of them as a neutral complex and the third one as an ion. The system is catalytically active. The affinity for ATP is high (apparent K_D is about $0.1 \mu\text{M}$ ^[59, 60]), and with ATP bound the enzyme is phosphorylated^[51–56, 66, 67] to an ADP sensitive form. This means that a high-energy phosphate bond is formed ($\text{E}_1 \sim \text{P}$), which closes the gate to the cytoplasmic side. The three Na^+ ions are occluded inside the carrier part.^[68] The following two steps involve conformational changes and rearrangements of the phosphate bond from a high-energy to a low-energy bond ($\text{E}_1' \sim \text{P}$ to $\text{E}_2 - \text{P}$);^[51–56, 66, 67] deocclusion and release of one Na^+ from the $\text{E}_1' \sim \text{P}$ form^[67] to the extracellular side of the membrane takes place. The transition to the $\text{E}_2 - \text{P}$ form deoccludes, opens the gate for the remaining two Na^+ ions, and the affinity decreases for Na^+ and increases for K^+ . By this, the two Na^+ are exchanged for two K^+ ions from the extracellular medium ($\text{E}_2 - \text{PNa}_2$ to $\text{E}_2 - \text{PK}_2$). With two K^+ ions bound the system is dephosphorylated; the extracellular gate closes and the two K^+ ions are

occluded ($\text{E}_2' \cdot (2\text{K})$).^[69, 70] ATP bound to $\text{E}_2' \cdot (2\text{K})$ on the cytoplasmic side, with an affinity which is low (apparent K_D is about 0.1 mM), increases the rate at which the cytoplasmic gate opens and decreases the affinity for K^+ ($\text{E}_2' \cdot (2\text{K}) \cdot \text{ATP}$ to $\text{E}_2' \cdot (2\text{K}) \cdot \text{ATP}$).^[69] The two K^+ ions are exchanged for three Na^+ ions from the cytoplasmic medium, and this closes the cycle.

The transport system is a very efficient pump. It pumps three Na^+ ions out of and two K^+ ions into the cell for each ATP hydrolyzed to ADP and P_i , and for this it uses 70–85% of the free energy of the hydrolysis of ATP.^[50] With the normal intra- and extracellular concentrations of Na^+ and K^+ , the activity of the pump is 10–15% of maximum; that is, the pump has a considerable reserve power. At 37°C and pH 7.4 the enzyme turns over at a rate of about 160 per second.

In the transport reaction one net positive charge is carried to the outside. The transport is therefore electrogenic, but due to a high permeability of the membrane to Cl^- it only adds a few millivolts to the membrane potential. Each of the two conformations E_1 and E_2 seem to have two negatively charged binding sites on which the positive charge of the cations are neutralized; in addition, the E_1 conformation, as mentioned above, binds a third Na^+ ion in ionic form.^[71, 72]

To obtain transport of the cations against an electrochemical gradient coupled to a chemical reaction, the hydrolysis of ATP, it is necessary that there is a tight coupling between each of the steps in the catalytic reaction and each of the corresponding steps in the translocation reaction, as suggested in the model. The shift in affinities, and the opening and closing of the gates, are coupled to the stepwise degradation of ATP from phosphorylation with a high-energy $\text{E}_1 \sim \text{P}$ bond to a low-energy $\text{E}_2 \sim \text{P}$ bond, and finally to dephosphorylation. This divides the translocation into steps in which local gradients are created along which the ions can flow, and thereby they are transported against their overall gradient.

The model is a working hypothesis which explain a good deal of the experimental observations. There are, however, observations which do not fit with the model. And even though we have much information about the structure, there are many unanswered questions, especially at the molecular level: What is the nature of the gates, where are the binding sites for the cations, and of what nature, and how does the system discriminate between Na^+ and K^+ , etc.

Conclusion

It may seem disappointing that 30 years of work, since the conclusion that the membrane-bound Na^+/K^+ -ATPase is identical with the $\text{Na}^+ - \text{K}^+$ pump, has not given us an understanding of the basic molecular events behind the transport. However, considering that the problem is to reveal how 1320 amino acids inside a volume of $60 \times 60 \times 100 \text{ \AA}^3$ can be assembled to a very efficient machine which can convert the chemical energy from the hydrolysis of ATP into work, namely the transport of cations against their electrochemical gradient, and which can distinguish between so closely related cations as Na^+ and K^+ , it can be of no surprise that progress is slow.

Thirty years ago it seemed impossible to find a way to purify the enzyme, not to speak of getting it into solution and reconstituting it into liposomes. Nobody dared to dream about knowing the sequence, and yet this and much more has been accomplished.

Received: April 17, 1998 [A278IE]

German version: *Angew. Chem.* **1988**, *110*, 2452–2461

Keywords: ATP • ion transport • membranes • Nobel lecture • sodium–potassium pump

- [1] E. Gorter, F. Grendel, *J. Exp. Med.* **1925**, *41*, 439.
- [2] J. F. Danielli, H. Davson, *J. Cell. Comp. Physiol.* **1935**, *5*, 495.
- [3] P. J. Boyle, E. F. Conway, *J. Physiol.* **1941**, *100*, 1.
- [4] R. B. Dean, *Biol. Symp.* **1941**, *3*, 331.
- [5] G. Hevesy, *Enzymologia* **1938**, *5*, 138.
- [6] H. H. Ussing, *Physiol. Rev.* **1949**, *29*, 129.
- [7] A. Krogh, *Proc. R. Soc. London B* **1946**, *133*, 140.
- [8] E. J. Conway, *Nature* **1946**, *157*, 715.
- [9] H. H. Ussing, *Nature* **1947**, *160*, 262.
- [10] T. Rosenberg, *Acta Chem. Scand.* **1948**, *2*, 14.
- [11] I. M. Glynn in *The Enzymes of Biological Membranes*, Vol. 3 (Ed.: A. N. Martonosi), New York, Plenum **1985**, p. 28.
- [12] J. C. Skou, *Biochim. Biophys. Acta* **1989**, *1000*, 435.
- [13] J. C. Skou in *Membrane Transport: People and Ideas* (Ed.: D. Tosteson), American Physiological Society **1989**, p. 155.
- [14] „Lokalanestetika“: J. C. Skou, Dissertation, Universitetsforlaget Aarhus, **1954**.
- [15] J. C. Skou, *J. Pharm. Pharmacol.* **1961**, *13*, 204.
- [16] H. H. Meyer, *Arch. Exp. Pathol. Pharmacol.* **1899**, *42*, 109.
- [17] E. Overton, *Studien über die Narkose zugleich ein Beitrag zur allgemeinen Pharmakologie*, Jena, Fischer, **1901**.
- [18] N. K. Adam, *The Physics and Chemistry of Surfaces*, London, Oxford University Press, **1941**.
- [19] A. L. Hodgkin, B. Katz, *J. Physiol. (London)* **1949**, *108*, 37.
- [20] B. Libet, *Fed. Proc.* **1948**, *7*, 72.
- [21] A. L. Hodgkin, R. D. Keynes, *J. Physiol. (London)* **1955**, *128*, 28.
- [22] G. Gardos, *Acta Physiol. Acad. Sci. Hung.* **1954**, *6*, 191.
- [23] A. L. Hodgkin, R. D. Keynes, *J. Physiol. (London)* **1956**, *131*, 592.
- [24] P. C. Caldwell, R. D. Keynes, *J. Physiol. (London)* **1957**, *137*, 12P.
- [25] R. D. Keynes, in *Metabolism of the Nervous System* (Ed.: D. Richter), Pergamon, London, **1957**, p. 159.
- [26] J. C. Skou, in *Metabolism of the Nervous System* (Ed.: D. Richter), Pergamon, London, **1957**, p. 173.
- [27] J. C. Skou, *Biochim. Biophys. Acta* **1957**, *23*, 394.
- [28] R. L. Post, P. Jolly, *Biochim. Biophys. Acta* **1957**, *25*, 118.
- [29] H. J. Schatzmann, *Helv. Physiol. Pharmacol. Acta* **1953**, *11*, 346.
- [30] J. C. Skou, *Biochim. Biophys. Acta* **1960**, *42*, 6.
- [31] J. C. Skou, *Biochim. Biophys. Acta* **1962**, *58*, 314.
- [32] M. Esmann, J. C. Skou, C. Christiansen, *Biochim. Biophys. Acta* **1979**, *567*, 410.
- [33] R. L. Post, C. R. Merritt, C. R. Kinsolving, C. D. Albright, *J. Biol. Chem.* **1960**, *235*, 1796.
- [34] J. C. Skou, *Physiol. Rev.* **1965**, *45*, 596.
- [35] A. L. Hodgkin, A. F. Huxley, *J. Physiol. (London)* **1952**, *116*, 449.
- [36] P. Mitchell, *Biochem. Soc. Symp.* **1963**, *22*, 142.
- [37] R. K. Crane, *Comp. Biochem.* **1983**, *35*, 43.
- [38] H. Kromphardt, H. Grobecker, K. Ring, E. Heinz, *Biochim. Biophys. Acta* **1963**, *74*, 549.
- [39] E. K. Hoffmann, L. O. Simonsen, *Physiol. Rev.* **1989**, *69*, 315.
- [40] B. J. R. Pitts, *J. Biol. Chem.* **1979**, *254*, 6232.
- [41] J. D. Robertson, *Biochem. Soc. Symp.* **1959**, *16*, 3.
- [42] S. J. Singer, G. L. Nicolson, *Science* **1972**, *175*, 720.
- [43] W. Hasselbach, M. Makinose, *Biochem. Z.* **1961**, *333*, 518.
- [44] H. J. Schatzmann, *Experientia* **1966**, *22*, 364.
- [45] B. Stewart, B. Wallmark, G. Sachs, *J. Biol. Chem.* **1981**, *256*, 2682.
- [46] J. D. Robinson, *Moving Questions. A History of Membrane Transport and Bioenergetics*, American Physiological Society, NY **1997**.
- [47] *Na/K-ATPase and Related Transport ATPases: Structure, Mechanism and Regulation*, Vol. 834 (Eds.: L. A. Beaugé, D. C. Gadsby, P. J. Garrahan), Annals New York Academy Science, New York, **1997**.
- [48] P. L. Jørgensen, in *Molecular Aspects of Transport Proteins* (Ed.: J. H. M. M. De Pont), Elsevier Service Publisher, **1992**, p. 1.
- [49] I. M. Glynn, *J. Physiol. (London)* **1993**, *462*, 1.
- [50] F. Cornelius, *Biomembranes* **1996**, *5*, 133.
- [51] R. W. Albers, *Annu. Rev. Biochem.* **1967**, *36*, 727.
- [52] R. W. Albers, S. Fahn, G. J. Koval, *Proc. Natl. Acad. Sci. USA* **1963**, *50*, 474.
- [53] S. Fahn, G. J. Koval, R. W. Albers, *J. Biol. Chem.* **1968**, *243*, 1993.
- [54] R. L. Post, S. Kume, *J. Biol. Chem.* **1973**, *248*, 6993.
- [55] R. L. Post, S. Kume, T. Tobin, B. Orcutt, A. K. Sen, *J. Gen. Physiol.* **1969**, *54*, 306S.
- [56] R. L. Post, A. K. Sen, A. S. Rosenthal, *J. Biol. Chem.* **1965**, *240*, 1437.
- [57] P. Läuger, *Biochim. Biophys. Acta* **1979**, *552*, 143.
- [58] D. C. Gadsby, R. F. Rakowsky, P. De Weer, *Science* **1993**, *260*, 100.
- [59] J. G. Nørby, J. Jensen, *Biochim. Biophys. Acta* **1971**, *233*, 104.
- [60] C. Hegyvary, R. L. Post, *J. Biol. Chem.* **1971**, *246*, 5234.
- [61] P. L. Jørgensen, *Biochim. Biophys. Acta* **1975**, *401*, 399.
- [62] S. J. D. Karlsh, D. W. Yates, *Biochim. Biophys. Acta* **1978**, *527*, 115.
- [63] S. J. D. Karlsh, D. W. Yates, I. M. Glynn, *Biochim. Biophys. Acta* **1978**, *525*, 252.
- [64] J. C. Skou, M. Esmann, *Biochim. Biophys. Acta* **1981**, *647*, 232.
- [65] S. J. D. Karlsh, *J. Bioenerg. Biomembr.* **1980**, *12*, 111.
- [66] J. G. Nørby, I. Klodos, N. O. Christiansen, *J. Gen. Physiol.* **1983**, *82*, 725.
- [67] S. Yoda, A. Yoda, *J. Biol. Chem.* **1986**, *26*, 1147.
- [68] I. M. Glynn, Y. Hara, D. E. Richards, *J. Physiol. (London)* **1984**, *351*, 531.
- [69] R. L. Post, C. Hegyvary, S. Kume, *J. Biol. Chem.* **1972**, *247*, 6530.
- [70] L. A. Beaugé, I. M. Glynn, *Nature* **1979**, *280*, 510.
- [71] M. Nakao, D. C. Gadsby, *Nature* **1986**, *323*, 628.
- [72] R. Goldschleger, S. J. D. Karlsh, A. Raphaeli, W. D. Stein, *J. Physiol. (London)* **1987**, *387*, 331.
- [73] J. C. Skou, *News Physiol. Sci.* **1992**, *7*, 95.