

from triplet states of a ligand-to-metal charge transfer character that mix with metal-centered (ds/dp) states modified by Au...Au interactions (LMMCT; S → Au), whereas the high-energy emission at ca. 500 nm is attributed to metal-perturbed intraligand phosphorescence.

Received: December 20, 1999 [Z14429]

- [1] a) V. W.-W. Yam, K. K.-W. Lo, *Chem. Soc. Rev.* **1999**, 323, and references therein; b) P. C. Ford, A. Vogler, *Acc. Chem. Res.* **1993**, 26, 220, and references therein; c) W.-F. Fu, K.-C. Chan, V. M. Miskowski, C.-M. Che, *Angew. Chem.* **1999**, 111, 2953; *Angew. Chem. Int. Ed.* **1999**, 38, 2783; d) P. D. Harvey, H. B. Gray, *J. Am. Chem. Soc.* **1988**, 110, 2145.
- [2] P. Pykkö, *Chem. Rev.* **1988**, 88, 563.
- [3] a) H. Schmidbaur, *Chem. Soc. Rev.* **1995**, 391; b) *Gold: Progress in Chemistry, Biochemistry, and Technology* (Ed.: H. Schmidbaur), Wiley, Chichester, **1999**, and references therein; c) P. Pykkö, *Chem. Rev.* **1997**, 97, 597.
- [4] a) S. Dehnen, D. Fenske, A. C. Deveson, *J. Cluster Sci.* **1996**, 7, 351; b) S. Dehnen, D. Fenske, *Chem. Eur. J.* **1996**, 2, 1407; c) A. C. Deveson, S. Dehnen, D. Fenske, *J. Chem. Soc. Dalton Trans.* **1997**, 4491; d) J. F. Corrigan, D. Fenske, *Angew. Chem.* **1997**, 109, 2070; *Angew. Chem. Int. Ed. Engl.* **1997**, 36, 1981; e) J. F. Corrigan, D. Fenske, W. P. Power, *Angew. Chem.* **1997**, 109, 1224; *Angew. Chem. Int. Ed. Engl.* **1997**, 36, 1176.
- [5] a) P. Betz, B. Krebs, G. Henkel, *Angew. Chem.* **1984**, 96, 293; *Angew. Chem. Int. Ed. Engl.* **1984**, 23, 311; b) G. Marbach, J. Strähle, *Angew. Chem.* **1984**, 96, 695; *Angew. Chem. Int. Ed. Engl.* **1984**, 23, 715.
- [6] V. W.-W. Yam, E. C.-C. Cheng, K.-K. Cheung, *Angew. Chem.* **1999**, 111, 193; *Angew. Chem. Int. Ed.* **1999**, 38, 197.
- [7] a) C. Kowala, J. M. Swan, *Aust. J. Chem.* **1966**, 19, 547; b) P. G. Jones, G. M. Sheldrick, E. Hädicke, *Acta. Crystallogr. Sect. B* **1980**, 36, 2777; c) C. Lensch, P. G. Jones, G. M. Sheldrick, *Z. Naturforsch.* **1982**, 37b, 944; d) K. Angermaier, H. Schmidbaur, *Z. Naturforsch.* **1996**, 51, 879; e) F. Canales, M. C. Gimeno, P. G. Jones, A. Laguna, *Angew. Chem.* **1994**, 106, 811; *Angew. Chem. Int. Ed. Engl.* **1994**, 33, 769; f) S. Canales, O. Crespo, M. C. Gimeno, P. G. Jones, A. Laguna, *Chem. Commun.* **1999**, 679.
- [8] a) V. W.-W. Yam, W.-K. Lee, T.-F. Lai, *Chem. Commun.* **1993**, 1571; b) V. W.-W. Yam, K. K.-W. Lo, K.-K. Cheung, *Inorg. Chem.* **1996**, 35, 3459; c) V. W.-W. Yam, K. K.-W. Lo, C.-R. Wang, K.-K. Cheung, *Inorg. Chem.* **1996**, 35, 5116; d) V. W.-W. Yam, K. K.-W. Lo, C.-R. Wang, K.-K. Cheung, *J. Phys. Chem. A* **1997**, 101, 4666; e) V. W.-W. Yam, K. K.-W. Lo, *Comments Inorg. Chem.* **1997**, 19, 209.
- [9] V. W.-W. Yam, C.-L. Chan, K.-K. Cheung, *J. Chem. Soc. Dalton Trans.* **1996**, 4019.
- [10] **1**: Elemental analysis calcd for $[\text{Au}_{10}(\mu\text{-PNP})_4(\mu_3\text{-S})_4](\text{PF}_6)_2 \cdot (\text{CH}_3)_2\text{CO}$ (found): C 32.08 (32.34), H 2.76 (2.75), N 1.38 (1.43); positive FAB-MS: m/z : 1903 $[\text{M} - 2\text{PF}_6]^{2+}$; positive ESI-MS: m/z : 1903 $[\text{M} - 2\text{PF}_6]^{2+}$; ^1H NMR (300 MHz, CD_2Cl_2 , 298 K): δ = 0.04 (m, 12H, CH_3), 0.27 (m, 4H, CCH_2C), 0.85 (m, 4H, CCH_2C), 2.33 (m, 4H, NCH_2), 2.80 (m, 4H, NCH_2), 6.83 (m, 8H, Ph), 7.12–7.83 (m, 72H, Ph); $^{31}\text{P}\{^1\text{H}\}$ NMR (202 MHz, CD_2Cl_2 , 298 K): δ = 83.1, 85.5 (dd, J_{PP} = 115 Hz); molar conductivity (acetone, 298 K): $\Lambda_{\text{M}} = 262 \Omega^{-1} \text{cm}^2 \text{mol}^{-1}$ (1:2 electrolyte).
- [11] a) Crystal data for **1**: $\text{Au}_{10}\text{S}_4(\text{C}_{27}\text{H}_{27}\text{P}_2\text{N})_4 \cdot 2\text{PF}_6$, $M_r = 4097.59$, tetragonal, space group $I4$ (no. 82), $a = 18.9950(14)$, $c = 17.7282(19) \text{ \AA}$, $V = 6396.5(10) \text{ \AA}^3$, $Z = 2$, $\rho_{\text{calcd}} = 2.127$, $\mu(\text{MoK}\alpha) = 11.670 \text{ mm}^{-1}$, $F(000) = 3792$, $T = 293 \text{ K}$; $R = 0.0347$, $wR = 0.0870$ for 7314 reflections with $I > 2\sigma(I)$. Crystal dimensions $0.20 \times 0.14 \times 0.12 \text{ mm}$; Bruker CCD area detector diffractometer, $\text{MoK}\alpha$ radiation ($\lambda = 0.71073 \text{ \AA}$); collection range $2\theta = 3\text{--}55^\circ$; Φ and Ω oscillation frames $0\text{--}180^\circ$; 1321 frames taken in four shells. Absorption correction applied by using the program SADABS^[11b]; the structure was determined by direct methods and subsequent difference Fourier syntheses, and refined by full-matrix least-squares on F^2 by using the programs of the Bruker Smart and Bruker Shelxtl package^[11b]. All non-hydrogen atoms were refined anisotropically. Crystallographic data (excluding structure factors) for the structure reported in this paper has been deposited with the Cambridge Crystallographic Data Centre as supplementary


publication no. CCDC-138838. Copies of the data can be obtained free of charge on application to CCDC, 12 Union Road, Cambridge CB2 1EZ, UK (fax: (+44) 1223-336-033; e-mail: deposit@ccdc.cam.ac.uk.); b) SADABS, Bruker Smart and Bruker Shelxtl Package, Bruker AXS GmbH, **1998**.

- [12] P. G. Jones, C. Lensch, G. M. Sheldrick, *Z. Naturforsch. B* **1982**, 37, 141.
- [13] J. Vicente, M. T. Chicote, P. González-Herrero, C. Grünwald, P. G. Jones, *Organometallics* **1997**, 16, 3381.
- [14] M. S. Balakrishna, V. S. Reddy, S. S. Krishnamurthy, *Coord. Chem. Rev.* **1994**, 129, 1.
- [15] a) H. Friebolin, *Basic One- and Two-Dimensional NMR Spectroscopy*, VCH, Weinheim, **1991**, p. 279; b) D. L. Griffith and J. D. Roberts, *J. Am. Chem. Soc.* **1965**, 87, 4089; c) E. Fahr, W. Rohlfing, R. Thiedemann, A. Mannschreck, G. Rissmann, W. Seitz, *Tetrahedron Lett.* **1970**, 3605.

Metal-Mediated Transfer of Electrons between Two Different C–C Single Bonds That Function as Electron-Donor and Electron-Acceptor Units**

Federico Franceschi, Euro Solari, Rosario Scopelliti, and Carlo Floriani*

Two complementary electron-storage and -releasing units are the fundamental constituents of storage cells^[1] and molecular devices devoted to long-range electron-transfer processes.^[2] The C–C single bonds formed by oxidative (Scheme 1a)^[3] or reductive coupling^[4] (Scheme 1b) can be



Scheme 1. Reversible two-electron oxidative (a) and reductive coupling (b) to form C–C bonds. R = Et.

[*] Prof. Dr. C. Floriani, Dr. F. Franceschi, Dr. E. Solari, Dr. R. Scopelliti
Institut de Chimie Minérale et Analytique
Université de Lausanne, BCH
1015 Lausanne (Switzerland)
Fax: (+41) 21-6923905
E-mail: carlo.floriani@icma.unil.ch

[**] This work was supported by the Fonds National Suisse de la Recherche Scientifique (Grant No. 20-53336.98) and Action COST D9 (European Program for Scientific Research, OFES No. C98.008).


Supporting information for this article is available on the WWW under <http://www.wiley-vch.de/home/angewandte/> or from the author.

used for storing and releasing pairs of electrons in a reversible manner.


The novelty of the system reported here lies in the nature of the electron-storage and -release units, both of which are ubiquitous C–C single bonds, and on how the communication between them is mediated by metal ions. In addition, the two complementary C–C units operate without being involved, at any stage, in chemical reactions.

Redox chemistry involving exclusively C–C single bonds was investigated for complexes **1**, **2a**, and **2b**, and the complementary behavior of the two different C–C bonds was demonstrated. Although the two complementary systems are chemically quite different, they both bind alkali metal cations at the periphery.^[5, 6] They form tight ion-pair structures, in which the pyrrolyl anions or the Schiff base oxygen atoms, respectively, function as binding sites for the alkali metal cations. The oxidizing properties of **1** were demonstrated, among others, in its reaction with LiMe,^[7] which led to the expected evolution of a mixture of CH₄ and C₂H₆ and the cleavage of the cyclopropane unit (Scheme 2). The partner reaction of **2a** or **2b** with CH₃I^[7] also resulted in the formation of C₂H₆ and cleavage of the C–C bond across the two salophen units, whereby the C–C bond functioned as a two-electron donor.

The complementarity of the two processes was then proved by adding a solution of **2a** or **2b** in THF to a solution of **1** in THF, which led to immediate precipitation of **5a** or **5b**, respectively (Scheme 3). The latter was recov-



Scheme 2. A C–C bond functioning as a two-electron oxidizing (a) and reducing agent (b).



Scheme 3. Electron transfer system between electron-rich and electron-poor C–C bonds assisted by alkali metal ions.

ered as the ion-separated compound **6**^[8] under the crystallization conditions, rather than in the ion-pair form. The crystal structure of **6** is shown in Figure 1^[9].

The occurrence of intermolecular electron transfer and the nature of the final compound emphasizes the assistance of both the central transition metal ion and the alkali metal ion in the redox process. The chemical transformation displayed in Scheme 3 could be exploited for energy storage (e.g., storage cells) or transfer. Although for both model compounds, **2** and **1**, the C–C bond is the site of storage or deficiency, respectively, of a pair of electrons, these functional groups are not involved in any chemical reactions; the metal atom assists electron transfer from the C–C bond to the substrate.^[4e, f] This chemical behavior ensures complete chemical and energetic reversibility of the system. Apart from the practical applicability of this system, the novel concept of exploiting ubiquitous C–C bonds for energy-storing and electron-transfer processes is noteworthy.




Figure 1. Crystal structure of **6** (hydrogen atoms and interstitial CH₃CN solvent molecules omitted for clarity).

Experimental Section

1: A solution of iodine (3.21 g, 12.7 mmol) in THF (100 mL) was added dropwise to a solution of [NiEt₈Py₄Na₂]·2Et₂O^[7] (10 g, 12.7 mmol; Py = pyrrolyl) in THF (100 mL). The resulting red suspension was stirred for 12 h and then concentrated to dryness. The residue was extracted with CH₂Cl₂ (60 mL) to give a suspension from which a red microcrystalline solid was collected and dried in vacuo (5.97 g, 79%). Elemental analysis calcd for C₃₆H₄₈N₄Ni: C 72.61, H 8.12, N 9.41; found: C 72.58, H 8.05, N 9.23; ¹H NMR (400 MHz, [D₆]benzene, 298 K): δ = 6.67 (d, *J* = 5.1 Hz, 2H, CH), 6.53 (d, *J* = 2.7 Hz, 2H, CH), 6.32 (d, *J* = 2.7 Hz, 2H, CH), 6.11 (d, *J* = 5.1 Hz, 2H, CH), 3.57 (q, *J* = 7.0 Hz, 2H, CH₂), 2.87 (q, *J* = 7.4 Hz, 2H, CH₂), 2.50 (q, *J* = 7.2 Hz, 2H, CH₂), 2.28 (m, 2H, CH₂), 1.84 (m, 4H, CH₂), 1.69 (q, *J* = 6.8 Hz, 2H, CH₂), 1.31–1.18 (m, 8H, CH₂, CH₃), 1.02 (t, *J* = 7.0 Hz, 3H, CH₃), 0.88 (m, 6H, CH₃), 0.70 (m, 6H, CH₃), 0.48 (t, *J* = 6.85 Hz, 3H, CH₃); ¹³C NMR (100.6 MHz, [D₆]benzene, 298 K): δ = 147.6, 139.3, 130.7, 130.0, 104.5, 104.0, 85.8, 67.1, 57.7, 51.8, 47.5, 44.8, 39.3, 33.3, 30.0, 27.9, 25.1, 11.2, 11.1, 11.0, 10.7, 10.3, 10.1.

5a: A red solution of **1** (0.75 g, 1.26 mmol) in THF (50 mL) was added to a dark green suspension of **2a**^[4a] (1.32 g, 1.26 mmol) in THF (50 mL). A brown-red precipitate separated immediately from the resulting brown solution. The suspension was stirred for 12 h then the solid was collected and dried in vacuo (1.41 g, 75%). Elemental analysis calcd for **5a**, C₈₄H₉₂Li₂N₈Ni₃O₆: C 65.87, H 6.05, N 7.31; found: C 65.38, H 5.75, N 7.30. IR (Nujol): $\tilde{\nu}$ = 1606 (s), 1578 (s), 1534 (s), 1488 (m), 1444 (s), 1326 (s), 1262 (w), 1196 (s), 1152 (m), 1129 (m), 1068 (w), 1042 (m), 950 (w), 929 (m), 880 (w), 840 (w), 813 (m), 747 (s), 735 (s), 721 (m), 700 (m), 622 (w), 570 (w), 547 (m), 507 (w), 450 (w), 408 cm⁻¹ (w).

5b and **6:** A red solution of **1** (0.87 g, 1.46 mmol) in THF (100 mL) was added to a dark green solution of **2b**^[4a] (1.79 g, 1.46 mmol) in THF (100 mL). A red precipitate separated immediately from the resulting dark orange-red solution. The suspension was stirred overnight then the solid was collected and dried in vacuo (1.81 g, 81%). Elemental analysis calcd for **5b**, C₈₄H₉₂N₈Na₂Ni₃O₆: C 65.87, H 6.05, N 7.31; found: C 65.38, H 5.75, N 7.30. IR (Nujol): $\tilde{\nu}$ = 1608 (s), 1578 (s), 1527 (s), 1483 (s), 1356 (m), 1330 (s), 1317 (s), 1286 (s), 1258 (m), 1235 (w), 1193 (s), 1151 (s), 1129 (m), 1053 (w), 1025 (w), 929 (m), 901 (w), 871 (w), 848 (w), 808 (w), 755 (s), 744 (s), 713 (s), 649 (w), 612 (w), 584 (w), 565 (s), 545 (s), 483 (w), 452 (m), 410 cm⁻¹ (m). The product was recrystallized from acetonitrile to give deep red crystals of **6** that were suitable for X-ray analysis. The crystals were collected, dried in vacuo, and analyzed. Elemental analysis calcd for **6**·CH₃CN, C₉₈H₉₃N₁₁Na₂Ni₄O₆: C 65.33, H 5.20, N 8.55; found: C 65.23, H 4.65, N 8.38.

Received: November 18, 1999 [Z14294]

- [1] a) A. Cisak, L. Werblan, *High-Energy Non-Aqueous Batteries*, Ellis Horwood, Chichester, **1993**; b) C. A. Vincent, B. Scrosati, *Modern Batteries*, 2nd ed., Arnold, London, **1997**; c) *Chemistry for the Energy Future* (Eds.: V. N. Parmon, H. Tributsch, A. V. Bridgewater, D. O. Hall), Blackwell Science, Oxford, UK, **1999**.
- [2] a) *Electron Transfer in Biology and the Solid State* (Eds.: M. J. Johnson, R. B. King, D. M. Kurtz, Jr., C. Kotal, M. L. Norton, R. A. Scott), American Chemical Society, Washington, DC, **1990**; b) A. G. Lippin, *Redox Mechanisms in Inorganic Chemistry*, Ellis Horwood, New York, **1994**.
- [3] a) S. De Angelis, E. Solari, C. Floriani, A. Chiesi-Villa, C. Rizzoli, *J. Am. Chem. Soc.* **1994**, *116*, 5691–5701; b) S. De Angelis, E. Solari, C. Floriani, A. Chiesi-Villa, C. Rizzoli, *J. Am. Chem. Soc.* **1994**, *116*, 5702–5713; c) C. Floriani in *Transition Metals in Supramolecular Chemistry* (Eds.: L. Fabbrizzi, A. Poggi), Kluwer, Amsterdam, **1994**, pp. 191–209; d) C. Floriani, *Chimia* **1996**, *50*, 608–611; e) C. Floriani, *Chem. Commun.* **1996**, 1257–1263; f) U. Piarulli, E. Solari, C. Floriani, A. Chiesi-Villa, C. Rizzoli, *J. Am. Chem. Soc.* **1996**, *118*, 3634–3642; g) R. Crescenzi, E. Solari, C. Floriani, A. Chiesi-Villa, C. Rizzoli, *Inorg. Chem.* **1998**, *37*, 6044–6051; h) R. Crescenzi, E. Solari, C. Floriani, A. Chiesi-Villa, C. Rizzoli, *J. Am. Chem. Soc.* **1999**, *121*, 1695–1706.
- [4] a) S. Gambarotta, F. Urso, C. Floriani, A. Chiesi-Villa, C. Guastini, *Inorg. Chem.* **1983**, *22*, 3966–3972; b) S. De Angelis, E. Solari, E. Gallo, C. Floriani, A. Chiesi-Villa, C. Rizzoli, *Inorg. Chem.* **1996**, *35*, 5995–6003; c) E. Gallo, E. Solari, N. Re, C. Floriani, A. Chiesi-Villa, C. Rizzoli, *J. Am. Chem. Soc.* **1997**, *119*, 5144–5154; d) E. Solari, C. Maltese, F. Franceschi, C. Floriani, A. Chiesi-Villa, C. Rizzoli, *J. Chem. Soc. Dalton Trans.* **1997**, 2903–2910; e) F. Franceschi, E. Solari, C. Floriani, A. Chiesi-Villa, C. Rizzoli, M. Rosi, *Chem. Eur. J.* **1999**, *5*, 708–721; f) M. Rosi, A. Sgamellotti, F. Franceschi, C. Floriani, *Chem. Eur. J.* **1999**, *5*, 2914–2920.
- [5] a) S. De Angelis, E. Solari, C. Floriani, A. Chiesi-Villa, C. Rizzoli, *J. Chem. Soc. Dalton Trans.* **1994**, 2467–2469; b) S. De Angelis, E. Solari, C. Floriani, A. Chiesi-Villa, C. Rizzoli, *Organometallics* **1995**, *14*, 4505–4512; c) D. Jacoby, S. Isoz, C. Floriani, K. Schenk, A. Chiesi-Villa, C. Rizzoli, *Organometallics* **1995**, *14*, 4816–4824; d) C. Floriani, *Pure Appl. Chem.* **1996**, *68*, 1–8; e) C. Floriani, E. Solari, G. Solari, A. Chiesi-Villa, C. Rizzoli, *Angew. Chem.* **1998**, *110*, 2367–2369; *Angew. Chem. Int. Ed.* **1998**, *37*, 2245–2248.
- [6] a) N. Bresciani-Pahor, M. Calligaris, P. Delise, G. Nardin, L. Randaccio, E. Zotti, G. Fachinetti, C. Floriani, *J. Chem. Soc. Dalton Trans.* **1976**, 2310–2316; b) J. C. Pessoa, J. A. L. Silva, A. L. Viera, L. Vilas-Boas, P. O'Brien, P. Thornton, *J. Chem. Soc. Dalton Trans.* **1992**, 1745–1749; c) S. C. Ball, I. Cragg-Hine, M. G. Davidson, R. P. Davies, M. I. Lopez-Solera, P. R. Raithby, D. Reed, R. Snaith, E. M. Vogl, *J. Chem. Soc. Chem. Commun.* **1995**, 2147–2149; d) D. A. Atwood, D. Rutherford, *Inorg. Chem.* **1995**, *34*, 4008–4010; e) H. Miyasaka, N. Matsumoto, H. Okawa, N. Re, E. Gallo, C. Floriani, *J. Am. Chem. Soc.* **1996**, *118*, 981–994; f) N. Re, E. Gallo, C. Floriani, H. Miyasaka, N. Matsumoto, *Inorg. Chem.* **1996**, *35*, 5964–5965; g) E. Gallo, E. Solari, C. Floriani, A. Chiesi-Villa, C. Rizzoli, *Inorg. Chem.* **1997**, *36*, 2178–2186.
- [7] For experimental details see Supporting Information.
- [8] In such complexes, the ligand distribution often gives rise to the least soluble complex in the solid state; hence, the [Ni(salophen)]/Na ratio is 3:2 rather than 1:1.^[4,6]
- [9] Crystal structure analysis of **6**: [C₆₆H₅₁N₉Na₂Ni₅O₆][C₃₆H₄₈N₄Ni]·3C₂H₃N, *M*_r = 2006.92, orthorhombic, space group *Pna*2₁, *a* = 20.868(3), *b* = 21.738(3), *c* = 20.9686(14) Å, *V* = 9512.0(19) Å³, *Z* = 4, ρ_{calcd} = 1.401 g cm⁻³, *F*(000) = 4200, MoK_α radiation (λ = 0.71073 Å), μ(MoK_α) = 0.855 mm⁻¹; crystal dimensions 0.20 × 0.14 × 0.10. For 13108 observed reflections (*I* > 2σ(*I*)) and 1227 parameters, the conventional *R* is 0.0926 (*wR*2 = 0.1572 for 15798 independent reflections). Crystallographic data (excluding structure factors) for the structure reported in this paper have been deposited with the Cambridge Crystallographic Data Centre as supplementary publication no. CCDC-137426. Copies of the data can be obtained free of charge on application to CCDC, 12 Union Road, Cambridge CB2 1EZ, UK (fax: (+44) 1223-336-033; e-mail: deposit@ccdc.cam.ac.uk).