

methanol, a drop of this suspension was then deposited on a sapphire. After evaporation of the solvent the sapphire was placed in the field of view of the NIRIM. A library of single-bead NIR–Raman spectra of each of the resins was first recorded (Supporting Information), then several regions were arbitrarily selected for multispectral imaging. Since each bead appears as a collection of pixels and each pixel is a NIR–Raman spectrum of that area of the bead, comparison of these pixel spectra with the library of single-bead spectra recorded on the authentic samples confirmed the automated assignments. These results were reproducible regardless of the size and shape of the beads (Supporting Information).

The principles of dual recursive deconvolution of resin-supported combinatorial libraries have been proposed and the feasibility of the key feature of this method, the identification of the first randomized position, has been demonstrated using NIR–Raman imaging of self-encoded resin beads. The reliability of this method is very high, it has been consistently accurate over the two and a half years that we have been practicing multispectral imaging. Furthermore, any imaging technique could be applicable to the DRED method as long as the beads used display unique spectral features, and provided the loaded material does not significantly alter their spectral signature. For instance, secondary-ion mass spectrometry (SIMS)^[10] and FT-IR^[11] imaging are alternative approaches that we are investigating. Although, several commercially available and chemically distinct solid supports are eligible to explore the scope of DRED in combinatorial chemistry, inhomogeneities in their physical and chemical properties (e.g. swelling, porosity, size, reactivity) prompted the design of the DRED beads, a new class of resins for solid-phase synthesis.^[12] The added value of the DRED method is that it is a non-invasive screening technique, it does not necessarily require sophisticated equipment, it is inexpensive, and it does not involve the development of any encoding chemistry since the DRED beads will soon become commercially available.

Received: August 4, 2000 [Z15581]

- [1] Á. Furka, L. K. Hamaker, M. L. Peterson in *Combinatorial Chemistry: A Practical Approach* (Ed.: H. Fenniri), Oxford University Press, Oxford, **2000**, pp. 1–32.
- [2] a) W. C. Still, *Acc. Chem. Res.* **1996**, 29, 155–163; b) W. L. Fitch, T. A. Baer, W. Chen, F. Holden, C. P. Holmes, D. MacLean, N. Shah, E. Sullivan, M. Tang, P. Waybourn, S. M. Fisher, C. A. Miller, L. R. Snyder, *J. Comb. Chem.* **1999**, 1, 188–194; c) M. C. Needels, D. G. Jones, E. H. Tate, G. L. Heinkel, L. M. Kochersperger, W. J. Dower, R. W. Barrett, M. A. Gallop, *Proc. Natl. Acad. Sci. USA* **1993**, 90, 10700–10704; d) K. S. Lam, M. Lebl, V. Krchnak, *Chem. Rev.* **1997**, 97, 411–448; e) X.-Y. Xiao, K. C. Nicolaou in *Combinatorial Chemistry: A Practical Approach* (Ed.: H. Fenniri), Oxford University Press, Oxford, **2000**, pp. 75–94.
- [3] a) A. Nefzi, J. M. Ostresh, R. A. Houghten, *Chem. Rev.* **1997**, 97, 449–472; b) D. L. Boger, W. Chai, Q. Jin, *J. Am. Chem. Soc.* **1998**, 120, 7220–7225; c) M. C. Pirrung, J. Chen, *J. Am. Chem. Soc.* **1995**, 117, 1240–1245.
- [4] A. D. Gift, J. Ma, K. S. Haber, B. L. McClain, D. Ben-Amotz, *J. Raman Spectrosc.* **1999**, 30, 757–765. The NIRIM instrument is currently housed in the Purdue University Laser Laboratory within the Chemistry Department. Web: <http://www.chem.purdue.edu/lfac/laser.html>

- [5] J. Ma, D. Ben-Amotz, *Appl. Spectrosc.* **1997**, 51, 1845–1848.
- [6] a) P. J. Treado, M. D. Morris, *Appl. Spectrosc. Rev.* **1994**, 29, 1–38; b) G. J. Puppels, M. Grond, J. Greve, *Appl. Spectrosc.* **1993**, 47, 1257; c) L. Markwort, B. Kip, E. D. Silva, B. Roussel, *Appl. Spectrosc.* **1995**, 49, 1411–1430.
- [7] 90 µm diameter polystyrene (PS; 1 % divinyl benzene(DVB)/PS) and 130 µm TentaGel-S-OH resin beads (Advanced ChemTech). See Supporting Information for the list of resin-supported amino acids studied. A detailed description of the design and operation of the NIRIM, and the processing software can be found in ref. [4] and in the Supporting Information.
- [8] This acquisition time could be decreased by at least an order of magnitude with a more powerful laser source and a more efficient optics setup.
- [9] “MultiSpec—A Tool for Multispectral Image Data Analysis”: L. Biehl, D. Langrebe, Pecora 13, Sioux Falls, SD, **1996**. The software can be downloaded at: <http://dynamo.ecn.purdue.edu/~biehl/multispec/>. See Supporting Information.
- [10] C. L. Brummel, J. C. Vickerman, S. A. Carr, M. E. Hemling, G. D. Roberts, W. Johnson, D. Gaitanopoulos, S. J. Benkovic, N. Winograd, *Anal. Chem.* **1996**, 68, 237–242.
- [11] H. Fenniri, J. Achkar, unpublished work.
- [12] H. Fenniri, PCT 60/180,939.

Polymorphism in *p*-Hydroxybenzoic Acid: The Effect of Intermolecular Hydrogen Bonding in Controlling Proton Order versus Disorder in the Carboxylic Acid Dimer Motif**

Benson M. Kariuki, Clare L. Bauer,
Kenneth D. M. Harris,* and Simon J. Teat

The phenomenon of polymorphism^[1–3] is of considerable contemporary interest in the field of organic solid-state chemistry, in part because comparisons between the properties of polymorphs provide an ideal basis on which to understand relationships between solid-state properties and crystal structure. Furthermore, in the quest to develop reliable computational techniques to predict the crystal structure(s) formed by a given type of organic molecule,^[4–8] studies of polymorphic systems provide stringent challenges for assessing the success of these methods. In general, the fact that the structures of organic molecular crystals usually arise from the interplay of several factors of comparable importance leads to intrinsic difficulties in attempting to predict and/or rationalize such structures. However, when one type of intermolecular interaction (or a small number of interactions) is dominant, a reliable rationalization of the observed molecular packing

[*] Prof. Dr. K. D. M. Harris, Dr. B. M. Kariuki, Dr. C. L. Bauer
School of Chemistry, University of Birmingham
Edgbaston, Birmingham B15 2TT (UK)
Fax: (+44) 121-414-7473
E-mail: K.D.M.Harris@bham.ac.uk
Dr. S. J. Teat
CCLRC Daresbury Laboratory
Daresbury, Warrington, Cheshire WA4 4AD (UK)

[**] We are grateful to the EPSRC and Daresbury Laboratory for financial support (CASE award to C.L.B.) and for the award of beam-time at Daresbury Laboratory.

arrangement in the crystal may be possible. In this regard, crystal structures based on strong intermolecular hydrogen bonding are often amenable to straightforward structural understanding.

The hydroxy derivatives of benzoic acid represent a potentially useful family of systems in which to explore polymorphism, as the combination of hydroxy groups and carboxylic acid groups should give rise to a variety of plausible intermolecular hydrogen bonding schemes and hence different crystal packing arrangements. Indeed, two polymorphs of *m*-hydroxybenzoic acid have been reported previously,^[9] and differ in their intermolecular hydrogen bonding arrangements. For *p*-hydroxybenzoic acid, on the other hand, only one crystal structure (hereafter referred to as form I) has been reported previously,^[10] although the possible existence of a second polymorph (which was neither isolated nor structurally characterized) was speculated^[11] on the basis of thermal analysis of samples of *p*-hydroxybenzoic acid.

Herein, we report a new polymorph (denoted form II) of *p*-hydroxybenzoic acid. Only a small number of low-quality microcrystals of this polymorph were obtained (as mixtures with other phases),^[12] and the crystal structure has been determined by exploiting synchrotron X-ray microcrystal diffraction techniques.^[13] Although the use of a microcrystal has led to a rather poorer quality of structural information^[13] than normally expected from single-crystal X-ray diffraction studies, unambiguous information on the crystal packing arrangement and other structural aspects is nevertheless obtained. Importantly, we find that form II exhibits interesting behavior with regard to the question of order versus disorder in the carboxylic acid dimer motif, contrasting markedly with the behavior of form I.

In the crystal structure^[13] of form II of *p*-hydroxybenzoic acid (Figure 1), the asymmetric unit comprises two independent molecules (denoted A and B). Each of these molecules forms an $R_2^2(8)$ centrosymmetric dimer about a crystallo-




Figure 1. Crystal structure of form II of *p*-hydroxybenzoic acid viewed along the *b* axis. Dotted lines represent hydrogen bonding interactions. Molecules of type A are labeled with oxygen atoms numbered 1, 2, and 3. Molecules of type B are labeled with oxygen atoms numbered 4, 5, and 6.

graphic inversion center, a hydrogen bond pattern (involving two $O-H \cdots O=C$ hydrogen bonds) that is commonly found in carboxylic acids.^[16, 17] Clearly the two molecules in a given dimer are of the same type (A or B). For the centrosymmetric dimer involving molecules of type A, the $O(1) \cdots O(2)$ distance is 2.664(6) Å, and for that involving molecules of type B, the $O(4) \cdots O(5)$ distance is 2.617(6) Å (the atom labeling is specified in Figure 1). In both cases, these $O \cdots O$ distances are consistent with $O-H \cdots O=C$ hydrogen bonding. A hydrogen bonding interaction involving the hydroxy group of molecule B and the carboxylic acid group of molecule A is also clearly identified. In this interaction, the $O \cdots O$ distance between the hydroxy oxygen atom O(6) of molecule B and one oxygen atom O(1) of the carboxylic acid group of molecule A is 2.946(6) Å and the $O-H \cdots O$ angle is about 165°. For the carboxylic acid group of molecule B, the shortest intermolecular $O \cdots O$ distances to hydroxy groups of other molecules are 3.427(8) Å and 3.439(9) Å—these $O \cdots O$ distances involve the same oxygen atom of the carboxylic acid group of molecule B and the hydroxy groups of two different molecules of type A (related to each other by translation along the *b* axis). However, these distances are rather long to be assigned as $O-H \cdots O=C$ hydrogen bonds^[19] and other geometric characteristics^[20] also deviate from those normally observed for strong $O-H \cdots O=C$ hydrogen bonds.^[19] In this structure, there are no short intermolecular contacts between hydroxy groups.

In the space-averaged and time-averaged crystal structures determined from diffraction data, $R_2^2(8)$ carboxylic acid dimer motifs are often found to be disordered,^[16] such that the carbon–oxygen bond lengths are intermediate between those characteristic of single bonds (corresponding to $C-OH$) and double bonds (corresponding to $C=O$) and the protons are disordered (often two sites with half occupancy are located, corresponding to equal amounts of $C-O-H \cdots O=C$ and $C=O \cdots H-O-C$ situations in the average crystal structure). We now assess the evidence for order versus disorder of the carboxylic acid dimers in form II of *p*-hydroxybenzoic acid. The carboxylic acid group in molecule A is clearly not disordered. The $C-O$ bond lengths are 1.230(5) Å for O(1) and 1.333(5) Å for O(2), leading to the unambiguous assignment^[21] of O(1) as $C=O$ and O(2) as $C-OH$ in an ordered carboxylic acid group. The fact that O(1) receives a hydrogen bond from the hydroxy group in molecule B is consistent with the preference for the $C=O$ oxygen atom in an ordered carboxylic acid group (rather than the $C-OH$ oxygen atom) to act as a hydrogen-bond acceptor. Clearly the intermolecular $O-H \cdots O=C$ hydrogen bond from the hydroxy group of molecule B to the carboxylic acid group of molecule A “locks” this carboxylic acid group into an ordered arrangement. Our ability to locate the hydrogen atoms of the carboxylic acid and hydroxy groups in the crystal structure determination is discussed in reference [13].

For molecule B, the carbon–oxygen bond lengths in the carboxylic acid group are 1.273(7) Å for O(4) and 1.299(7) Å for O(5). The fact that these bond lengths in the average crystal structure are intermediate between $C-O$ single and $C=O$ double bonds suggests that this carboxylic acid group is disordered. Presumably any intermolecular interactions in-

volving the carboxylic acid group of molecule B (in particular, the rather long contacts to the hydroxy groups of molecules of type A discussed above) are insufficient to "lock" the carboxylic acid group of molecule B into an ordered arrangement.

For comparison, the crystal structure of form I of *p*-hydroxybenzoic acid^[10] also contains carboxylic acid dimers with the $R_2^2(8)$ hydrogen bond motif, but there is no direct hydrogen bonding between hydroxy groups and carboxylic acid groups (Figure 2). Instead, the hydroxy groups engage in




Figure 2. Crystal structure of form I of *p*-hydroxybenzoic acid (data from reference [10]).

O—H...O—H hydrogen bonds with other hydroxy groups, constructing chains that extend through the crystal.^[22] The two carbon–oxygen bond lengths in the carboxylic acid group are equal (1.266 Å) (there is one molecule in the asymmetric unit), and the hydrogen atom of the carboxylic acid group is found to be distributed (with half occupancy) between two sites. These facts clearly indicate that the carboxylic acid group is disordered.

The subject of proton disorder in carboxylic acid dimers has aroused much interest from structural and dynamic viewpoints,^[23–30] and it is clear that the two polymorphs of *p*-hydroxybenzoic acid represent an ideal basis for systematic studies of this property. Thus, from the X-ray diffraction data reported here, form I exhibits proton disorder, whereas form II has two crystallographically independent carboxylic acid dimers, one of which is clearly ordered and the other disordered.^[31] Future neutron diffraction studies, also as a function of temperature, will provide more details on structural aspects of the proton order/disorder, whereas other techniques (solid-state NMR and inelastic and quasielastic neutron scattering) will lead to an understanding of dynamic aspects. Inter alia, these variable temperature studies will allow the static versus dynamic character of the disorder to be probed, as well as the question of whether polymorphic phase transformations are observed in this system. Such future advances will be facilitated by the ability to prepare form II as a monophasic sample, ideally comprising larger single crystals than the microcrystals available in the present work.

Received: February 14, 2000
Revised: August 30, 2000 [Z14700]

- [1] J. D. Dunitz, *Pure Appl. Chem.* **1991**, 63, 177.
[2] M. R. Caira, *Top. Curr. Chem.* **1998**, 198, 164.

- [3] J. Bernstein, R. J. Davey, J.-O. Henck, *Angew. Chem.* **1999**, 111, 3646; *Angew. Chem. Int. Ed.* **1999**, 38, 3441.
[4] H. R. Karfunkel, R. J. Gdanitz, *J. Comput. Chem.* **1992**, 13, 1171.
[5] S. L. Price in *Computer Modelling in Inorganic Crystallography* (Ed.: C. R. A. Catlow), Academic Press, San Diego, **1997**, pp. 269–293.
[6] R. J. Gdanitz in *Theoretical Aspects and Computer Modeling of the Molecular Solid State* (Ed.: A. Gavezzotti), Wiley, New York, **1997**, pp. 185–201.
[7] A. Gavezzotti, *Crystallogr. Rev.* **1998**, 7, 5.
[8] P. Verwer, F. J. J. Leusen, *Rev. Comp. Chem.* **1998**, 12, 327.
[9] G. V. Gridunova, N. G. Furmanova, Yu. T. Struchkov, Z. I. Ezhkova, L. P. Grigor'eva, B. A. Chayanov, *Kristallografiya* **1982**, 27, 267.
[10] E. A. Heath, P. Singh, Y. Ebisuzaki, *Acta Crystallogr. Sect. C* **1992**, 48, 1960.
[11] It was reported in reference [10] that differential scanning calorimetry of some samples of *p*-hydroxybenzoic acid showed two transition peaks (separated by 1 K) around the melting point, which was ascribed to the existence of two different crystalline forms. No solid–solid transition was observed. Form II reported herein is possibly the "high temperature form" mentioned in reference [10], but we can make no further speculation in this regard until a pure phase of form II has been obtained and subjected to detailed thermal analysis.
[12] Microcrystals (for typical crystal size, see reference [13]) of form II of *p*-hydroxybenzoic acid were obtained during preparation of inclusion compounds between deoxycholic acid and *p*-hydroxybenzoic acid, using methanol as solvent. The crystals were very small and of very poor quality.
[13] Single-crystal X-ray diffraction data were recorded for a microcrystal of form II of *p*-hydroxybenzoic acid on Station 9.8 at the Synchrotron Radiation Source (Daresbury Laboratory). This microcrystal diffraction station comprises a Siemens SMART CCD detector and goniometer system. Crystal size $0.10 \times 0.07 \times 0.05$ mm³; $T = 296(2)$ K; $\lambda = 0.68850$ Å; monoclinic, $P2_1/n$; $a = 18.66(2)$, $b = 3.860(4)$, $c = 18.82(3)$ Å, $\beta = 93.511(6)^\circ$; $V = 1353(3)$ Å³; $Z = 8$. The structure was solved by using SHELXS^[14] and refined by using SHELXL^[15] ($R_1 = 0.131$; $R_w = 0.365$). The aryl rings were refined as idealized hexagons (including hydrogen atoms). In the difference Fourier map, the strongest peak ($0.5 \text{ e } \text{\AA}^{-3}$) represented the position of the hydrogen atom of the carboxylic acid group of molecule A (the ordered carboxylic acid group—see text), and this hydrogen atom was then included in the refinement in a fixed position corresponding to standard geometry. The hydrogen atom of the carboxylic acid group of molecule B and the hydrogen atoms of the hydroxy groups were not located among the next highest peaks in the difference Fourier map. As only microcrystals of form II of *p*-hydroxybenzoic acid were available for the X-ray diffraction experiments, it is necessary to recognize that the intrinsically poorer data quality has resulted in relatively high values of R factors. Nevertheless, by maintaining the discussion and interpretation at a level commensurate with the reliability of the refined structural information, reliable conclusions on structural properties are nevertheless reached.
[14] G. M. Sheldrick, SHELXS86, Program for the Solution of Crystal Structures, University of Göttingen, Germany, **1986**.
[15] G. M. Sheldrick, SHELXL92, Program for the Refinement of Crystal Structures, University of Göttingen, Germany, **1993**.
[16] L. Leiserowitz, *Acta Crystallogr. Sect. B* **1976**, 32, 7755.
[17] The graph set notation is defined and applied in the following papers: M. C. Etter, *Acc. Chem. Res.* **1990**, 23, 120; J. Bernstein, R. E. Davis, L. Shimoni, N. L. Chang, *Angew. Chem.* **1995**, 107, 1689; *Angew. Chem. Int. Ed. Engl.* **1995**, 34, 1555.
[18] The geometry of the C(11)–O(6)–H...O(1)–C(7)–C(1) hydrogen bond is: O(6)...O(1) distance 2.946(6) Å; O(6)...O(1)–C(7) angle $139.6(3)^\circ$; C(11)–O(6)...O(1) angle $120.1(2)^\circ$; O(6)...O(1)–C(7)–C(1) dihedral angle $-35.6(6)^\circ$. The O(6)–H...O(1) angle, for the hydrogen atom of the hydroxy group inserted according to standard geometry, is about 165° . Note that C(11)–O(6)–H is the hydroxy group of molecule B and O(1)–C(7) is in the carboxylic acid group of molecule A, and is assigned (see text) as an O=C bond.
[19] G. A. Jeffrey, W. Saenger, *Hydrogen Bonding in Biological Structures*, Springer, Berlin, **1994**.
[20] The geometries of these C(4)–O(3)–H...O(4)–C(14)–C(8) contacts are: a) O(3)...O(4) distance 3.439(9) Å; O(3)...O(4)–C(14) angle

112.2(4)°; C(4)–O(3)···O(4) angle 107.6(3)°; O(3)···O(4)–C(14)–C(8) dihedral angle –60.3(6)°; b) O(3)···O(4) distance 3.427(8) Å; O(3)···O(4)–C(14) angle 153.3(4)°; C(4)–O(3)···O(4) angle 123.3(4)°; O(3)···O(4)–C(14)–C(8) dihedral angle 25.5(1.3)°. Note that C(4)–O(3)–H is the hydroxy group of molecule A and O(4)–C(14) is in the carboxylic acid group of molecule B.

- [21] F. H. Allen, O. Kennard, D. G. Watson, L. Brammer, A. G. Orpen, R. Taylor, *J. Chem. Soc. Perkin Trans. 2* **1987**, S1.
- [22] C. P. Brock, L. L. Duncan, *Chem. Mater.* **1994**, 6, 1307.
- [23] B. H. Meier, F. Graf, R. R. Ernst, *J. Chem. Phys.* **1982**, 76, 767.
- [24] K. Furic, *Chem. Phys. Lett.* **1984**, 108, 518.
- [25] B. H. Meier, R. Meyer, R. R. Ernst, A. Stöckli, A. Furrer, W. Hälg, I. Anderson, *Chem. Phys. Lett.* **1984**, 108, 522.
- [26] S. Nagaoka, T. Terao, F. Imashiro, N. Hirota, S. Hayashi, *Chem. Phys. Lett.* **1984**, 108, 524.
- [27] W. Scheubel, H. Zimmerman, U. Haeberlen, *J. Mag. Res.* **1988**, 80, 401.
- [28] C. C. Wilson, N. Shankland, A. J. Florence, *J. Chem. Soc. Faraday Trans.* **1996**, 92, 5051.
- [29] M. Neumann, D. F. Broughman, C. J. McGloin, M. R. Johnson, A. J. Horsewill, H. P. Trommsdorff, *J. Chem. Phys.* **1998**, 109, 7300.
- [30] M. A. Neumann, S. Craciun, A. Corval, M. R. Johnson, A. J. Horsewill, V. A. Benderskii, H. P. Trommsdorff, *Phys. Chem. Chem. Phys.* **1998**, 102, 317.
- [31] Form II of *p*-hydroxybenzoic acid represents a very rare example of a structure containing two crystallographically independent carboxylic acid dimer motifs in which one motif is ordered and the other is disordered. In support of this statement, we have carried out a search of the Cambridge Structural Database (CSD) to find structures containing both an unambiguously ordered carboxylic acid dimer (i.e. with one C–O bond in each carboxylic acid group in the range 1.21–1.25 Å representing C=O, and the other C–O bond in each carboxylic acid group in the range 1.31–1.35 Å representing C–OH) and an unambiguously disordered carboxylic acid dimer (i.e. with both C–O bonds in each carboxylic acid group in the range 1.26–1.30 Å). We note first that there is no overlap in the ranges used to distinguish the ordered and disordered carboxylic acid dimers, and second that for the structure reported here for form II of *p*-hydroxybenzoic acid, one carboxylic acid dimer is classed as ordered and the other is classed as disordered on the basis of the above distance criteria. In the CSD search, only one structure (octa-*cis*-2,trans-6-diene-1,8-dioic acid^[32]) was found containing carboxylic acid dimers of both types (note that six structures were found containing individual carboxylic acid groups of both types, but it is only in the above structure that these carboxylic acid groups are present in ordered and disordered dimers). However, the estimated standard deviations in the C–O bond lengths in the structure of octa-*cis*-2,trans-6-diene-1,8-dioic acid (which was determined from Weissenberg photographic data) were in the range 0.02–0.04 Å, from which the reliability of the classification on the above distance criteria becomes significantly blurred. We note that these estimated standard deviations are substantially larger than those reported herein.
- [32] E. Martuscelli, L. Mazzarella, *Acta Crystallogr. Sect. B* **1969**, 25, 2022.

Structure Determination of an Oligopeptide Directly from Powder Diffraction Data**

Emilio Tedesco, Giles W. Turner, Kenneth D. M. Harris,* Roy L. Johnston, and Benson M. Kariuki

In recent decades, single-crystal X-ray diffraction techniques have underpinned many major breakthroughs in structural biology, although it is well recognized that an intrinsic limitation of these techniques is the requirement to prepare single crystals of sufficient size, quality, and stability to allow diffraction data of sufficiently high quality to be measured. In solid-state and materials sciences, similar limitations are encountered in the scope and applicability of single-crystal diffraction methods, but much progress has been made in recent years in the use of powder diffraction data as an alternative means of obtaining structural information. Herein, we apply a new methodology for structure determination from powder diffraction data to determine the crystal structure of the oligopeptide Phe-Gly-Gly-Phe (Figure 1), which demonstrates the opportunity for future extensions to more complex systems of interest in structural biology. In this regard, we note that knowledge of the structural properties and interactions in model oligopeptide systems can yield fundamental insights towards understanding structural properties of polypeptide sequences in proteins.^[1–6]




Figure 1. The molecular structure of Phe-Gly-Gly-Phe, showing the variable torsion angles in the genetic algorithm structure solution calculation.

In principle, powder diffraction data contains the same information as single-crystal diffraction data, but with the three-dimensional diffraction data compressed into one dimension. As a consequence, there is generally substantial overlap of peaks in the powder diffraction pattern, leading to severe difficulties in extracting the same quality of diffraction data (and hence structural information). Although the challenges in determining structures directly from powder diffraction data are considerable, new techniques and strat-

[*] Prof. K. D. M. Harris, Dr. E. Tedesco, G. W. Turner, Dr. R. L. Johnston, Dr. B. M. Kariuki
School of Chemistry, University of Birmingham
Edgbaston, Birmingham B15 2TT (UK)
Fax: (+44) 121-414-7473
E-mail: K.D.M.Harris@bham.ac.uk

[**] This work was supported by the EPSRC, the University of Birmingham, Wyeth-Ayerst plc, and Ciba Specialty Chemicals. We are grateful to Professor P. Balaram (Indian Institute of Science, Bangalore) for valuable discussions.

Supporting information for this article is available on the WWW under <http://www.wiley-vch.de/home/angewandte/> or from the author.