


Novel Three-Component Coupling Using Aluminum Tris(2,6-diphenylphenoxide) (ATPH): The Same Synthetic Strategy Leads to *trans*- and *cis*-Jasmonates


Susumu Saito, Satoko Yamazaki, and Hisashi Yamamoto*

One-pot three-component coupling^[1] is an important procedure in organic synthesis and provides a powerful and rapid means for construction of the prostaglandin^[2] and jasmonate^[3] families. Here we report on a novel strategy for three-component coupling (Scheme 1), which involves the




Scheme 1. Three-component coupling using ATPH. TBS = *tert*-butyldimethylsilyl.

combined use of: 1) an organolithium reagent (RLi); 2) aluminum tris(2,6-diphenylphenoxide) (ATPH;^[4] see space-filling model) cyclopentenone complexes **1a** and **1b**; and 3) 2,5-dihydrofuran (DHF)·BCl₃ complex (**2**). This approach leads efficiently to both *trans*- and *cis*-jasmonate derivatives.



Precomplexation of a solution of ATPH (1.05 equiv) in toluene with 2-cyclopenten-1-one (**3a**; 1.0 equiv) at -78°C was followed by treatment with a solution of isopropenyllithium (1.1 equiv) in DME. After 1 h, sequential addition of DHF (5.0 equiv) and BCl₃ (2 equiv) gave three-component

coupling product **4**, which was isolated in a yield of 71 % (2,3-*trans*-**4**/2,3-*cis*-**4** = 3.5:1; 2'*E*/2'*Z* = 1:99). The formation of **4** was not observed before treatment with BCl₃.^[5] The addition of a mixture of DHF and BCl₃ gave a yield and selectivity comparable to those achieved by sequential addition. Other alkenyl- and alkynyllithiums, as well as lithium enolates, are well suited to this process and give the corresponding products in moderate to high yields with complete retention of the original *Z* configuration (Table 1). The even more labile cyclopentenone **3b** was also compatible with these basic conditions (entries 3 and 6), but scant 2,3-*cis/trans* selectivity was observed with 2-alkylethenyllithiums (entries 2 and 3). The 2,3-*cis* selectivity increases in the order isopropenyl- < crotyl- < alkynyllithium, that is, with decreasing size of the organolithium reagents (each prepared in DME) (entries 1, 2, and 5).^[6]



Tetrahydrofuran was also used as an alkylating agent in place of DHF (Table 2), but in some cases the stereoselectivity was different from that with DHF. For instance, in the case of isopropenyllithium, the 2,3-*cis* product predominated with THF (Table 2, entries 2–4),^[6, 7] while the 2,3-*trans* product was formed selectively with DHF (Table 1, entries 4–6). However, both DHF and THF gave the 2,3-*cis* product almost exclusively with 1-hexynyllithium (Table 1, entry 1; Table 2, entry 1).^[6, 7] The TBSOTf·THF complex gives better 2,3-*cis* selectivity than BCl₃·THF (Table 2, entries 2–4). Both THF and DHF showed a general trend to give high 2,3-*trans* selectivity with lithium enolates (Table 1, entries 8 and 9; Table 2, entries 5–7).^[8]

With this new method in hand, various *trans*- and *cis*-jasmonate derivatives can be obtained in a few steps by assembling commercially available compounds. For example, stereo- and regioselective methylation of diphenylphosphate **8** (Scheme 2), readily obtained from *trans*-**7**, was effected by using trimethylzincate and a catalytic amount of Fe^{III} (Me₂Zn + MeMgBr (2 + 2 equiv); [Fe(acac)₃]: 1 mol %)^[9] at -78°C to give *tert*-butyl jasmonate **9** (2,3-*trans/cis* > 20:1) together with small amounts of stereo- (2'*Z*/2'*E* = 97:3; **10**, 2 %) and regioisomers (S_N2/S_N2' = 87:13; **11**, 11 %); the total yield of isomers was 87 %, and the isomeric purity of **9** was 85 %).^[10] The ketone and ester groups remained intact under these conditions. Subsequent displacement of a *tert*-butyl group (HCO₂H, RT) led to quantitative formation of *trans*-jasmonic acid.

Likewise, methyl *cis*-jasmonate was easily accessible from three-component coupling product **12** (2,3-*cis/trans* = 7.5:1 and 71 % yield from **3a**). After separation of the diastereomers (to 2,3-*cis/trans* = 99:1), a similar phosphorylation–methylation^[9] sequence (Scheme 3) caused only slight epimerization at C2 and gave **13** (2,3-*cis/trans* = 31:1) in a reasonable overall yield together with moderate amounts of stereo-
















[*] Prof. Dr. H. Yamamoto, Dr. S. Saito, S. Yamazaki
 Graduate School of Engineering, Nagoya University, SORST
 Japan Science and Technology Corporation (JST), Chikusa
 Nagoya 464-8603 (Japan)
 Fax: (+81) 52-789-3222
 E-mail: yamamoto@cc.nagoya-u.ac.jp

Table 1. Three-component coupling of RLi, **1a** (or **1b**), and DHF·BCl₃ complex.^[a]


3a: X = H 3b: X = OTBS					
Entry	RLi [equiv] ^[b]	Ketone	Conditions [°C, h] ^[c, d]	Products	Yield [%] (2,3- <i>trans</i> /2,3- <i>cis</i>) ^[e]
1	(1.1)	3a	−78, 2.5; −40, 2		44 (1:11)
2	(1.1)	3a	−78, 5; −40, 0.5		57 (1.2:1)
3	(1.2)	3b	−78, 20; −40, 10		46 (1:1) ^[f]
4	(1.1)	3a	−78, 17.5; −40, 3		59 (4.4:1)
5	(1.1)	3a	−78, 1.5; −40, 3		71 (3.5:1)
6	(1.1)	3b	−78, 168		66 (79:11:10) ^[g]
7	(2.0)	3a	−78, 18; −40, 2		69 (>20:1) ^[h]
8	(2.0)	3a	−78, 19		47 (>20:1)
9	(2.0)	3a	−78, 19; −40, 3		32 (>20:1)

[a] Unless otherwise specified, the reaction was performed with a solution of ATPH (1.05 equiv) in toluene, cyclopentenone (1.0 equiv), organolithium (RLi), BCl₃ or *tert*-butyldimethylsilyl triflate (TBSOTf), and DHF. [b] Each was prepared from the corresponding alkyne, bromide, or tin reagent with BuLi (entries 1–6) or from carbonyl compounds and lithium diisopropylamide (LDA; entries 7–9) in DME (entries 1–3 and 5) or diethyl ether (entries 4 and 6–9). [c] The following conditions were used: BCl₃ (2.0 equiv) + DHF (5.0 equiv). [d] The indicated reaction temperatures and times are for the three-component coupling process. [e] Yields of isolated, purified product. The diastereoselectivity was determined by ¹H and ¹³C NMR spectroscopy. See also ref. [7]. [f] Elimination product **5** (7%) was also obtained. [g] 2,3-*trans*,3,4-*trans*/2,3-*cis*,3,4-*trans*/2,3-*trans*,3,4-*cis*. [h] Protonated product **6** (27%) was obtained.

Table 2. Three-component coupling of RLi, ATPH·3a (1a), and THF·Lewis acid.^[a]

					
Entry	RLi [equiv] ^[b]	Alkylating agent ^[c]	Conditions [°C, h] ^[d]	Products	Yield [%] (2,3- <i>trans</i> /2,3- <i>cis</i>) ^[e]
1	 (1.1)	A	−78, 2.5; −40, 2		44 (1: >20)
2	 (1.1)	A ^[f]	−78, 7.5		77 (1:2.7)
3	 (1.1)	A	−78, 2		85 (1:3.2)
4	 (1.1)	B	−78, 2; −40, 2.5		74 (1:4.0)
5	 (2.0)	A	−78, 15		43 (>20:1)
6	 (2.0)	A	−78, 15		54 (>20:1)
7	 (2.0)	A	−78, 15		57 (>20:1)


[a] Unless otherwise specified, the reaction was performed with a solution of ATPH (1.05 equiv) in toluene, cyclopentenone (1.0 equiv), organolithium (RLi, THF solution), and BCl₃ or TBSOTf. [b] Each was prepared from the corresponding alkyne, bromide, or tin reagent with BuLi (entries 1–4). Enolates were prepared by using LDA (entries 5–7). [c] The following conditions were used: A) BCl₃ (2.0 equiv) + THF (123 equiv); B) TBSOTf (7.0 equiv) + THF (123 equiv). Organolithiums or enolates were prepared in THF. [d] The indicated reaction temperatures and times are for the three-component coupling process. [e] Yields of isolated, purified products. The diastereoselectivity was determined by ¹H and ¹³C NMR spectroscopy. See also ref. [7]. [f] BCl₃ (1.1 equiv) was used.



Scheme 2. Synthesis of *trans*-jasmonic acid. Reagents and conditions: a) Et₃N, ClPO(OPh)₂, DMAP, CH₂Cl₂, −20°C, 90%; b) Me₃ZnMgBr, [Fe(acac)₃] (1 mol %), THF, −78°C, <87%; c) HCO₂H, RT, >99%. DMAP = 4-(dimethylamino)pyridine, acac = acetyl acetone.

(2'Z/2'E = 90:10; **14**, 7%) and regioisomers (S_N2/S_N2' = 84:16; **15**, 14%); the total yield of isomers was 86%, and the isomeric purity of **13** was 73%.^[10] Reduction of the ketone carbonyl group was followed by hydroboration of the trimethylsilylacetylene^[11] and subsequent oxidation. The resulting crude carboxylic acid was selectively methylated, and reoxidation of the remaining hydroxy group would lead to methyl *cis*(*epi*)-jasmonate^[12] by the method previously described.^[13]

In summary, the present coupling was successful under strongly basic conditions, which is



Scheme 3. Synthesis of methyl *epi(cis)*-jasmonate. Reagents and conditions: a) Et_3N , $\text{ClPO}(\text{OPh})_2$, -20°C , 84%; b) Me_3ZnMgBr , $[\text{Fe}(\text{acac})_3]$ (5 mol %), THF, -78°C , <86%; c) NaBH_4 , MeOH, 0°C , 91% (1,2-*cis*;2,3-*cis*/1,2-*trans*;2,3-*cis* = 3.2:1); d) $(\text{C}_6\text{H}_{11})_2\text{BH}$, THF, 0°C ; H_2O_2 , NaOH, MeOH/benzene, 0°C ; TMSCHN_2 , MeOH, 0°C , 48%; e) see ref. [12]. TMS = trimethylsilyl.

otherwise difficult to achieve. The synthetic potential of ATPH may originate from: 1) The effective Michael addition of organolithium reagents due to both steric (prevention of 1,2-addition and α - and/or γ -deprotonation) and electronic (activation of the β -position) influence of ATPH;^[4e] and 2) the novel reactivity of the resulting ATPH enolates, which undergo either 2,3-*cis*- or -*trans*-selective alkylation by the DHF/Lewis acid complexes. This eventually accommodated the *Z* geometry of the new double bond in the α chain.

Experimental Section

Representative procedure for the reaction of ATPH·**3a**, isopropenyllithium, and BCl_3 ·DHF: **3a** (42 μL , 0.5 mmol) was added to a solution of ATPH (0.53 mmol) in toluene (5.0 mL) at -78°C under an argon atmosphere. After 20 min, isopropenyllithium [generated by treatment of a solution of isopropenyltributyltin (194 mg, 0.59 mmol) in DME (5.0 mL) with a solution of *n*BuLi (1.66 M, 0.33 mL, 0.55 mmol) in hexane at -78°C followed by stirring at 0°C for 12 min and cooling to -78°C] was transferred by cannula to the solution of ATPH·**3a** at -78°C . The reaction mixture was stirred at this temperature for 45 min. To this mixture were then added DHF (0.189 mL, 2.5 mmol) and a solution of BCl_3 (1.0 M, 1.0 mL, 1.0 mmol) in hexane. After 1.5 h, the reaction mixture was warmed to -40°C , stirred for 3 h, quenched with aqueous NaHCO_3 , filtered through a Celite pad, and extracted with Et_2O . The organic layer was dried over Na_2SO_4 , and concentrated. The residue was purified by column chromatography on silica gel to give **4** (69.4 mg, 71%) as a pale yellow oil. **trans-4**: IR (neat): $\tilde{\nu}$ = 3434, 2950, 1734, 1456, 1377, 1148, 1015, 895 cm^{-1} ; ^1H NMR (CDCl_3 , 300 MHz): δ = 5.74–5.63 (m, 1H), 5.58–5.41 (m, 1H), 4.87 (dt, 2H, J = 12.3, 1.2 Hz), 4.16 (dd, 2H, J = 6.6, 2.1 Hz), 2.57–2.03 (m, 8H), 1.76 (s, 3H), 1.78–1.63 (m, 1H); ^{13}C NMR (CDCl_3 , 75 MHz): δ = 219.8, 144.9, 130.5, 128.6, 112.0, 58.0, 52.1, 48.8, 37.8, 26.4, 24.5, 19.0; elemental analysis (%) calcd for $\text{C}_{12}\text{H}_{18}\text{O}_2$: C, 74.19; H, 9.34; found: C, 74.09; H, 9.64. **cis-4**: ^1H NMR (CDCl_3 , 300 MHz): δ = 4.90 (s, 1H), 4.72 (s, 1H), 4.18–4.04 (m, 2H), 2.97–2.90 (m, 1H), 1.70 (s, 3H); ^{13}C NMR (CDCl_3 , 75 MHz): δ = 219.8, 144.2, 130.1, 129.8, 112.4, 58.1, 51.7, 46.4, 36.4, 24.5, 23.4, 22.4.

Received: June 7, 2001 [Z17236]

[1] a) R. Noyori, M. Suzuki, *Angew. Chem.* **1984**, 96, 854; *Angew. Chem. Int. Ed. Engl.* **1984**, 23, 847; b) P. W. Collins, S. W. Djuric, *Chem. Rev.*

- 1993**, 93, 1533; c) B. H. Lipshutz, M. R. Wood, *J. Am. Chem. Soc.* **1994**, 116, 11689; for reviews on domino reactions and multicomponent coupling, see d) L. F. Tietze, U. Beifuss, *Angew. Chem.* **1993**, 105, 137; *Angew. Chem. Int. Ed.* **1993**, 32, 131; e) A. Dömling, I. Ugi, *Angew. Chem.* **2000**, 112, 3300; *Angew. Chem. Int. Ed.* **2000**, 39, 3168.
- [2] a) M. Suzuki, A. Yanagisawa, R. Noyori, *J. Am. Chem. Soc.* **1985**, 107, 3348; b) M. Suzuki, Y. Morita, H. Koyano, M. Koga, R. Noyori, *Tetrahedron* **1990**, 46, 4809; c) S. Chen, K. D. Janda, *J. Am. Chem. Soc.* **1997**, 119, 8724; d) A. Fürstner, K. Grela, C. Mathes, C. W. Lehmann, *J. Am. Chem. Soc.* **2000**, 122, 11799; e) L. A. Arnold, R. Naasz, A. J. Minnaard, B. L. Feringa, *J. Am. Chem. Soc.* **2001**, 123, 5841. For a 4-oxa-prostaglandin derivative (ZK118182), see: f) S. Okamoto, S. Katayama, N. Ono, F. Sato, *Tetrahedron: Asymmetry* **1992**, 3, 1525.
- [3] a) T. K. Sarker, B. K. Ghorai, *J. Indian Chem. Soc.* **1999**, 76, 693, and references therein; b) F.-T. Luo, E. Negishi, *Tetrahedron Lett.* **1985**, 26, 2177; c) W. Oppolzer, M. Guo, K. Baettig, *Helv. Chim. Acta* **1983**, 66, 2140; d) H. Gerlach, P. Kunzler, *Helv. Chim. Acta* **1978**, 61, 2503; two industrial syntheses: e) H. Kataoka, T. Yamada, K. Goto, J. Tsuji, *Tetrahedron* **1987**, 43, 4107; f) F. Näf, R. Decorzant, *Helv. Chim. Acta* **1978**, 61, 2524.
- [4] For review, see: a) S. Saito, H. Yamamoto, *Chem. Commun.* **1997**, 1585; b) S. Saito, H. Yamamoto, *Chem. Eur. J.* **1999**, 5, 1959; c) S. Saito, M. Shiozawa, T. Nagahara, M. Nakadai, H. Yamamoto, *J. Am. Chem. Soc.* **2000**, 122, 7847; d) S. Saito, T. Sone, M. Murase, H. Yamamoto, *J. Am. Chem. Soc.* **2000**, 122, 10216; for Michael addition to α,β -unsaturated carbonyl compounds, see: e) K. Maruoka, H. Imoto, S. Saito, H. Yamamoto, *J. Am. Chem. Soc.* **1994**, 116, 4131; for novel reactivity and selectivity in the formation and reaction of ATPH enolate complexes, see: f) S. Saito, S. Yamazaki, M. Shiozawa, H. Yamamoto, *Synlett* **1999**, 581; g) S. Saito, M. Shiozawa, H. Yamamoto, *Angew. Chem.* **1999**, 111, 1884; *Angew. Chem. Int. Ed.* **1999**, 38, 1769; h) S. Saito, M. Shiozawa, M. Ito, *J. Am. Chem. Soc.* **1998**, 120, 813; i) S. Saito, M. Ito, H. Yamamoto, *J. Am. Chem. Soc.* **1997**, 119, 611; j) S. Saito, M. Ito, K. Maruoka, H. Yamamoto, *Synlett* **1997**, 357; k) K. Maruoka, I. Shimada, H. Imoto, H. Yamamoto, *Synlett* **1994**, 519.
- [5] This means that neither Li^+ nor Al^{III} is the activating agent for DHF.
- [6] At present, we do not have a reasonable explanation for these unprecedented 2,3-*cis* selectivities, which might be ascribed to steric influence of ATPH.
- [7] The diastereoselectivity (2,3-*cis*:2,3-*trans*) of three-component coupling products was determined by ^1H and ^{13}C NMR spectroscopy, as well as several *cis* and *trans* equilibration experiments (HCl, MeOH, RT). We found that ^{13}C NMR spectroscopy could be a powerful tool for establishing the relative stereochemistry at the 2,3-vicinal carbon atoms of 2,3-disubstituted-1-cyclopentanones. In general, ^{13}C NMR chemical shifts of $\text{C}1'$ and $\text{C}1''$ of the *cis* adducts appear upfield due to the greater shielding effects compared to those in the *trans* products. For example, *cis*- and *trans*-2,3-dimethylcyclopentanones show δ = 9.5 (2-Me ($\text{C}1'$)) and 14.8 (3-Me ($\text{C}1''$)), and δ = (2-Me ($\text{C}1'$)) and 19.1 (3-Me ($\text{C}1''$)), respectively (J. B. Stothers, C. T. Tan, *Can. J. Chem.* **1974**, 52, 308); methyl *epi*-jasmonate, *cis*: δ = 23.0 (2- CH_2 ($\text{C}1'$)), 33.7 (3- CH_2 ($\text{C}1''$)); *trans*: δ = 25.6 (2- CH_2 ($\text{C}1'$)), 37.7 (3- CH_2 ($\text{C}1''$)) (L. Crombie, K. M. Mistry, *J. Chem. Soc. Perkin Trans. 1* **1991**, 1981). We also found that the equilibrium reached an inherent static point depending on the size of the 3-substituent. Thus, the equilibrium lies on the *trans* side in the order of bulkiness: isopropenyl (*cis:trans* = 1:25) > crotyl (*cis:trans* = 1:11) > 1-hexynyl (*cis:trans* = 1:1.6).
- [8] In general, the concomitant formation of Michael addition products such as **6** through enolate protonation was a major side reaction. Considerable formation of this type of product (ca. 30%) was observed in several cases (Table 1, entries 8, and 9; Table 2, entries 1, and 5–7). It is highly unlikely that the *trans* selectivity is due to the work-up procedure, since the same conditions were used throughout the work (see Experimental Section).
- [9] The details of this new stereo- and regioselective method for the alkylation of allylic phosphates by the combined use of trialkyl- or tetraalkylzincates and Fe^{III} or Fe^I catalysts will be reported elsewhere.
- [10] These selectivities ($\text{S}_{\text{N}}2$ vs $\text{S}_{\text{N}}2'$; 2'E vs 2'Z; 2,3-*cis* vs 2,3-*trans*) were determined by ^1H and ^{13}C NMR spectroscopy and GC-MS.
- [11] J. A. Miller, G. Zweifel, *Synthesis* **1981**, 288.
- [12] Perhaps the most efficient asymmetric synthesis of methyl *cis*-jasmonate reported so far was reported by C. Fehr, J. Galindo,

Angew. Chem. 2000, 112, 581; Angew. Chem. Int. Ed. 2000, 39, 569. Other approaches to methyl *cis*-jasmonate are cited therein. See also ref. [3a].


- [13] a) T. K. Sarkar, B. Mukherjee, S. K. Ghosh, *Tetrahedron* 1998, 54, 3243; b) H. Stadtmüller, P. Knochel, *Synlett* 1995, 463; c) L. Crombie, K. M. Mistry, *J. Chem. Soc. Perkin Trans. 1* 1991, 1981.

The First Platinum Formyl, a Member of a Series of Hexanuclear Clusters Exhibiting a Rare Structure**

Piero Leoni,* Fabio Marchetti, Lorella Marchetti, Marco Pasquali, and Silvia Quaglierini

A large number of platinum clusters have been reported in the past decades.^[1] Nevertheless, only two hexanuclear structures have been authenticated by X-ray crystallography: the trigonal-prismatic $[\text{Pt}_6(\mu\text{-CO})_6(\text{CO})_6]^{2-}$ ^[2a] and the octahedral $[\text{Pt}_6(\text{CO})_6(\mu\text{-dppm})_3]^{2+}$ (dppm = bis(diphenylphosphanyl)methane).^[2b] In addition, the decanuclear $[\text{Fe}_4\text{Pt}_6(\text{CO})_{22}]^{2-}$,^[3] arising from the orthogonal dimerization of two $[\text{Pt}_3(\text{CO})_3\{\mu_2\text{-Fe}(\text{CO})_4\}_2]^-$ triangles, contains a tetrahedral Pt core, two opposite edges of which are each bridged by another platinum atom. We have discovered a facile route to platinum clusters having this unusual structure.^[4] The precursor $[\text{Pt}_6(\text{CO})_6(\mu\text{-PtBu}_2)_4](\text{CF}_3\text{SO}_3)_2$ (**2**) was prepared by treating the trinuclear hydrido cluster $[\text{Pt}_3(\text{H})(\text{CO})_2(\mu\text{-PtBu}_2)_3]$ (**1**)^[5] with an excess of triflic acid under a CO atmosphere (Scheme 1).

Figure 1 shows an ORTEP view of the cation **2**²⁺, which exhibits overall D_{2d} symmetry.^[7] Complex **2** is air-stable, thermally robust (decomp 228 °C), and remains unchanged, even under forcing conditions, in the presence of CO (100 °C, 100 atm, 12 h) or H₂ (80 °C, 90 atm, 6 h). Interestingly, **2** reacts with ¹³CO to give $[\text{Pt}_6(\text{CO})_4(^{13}\text{CO})_2(\mu\text{-PtBu}_2)_4](\text{CF}_3\text{SO}_3)_2$ (**2***), the product of selective substitution of the carbonyl ligands attached to the two “apical” atoms Pt2 and Pt5. A similar remarkable regioselectivity was also observed in the reactions of **2** with PMe₃ or NaBH₄ to give $[\text{Pt}_6(\text{CO})_4(\text{PMe}_3)_2(\mu\text{-PtBu}_2)_4](\text{CF}_3\text{SO}_3)_2$ (**3**), and $[\text{Pt}_6(\text{CO})_4(\text{CHO})_2(\mu\text{-PtBu}_2)_4]$ (**4**), respectively. The structures of **2***, **3**, and **4** can be clearly assigned on the basis of elemental analyses and IR and NMR spectra. For example, the four bridging P nuclei are still



Scheme 1. Synthesis and further reactions of **2**.




Figure 1. Molecular structure of **2**²⁺.^[7] Methyl groups are omitted for clarity (thermal ellipsoids at 30% probability; symmetry code: ' = x, -y, z). Selected bond lengths [Å] and angles [°]: Pt1-Pt2 2.758(1), Pt1-Pt3 2.667(1), Pt1-Pt4 2.841(1), Pt2-Pt3 2.762(1), Pt3-Pt4 2.841(1), Pt4-Pt4' 2.666(1), Pt4-Pt5 2.756(1); Pt3-Pt1-Pt2 61.17(3), Pt1-Pt2-Pt3 57.79(3), Pt1-Pt3-Pt2 61.04(3), Pt3-Pt1-Pt4 62.00(3), Pt1-Pt3-Pt4 62.00(3), Pt1-Pt4-Pt3 56.00(3), Pt1-Pt4-Pt4' 62.01(2), Pt3-Pt4-Pt4' 62.01(2), Pt4'-Pt4-Pt5 61.071(15), Pt4-Pt5-Pt4' 57.86(3).

isochronous and in the ³¹P{¹H} NMR spectrum they give rise to a central singlet (doublet in **3**, ²J(P_μ, P_{term}) = 13 Hz) flanked by a complex set of ¹⁹⁵Pt satellites, very similar to those observed for **2**. Strong similarities were also observed in the main features of the ¹⁹⁵Pt{¹H} NMR spectra, which still show only two signals and hence rule out all other possible isomers.

Complex **4** is of particular interest, being the first platinum formyl, and its structure was unequivocally confirmed by comparison of its spectra with those of the labeled $[\text{Pt}_6(\text{CO})_4(^{13}\text{CHO})_2(\mu\text{-PtBu}_2)_4]$ (**4***). In particular, the triplet for the formyl group at δ = 18.9 (³J(H,P) = 14 Hz) in the ¹H NMR spectrum of **4** splits as expected in **4*** (brdt, ¹J(C,H) = 152 Hz) for coupling with the adjacent ¹³C nucleus (Figure 2a, b). The same coupling is exhibited by a doublet at δ = 244.6 for the formyl C atom in the proton-coupled ¹³C NMR spectrum of **4*** (Figure 2c, d).^[6a] Furthermore, a

[*] Prof. P. Leoni, Dr. L. Marchetti, Prof. M. Pasquali, Dr. S. Quaglierini
Dipartimento di Chimica e Chimica Industriale
Via Risorgimento 35, 56126 Pisa (Italy)
Fax: (+390) 50-20237
E-mail: leoni@ccci.unipi.it

Prof. F. Marchetti
Dipartimento di Ingegneria Chimica dei Materiali
delle Materie Prime e Metallurgia
Via del Castro Laurenziano 7, 00161 Roma (Italy)

[**] We thank the Ministero dell'Università e della Ricerca Scientifica e Tecnologica (MURST), Programmi di Interesse Nazionale, 2000–2001, for financial support.

Supporting information for this article is available on the WWW under <http://www.angewandte.com> or from the author.