

Methyleneimine $\text{CH}_2=\text{NH}$ as a Unidentate Ligand in Rhenium Complexes**


Gabriele Albertin,* Stefano Antoniutti, Alessia Bacchi, Emilio Bordignon, Maria Teresa Giorgi, and Giancarlo Pelizzi

Coordinated hydrazines RNHNH_2 are reported to react with oxidizing agents, such as $[\text{Pb}(\text{OAc})_4]$ and H_2O_2 , to give the corresponding diazenes $\text{RN}=\text{NH}$, the stabilization of which on an appropriate metal fragment allows their separation as coordinated species.^[1–3] We now report a new reaction of coordinated methylhydrazine, which reacts with $[\text{Pb}(\text{OAc})_4]$ to give a $\eta^1\text{-NH}=\text{CH}_2$ methyleneimine derivative.

The $\text{CH}_2=\text{NH}$ molecule is a reactive species which was first obtained in 1933 from the low-temperature reaction of HCN with hydrogen.^[4] It has been detected in several galactic objects^[5] and proposed as a possible precursor^[6] of the simplest α -amino acid, glycine. As a ligand, it is present in only one case, through π coordination^[7] to an osmium center; no other report has been found on this molecule, which displays a simple constitution and structure, and has still unknown properties.

The reaction of the hydride^[8] $[\text{ReH}(\text{CO})\{\text{P}(\text{OEt})_3\}_4]$ with triflic acid (TfOH) gives the thermally unstable $[\text{Re}(\eta^2\text{-H}_2)(\text{CO})\{\text{P}(\text{OEt})_3\}_4]^+(\text{CF}_3\text{SO}_3)^-$ species, which loses H_2 , affording the compound $[\text{Re}(\kappa^1\text{-OTf})(\text{CO})\{\text{P}(\text{OEt})_3\}_4]$. Substitution of the weakly bound triflate ligand with methylhydrazine gives *trans*- $[\text{Re}(\text{CH}_3\text{NHNH}_2)(\text{CO})\{\text{P}(\text{OEt})_3\}_4]^+$ (**1**), which was isolated as a BPh_4 salt (**1-BPh₄**) in about 70% yield (Scheme 1).

Complex **1-BPh₄** was characterized by standard methods (IR, NMR, Λ_M , elemental analysis). The IR spectra show the ν_{NH} bands at 3343 and 3291 cm^{-1} of the methylhydrazine



Scheme 1. $\text{P} = \text{P}(\text{OEt})_3$.

[*] Prof. G. Albertin, S. Antoniutti, E. Bordignon, Dr. M. T. Giorgi
Dipartimento di Chimica
Università Ca' Foscari di Venezia
Dorsoduro 2137, 30123 Venezia (Italy)
Fax: (+39)041-234-8917
E-mail: albertin@unive.it


Prof. A. Bacchi, G. Pelizzi
Dipartimento di Chimica Generale ed Inorganica, Chimica Analitica
Chimica Fisica, Università di Parma
Parco Area delle Scienze 17/a, 43100 Parma (Italy)

[**] This work was supported by MIUR (Rome)—Programmi di Ricerca Scientifica di Rilevante Interesse Nazionale, Cofinanziamento 2000–2001. We thank Daniela Baldan for technical assistance.

Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.

ligand, whereas the ^1H NMR spectrum exhibits resonance signals at $\delta = 4.35$ (s, br; $\text{ReNH}_2\text{NHCH}_3$), 3.93 (m, br; $\text{ReNH}_2\text{NHCH}_3$), and 2.49 ppm (d; $\text{ReNH}_2\text{NHCH}_3$) of the CH_3NHNH_2 group.

Treatment of methylhydrazine complex **1-BPh₄** with an equimolar amount of $[\text{Pb}(\text{OAc})_4]$ at low temperature (-40°C) in CH_2Cl_2 gives a mixture of methyldiazene $[\text{Re}(\text{CH}_3\text{N}=\text{NH})(\text{CO})\{\text{P}(\text{OEt})_3\}_4]\text{BPh}_4$ (**2-BPh₄**) and methyleneimine $[\text{Re}(\eta^1\text{-NH}=\text{CH}_2)(\text{CO})\{\text{P}(\text{OEt})_3\}_4]\text{BPh}_4$ (**3-BPh₄**) derivatives (Scheme 2). These were separated by fractional crystallization in moderate yields (42% for **2-BPh₄**, 24% for **3-BPh₄**) as analytically pure white crystalline solids.



Scheme 2. $\text{P} = \text{P}(\text{OEt})_3$.

The complexes were characterized by spectroscopy and in two X-ray diffraction studies.^[9–12] Figure 1 shows the crystal structure of the cation $[\text{Re}(\eta^1\text{-NH}=\text{CH}_2)(\text{CO})\{\text{P}(\text{OEt})_3\}_4]^+$ (**3**). The most relevant feature of the complex is the presence




Figure 1. Structure of the core of the cation **3** (thermal ellipsoids drawn at the 30% level; ethoxy groups are omitted for clarity). Selected bond lengths [$\text{\AA}$] and angles [$^\circ$]: $\text{Re}-\text{C}50$ 1.956(8), $\text{Re}-\text{N}1$ 2.32(1), $\text{Re}-\text{P}3$ 2.362(2), $\text{Re}-\text{P}1$ 2.362(2), $\text{Re}-\text{P}4$ 2.374(2), $\text{Re}-\text{P}2$ 2.378(2), $\text{O}13-\text{C}50$ 1.108(8), $\text{N}1-\text{C}51$ 1.26(1), $\text{C}50-\text{Re}-\text{N}1$ 175.9(3), $\text{C}50-\text{Re}-\text{P}3$ 87.7(2), $\text{N}1-\text{Re}-\text{P}3$ 95.9(2), $\text{C}50-\text{Re}-\text{P}1$ 86.9(2), $\text{N}1-\text{Re}-\text{P}1$ 89.5(2), $\text{P}3-\text{Re}-\text{P}1$ 174.44(6), $\text{C}50-\text{Re}-\text{P}4$ 94.4(2), $\text{N}1-\text{Re}-\text{P}4$ 87.6(2), $\text{P}3-\text{Re}-\text{P}4$ 90.00(7), $\text{P}1-\text{Re}-\text{P}4$ 89.25(7), $\text{C}50-\text{Re}-\text{P}2$ 92.9(2), $\text{N}1-\text{Re}-\text{P}2$ 85.0(2), $\text{P}3-\text{Re}-\text{P}2$ 91.44(6), $\text{P}1-\text{Re}-\text{P}2$ 90.01(6), $\text{P}4-\text{Re}-\text{P}2$ 172.56(6), $\text{O}13-\text{C}50-\text{Re}$ 177.2(6), $\text{C}51-\text{N}1-\text{Re}$ 134(1).

of the methyleneimine ligand, *trans* to the carbonyl group, and coordinated with the metal in a bent mode, as required by the sp^2 character of the N atom ($\text{Re}-\text{N}-\text{C}$ 139(1) $^\circ$), with $\text{Re}-\text{N}$ 2.32(1) and $\text{N}-\text{C}$ 1.26(1) $\text{\AA}$. This is, in fact, the first example of η^1 coordination of a $\text{CH}_2=\text{NH}$ molecule to a transition metal, the only other similar case being the deprotonated $\text{CH}_2=\text{N}=\text{M}$ fragment found in (μ^2 -methyleneamido)tricarbonylbis-

(η^5 -pentamethylcyclopentadienyl)methyleneamidodimolybdenum,^[13] in which the system is practically linear (M–N–C 163°). The bent geometry found for our terminal methyleneimine group fits the common structural features of alkylic and aryl $R_2C=NH$ ligands, which show similar M–N–C angles and generally larger N–C distances (ranging from 1.25 to 1.30 Å; the shortest ones are found in the catenabis(isopropylideneamine)gold trifluoromethanesulfonate complex at 173 K^[14]). The plane of the methyleneimine ligand (Re–N1–C51) forms a dihedral angle of 38(1)° with the equatorial coordination plane containing the N donor (Re–C50–P1–P3–N1).

The ¹H NMR spectra of **3**–BPh₄ are diagnostic for the presence of the methyleneimine ligand, showing a broad high-frequency signal at δ = 13.98 ppm, which is attributed to the =NH imine proton. Substituted imine $R_2C=NH$, and $RHC=NH$ bonded to a metal center^[14, 15] are also reported to give rise to a high-frequency NH proton resonance signal. A slightly broad multiplet is also present at δ = 3.66 ppm, which is coupled with the imine proton and was assigned to one of the two protons of the methylene =CH₂ group. The other is probably masked by the methylene signals of the P(OCH₂CH₃)₃ ligands. In the temperature range between +30 and –80 °C the ³¹P{¹H} NMR spectrum displays a sharp singlet, which is assigned to a *trans* geometry like that found in the solid state.

In the crystal structure of **2**–BPh₄, the methyldiazene and carbonyl ligands in the cation are exchanged between two *trans* coordination positions, with 50 % substitutional disorder, and their refinement was possible only by restraining them to conform to a plausible geometry.^[16]

The ¹H NMR spectra of **2**–BPh₄ further support the presence of the CH₃N=NH ligand, showing the NH resonance signal at δ = 15.99 ppm and one doublet at δ = 4.37 ppm, attributed to the methyl group. A mutual *trans* position of carbonyl and methyldiazene ligands is also suggested in solution by the presence of only one singlet at δ = 116.7 ppm in the ³¹P{¹H} NMR spectrum.

Other methylhydrazine complexes, such as dicarbonyls [Re(CH₃NHNH₂)(CO)₂P₃]BPh₄ (P = P(OEt)₃ or PPh(OEt)₂), were prepared, and their reaction with [Pb(OAc)₄] led, at –40 °C, to a mixture of methyldiazene [Re(CH₃N=NH)(CO)₂P₃]⁺ and methyleneimine [Re(η^1 -NH=CH₂)(CO)₂P₃]⁺ derivatives which, in the case of P(OEt)₃, were separated in pure form or, for PPh(OEt)₂, were detected by spectroscopy. The reaction affording the coordinated η^1 -NH=CH₂ molecule seems to be general for the [Re(CO)_nP_{5–n}] (*n* = 1, 2) fragment containing a methylhydrazine ligand, but appears to be specific for [Pb(OAc)₄], as attempts to carry out the reaction with other oxidants such as MnO₂ or H₂O₂ were unsuccessful.

The formation of species **2**–BPh₄ and **3**–BPh₄ from the reaction of methylhydrazine complexes **1**–BPh₄ (Scheme 2) suggests that [Pb(OAc)₄] gives rise to two parallel reactions involving selective oxidation of CH₃NHNH₂ to methyldiazene CH₃N=NH, giving **2**–BPh₄, in one case, whereas a completely new reaction involving cleavage of the N=N bond and formation of the CH₂=NH moiety takes place in the other. Although coordinated hydrazine is known to undergo oxidation by [Pb(OAc)₄] or other reagents to the corresponding diazene,^[1–3] the reaction affording coordinated η^1 -NH=CH₂ is

new, unexpected, and interesting—not only because it allows us to prepare, and stabilize by coordination, an elusive molecule such as methyleneimine, but also because, whatever the mechanism^[17] may be, cleavage of the N=N bond^[18] of a coordinated hydrazine^[19] takes place in the presence of an oxidizing species.

Studies are currently in progress to explore the reaction chemistry of the M–NH=CH₂ systems, mainly in terms of deprotonation and substitution reactions.

Experimental Section

All reactions were carried out under an inert atmosphere using dry, air-free solvents.

1–BPh₄: CF₃SO₃H (TfOH) (0.23 mmol, 20 μ L) was added to a solution of [ReH(CO){P(OEt)₃}]^[8] (200 mg, 0.23 mmol) in CH₂Cl₂ (5 mL) cooled to –196 °C, and the reaction mixture was allowed to warm to room temperature, and stirred for 1 h. CH₃NHNH₂ (0.6 mmol, 32 μ L) was added and stirring was continued for 24 h. The solvent was removed under reduced pressure to give an oil which was triturated with ethanol (3 mL) containing NaBPh₄ (0.6 mmol, 205 mg). A white solid slowly separated out, which was crystallized from CH₂Cl₂ and ethanol to give **1**–BPh₄ (210 mg; yield 73 %). IR (KBr): $\tilde{\nu}$ = 3343 (m), 3291 (m) (ν_{NH}), 1880 cm^{–1} (s) (ν_{CO}); ¹H NMR (200 MHz, CD₂Cl₂, 293 K, TMS): δ = 7.40–6.86 (m, 20H; Ph), 4.35 (s, br, 2H; NH₂), 4.05 (m, 24H; CH₂); 3.93 (m, br, 1H; NH), 2.49 (d, ³J_{H,H} = 6 Hz, 3H; CH₃N), 1.29 ppm (t, 36H; CH₃); ³¹P{¹H} (200 MHz, CD₂Cl₂, 293 K, H₃PO₄ 85 % ext.): δ = 117.9 ppm (s); elemental analysis (%) calcd for C₅₀H₈₆BN₂O₁₃P₄Re (1244.14): C 48.27, H 6.97, N 2.25; found: C 48.15, H 7.01, N 2.13.

2–BPh₄, **3**–BPh₄: A sample of **1** (124 mg, 0.1 mmol) was placed in a three-necked 25-mL flask fitted with a solid-addition sidearm containing [Pb(OAc)₄] (0.1 mmol, 44 mg). The system was evacuated, CH₂Cl₂ (8 mL) was added, the solution cooled to –40 °C, and [Pb(OAc)₄] was added portionwise over 10–20 min to the cold stirring solution. The reaction mixture was then allowed to warm to 0 °C, stirred for 10 min, and the solvent removed under reduced pressure. The oil obtained was treated at 0 °C with ethanol (2 mL) containing NaBPh₄ (0.2 mmol, 68 mg). A white solid slowly separated out which was filtered and crystallized fractionally. A typical separation involved slow cooling from +20 to –25 °C of a saturated solution of the complexes prepared by adding ethanol (8 mL) to the white solid and enough CH₂Cl₂ to obtain a saturated solution at room temperature. The first crystals are of **2**–BPh₄, the second a mixture of **2**–BPh₄ and **3**–BPh₄ which was recrystallized. A total of 52 mg of **2**–BPh₄ (yield 42 %) was separated. By further cooling of the solution, 29 mg of white crystals of **3**–BPh₄ (yield 24 %) were obtained. Pure samples of **2**–BPh₄ and **3**–BPh₄ can also be obtained by Pasteur separation of crystals obtained by cooling a saturated solution of the reaction product in ethanol to –25 °C.

2–BPh₄: IR (KBr): $\tilde{\nu}$ = 1890 cm^{–1} (s) (ν_{CO}); ¹H NMR (200 MHz, CD₂Cl₂, 293 K, TMS): δ = 15.99 (s, br, 1H; NH), 7.40–6.70 (m, 20H; Ph), 4.37 (d, 3H; =NCH₃), 4.06 (m, 24H; CH₂), 1.33 ppm (t, 36H; CH₃); ³¹P{¹H} (200 MHz, CD₂Cl₂, 293 K, H₃PO₄ ext.): δ = 116.7 ppm (s); elemental analysis (%) calcd for C₅₀H₈₄BN₂O₁₃P₄Re (1242.12): C 48.35, H 6.82, N 2.26; found: C 48.19, H 6.95, N 2.30;

3–BPh₄: IR (KBr): $\tilde{\nu}$ = 1894 cm^{–1} (s) (ν_{CO}); ¹H NMR (200 MHz, CD₂Cl₂, 293 K, TMS): δ = 13.98 (s, br, 1H; NH), 7.60–6.80 (m, 20H; Ph), 4.06 (m, 24H; CH₂), 3.66 (m, br, 1H; N=CH₂), 1.34 ppm (t, 36H; CH₃); ³¹P{¹H} (200 MHz, CD₂Cl₂, 293 K, H₃PO₄ ext.): δ = 123.6 ppm (s); elemental analysis (%) calcd for C₅₀H₈₃BN₂O₁₃P₄Re (1227.11): C 48.94, H 6.82, N 1.14; found: C 49.08, H 6.96, N 1.10.

Received: March 13, 2002 [Z18883]

- [1] a) D. Sellmann, A. Brandl, R. Endell, *Angew. Chem.* **1973**, 85, 1122; *Angew. Chem. Int. Ed. Engl.* **1973**, 12, 1019; b) D. Sellmann, A. Brandl, R. Endell, *J. Organomet. Chem.* **1973**, 49, C22; c) G. Huttner, W. Gartzke, K. Allinger, *Angew. Chem.* **1974**, 86, 860; *Angew. Chem. Int. Ed. Engl.* **1974**, 13, 822; d) D. Sellmann, K. Jödden, *Angew. Chem.* **1977**, 89, 480; *Angew. Chem. Int. Ed. Engl.* **1977**, 16, 464; e) D.

- Sellmann, E. Böhlen, M. Waeber, G. Huttner, L. Zsolnai, *Angew. Chem.* **1985**, 97, 984; *Angew. Chem. Int. Ed. Engl.* **1985**, 24, 981; f) D. Sellmann, W. Soglowek, F. Knoch, M. Moll, *Angew. Chem.* **1989**, 101, 1244; *Angew. Chem. Int. Ed. Engl.* **1989**, 28, 1271; g) J. P. Collman, J. E. Hutchison, M. A. Lopez, R. Guillard, R. A. Reed, *J. Am. Chem. Soc.* **1991**, 113, 2794; h) D. Sellmann, J. Käppler, M. Moll, F. Knoch, *Inorg. Chem.* **1993**, 32, 960; i) D. Sellmann, K. Engl, F. W. Heinemann, J. Sieler, *Eur. J. Inorg. Chem.* **2000**, 1079.
- [2] a) M. R. Smith III, R. L. Keys, G. L. Hillhouse, A. L. Rheingold *J. Am. Chem. Soc.* **1989**, 111, 8312; b) M. R. Smith III, T.-Y. Cheng, G. L. Hillhouse, *J. Am. Chem. Soc.* **1993**, 115, 8638; c) T.-Y. Cheng, A. Ponce, A. L. Rheingold, G. L. Hillhouse, *Angew. Chem.* **1994**, 106, 703; *Angew. Chem. Int. Ed. Engl.* **1994**, 33, 657; d) T.-Y. Cheng, J. C. Peters, G. L. Hillhouse, *J. Am. Chem. Soc.* **1994**, 116, 204; e) D. Sutton, *Chem. Rev.* **1993**, 93, 995.
- [3] a) G. Albertin, S. Antoniutti, A. Bacchi, E. Bordignon, F. Busatto, G. Pelizzi, *Inorg. Chem.* **1997**, 36, 1296; b) G. Albertin, S. Antoniutti, E. Bordignon, S. Pattaro, *J. Chem. Soc. Dalton Trans.* **1997**, 4445; c) G. Albertin, S. Antoniutti, A. Bacchi, M. Bergamo, E. Bordignon, G. Pelizzi, *Inorg. Chem.* **1998**, 37, 479.
- [4] a) K. H. Geib, P. Harteck, *Ber. Dtsch. Chem. Ges. B* **1933**, 66, 1815; b) K. H. Geib, P. Harteck, *Trans. Faraday Soc.* **1934**, 30, 131.
- [5] a) J. E. Dickens, W. M. Irvine, C. H. DeVries, M. Ohishi, *Astrophys. J.* **1977**, 497, 307; b) G. Winnewisser, C. Kramer, *Space Sci. Rev.* **1999**, 90, 181.
- [6] F. Hoyle, N. C. Wickramasinghe, *Nature* **1976**, 264, 45.
- [7] P. A. Shapley, J. M. Shusta, J. C. Hunt, *Organometallics* **1996**, 15, 1622.
- [8] G. Albertin, S. Antoniutti, S. Garcia-Fontán, R. Carballo, F. Padoan, *J. Chem. Soc. Dalton Trans.* **1998**, 2071.
- [9] X-ray structural analysis: Philips PW1100 diffractometer equipped with a scintillation counter, graphite-monochromated Mo_{Kα} radiation ($\lambda = 0.71069 \text{ \AA}$). Data correction for absorption effects by the ψ scan method^[10] for both compounds, and intensity decay correction (40%) for **2-BPh₄**. Structural determination: direct methods^[11] and full-matrix least-squares refinement on all F^2 .^[12] Anisotropic displacement parameters refined in both cases for all non-hydrogen atoms; hydrogen atoms were introduced in idealized positions. Phosphite and phenyl groups were restrained to agree with typical bonding geometry from the literature. Crystal data for **2-BPh₄**: C₅₀H₈₄BN₂O₁₃P₄Re, $M_w = 1242.12$, crystal dimensions $0.3 \times 0.2 \times 0.2 \text{ mm}^3$, space group $P2_1/c$, monoclinic, $a = 13.002(2)$, $b = 24.570(5)$, $c = 20.054(4) \text{ \AA}$, $\beta = 95.49(2)^\circ$, $V = 6377(2) \text{ \AA}^3$, $Z = 4$, $\rho_{\text{calcd}} = 1.308 \text{ g cm}^{-3}$, $\theta_{\text{max}} = 30^\circ$, 18990 measured reflections (18537 unique), 4388 unique observed ($I > 2\sigma(I)$), $R_1 = 0.095$, $wR_2 = 0.26$ (on observed data), 176 restraints, 601 parameters, $GOF = 0.845$. Crystal data for **3-BPh₄**: C₅₀H₈₃BN₂O₁₃P₄Re, $M_w = 1227.11$, crystal dimensions $0.4 \times 0.3 \times 0.2 \text{ mm}^3$, space group $P\bar{1}$, triclinic, $a = 15.393(5)$, $b = 16.977(5)$, $c = 12.916(5) \text{ \AA}$, $\alpha = 100.02(5)$, $\beta = 91.63(5)$, $\gamma = 71.08(5)^\circ$, $V = 3143(2) \text{ \AA}^3$, $Z = 2$, $\rho_{\text{calcd}} = 1.290 \text{ g cm}^{-3}$, $\theta_{\text{max}} = 28^\circ$, 15138 measured unique reflections, 8634 unique observed ($I > 2\sigma(I)$), $R_1 = 0.048$, $wR_2 = 0.115$ (on observed data), 611 parameters, 79 restraints, $GOF = 0.912$. CCDC-181120 (**2-BPh₄**) and CCDC-181121 (**3-BPh₄**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/conts/retrieving.html (or from the Cambridge Crystallographic Data Centre, 12, Union Road, Cambridge CB2 1EZ, UK; fax: (+44) 1223-336-033; or deposit@ccdc.cam.ac.uk).
- [10] A. C. T. North, D. C. Phillips, F. S. Mathews, *Acta. Crystallogr. Sect. A* **1968**, 24, 351.
- [11] SIR97: A. Altomare, G. Cascarano, C. Giacovazzo, A. Guagliardi, M. C. Burla, G. Polidori, M. Camalli, *J. Appl. Crystallogr.* **1994**, 27, 435.
- [12] G. M. Sheldrick, SHELXL-97, Program for structure refinement, University of Göttingen, Göttingen (Germany), **1997**.
- [13] W. A. Herrmann, L. K. Bell, M. L. Ziegler, H. Pfisterer, C. Pahl, *J. Organomet. Chem.* **1983**, 247, 39.
- [14] J. Vicente, M. T. Chicote, M. D. Abrisqueta, R. Guerrero, P. G. Jones, *Angew. Chem.* **1997**, 109, 1252; *Angew. Chem. Int. Ed. Engl.* **1997**, 36, 1203.
- [15] D. A. Knight, M. A. Dewey, G. A. Stark, B. K. Bennett, A. M. Arif, J. A. Gladysz, *Organometallics* **1993**, 12, 4523.
- [16] Coordination geometry for **2**: Re–N 2.12(1), Re–CO 2.14(1), Re–P 2.354(4), NH–N 1.251(4), N–C 1.36(2), C–O 1.12(1) Å; Re–N–N 145(2), N–N–C 123(2)°.
- [17] Preliminary investigations show the presence of traces of ammonia in the final reaction mixture, but no other nitrogen-containing compound was unambiguously identified, and therefore no reaction path may be reasonably proposed.
- [18] Metal-mediated N–N or N=N bond activation is a topic of current interest. For some recent examples see: A. K. Verma, S. C. Lee, *J. Am. Chem. Soc.* **1999**, 121, 10838; R. G. Peters, B. P. Warner, C. J. Burns, *J. Am. Chem. Soc.* **1999**, 121, 5585; M. A. Aubart, R. G. Bergman, *Organometallics* **1999**, 18, 811; F. Maseras, M. A. Lockwood, O. Eisenstein, I. P. Rothwell, *J. Am. Chem. Soc.* **1998**, 120, 6598.
- [19] Interest in cleavage of the N–N bond of hydrazine stems from its importance to inorganic and bioinorganic reducing system(s): a) A. E. Shilov, *Metal Complexes in Biomimetic Chemical Reactions*, CRC, Boca Raton, FL, **1997**; b) R. R. Eady, *Chem. Rev.* **1996**, 96, 3013; c) G. J. Leigh, *Science* **1995**, 268, 827; d) R. R. Schrock, T. E. Glassman, M. G. Vale, *J. Am. Chem. Soc.* **1991**, 113, 725; e) S. M. Malinak, K. D. Demadis, D. Coucouvanis, *J. Am. Chem. Soc.* **1995**, 117, 3126; f) D. Sellmann, J. Sutter, *Acc. Chem. Res.* **1997**, 30, 460.

Total Synthesis of the Amaryllidaceae Alkaloid (+)-Plicamine and Its Unnatural Enantiomer by Using Solid-Supported Reagents and Scavengers in a Multistep Sequence of Reactions**

Ian R. Baxendale, Steven V. Ley,* and Claudia Piutti

Amaryllidaceae alkaloids are an important class of natural products especially as many members of the series display a wide range of potent biological activity. These properties include anticholinergic, antitumor, immunosuppressive, and analgesic activity, and they have also been shown to inhibit various cell cycle mechanisms (including HIV-1 activity), and have found recent application in the therapeutic treatment of Alzheimer's disease.^[1] Thus extensive synthetic studies of this family have been carried out over a number of years.^[2, 3] Furthermore, the search for new members of the series has proved to be extremely profitable.^[3, 4] The recently isolated compound (+)-plicamine (**1**) is especially attractive as it exemplifies many of the structural features of these natural

[*] Prof. S. V. Ley, I. R. Baxendale
Department of Chemistry
University of Cambridge
Lensfield Road, Cambridge CB2 1EW (UK)
Fax: (+44) 1223-336-442
E-mail: svl1000@cam.ac.uk
C. Piutti
Department of Chemistry
Pharmacia S.p.A
Discovery Research Oncology
Viale Pasteur, 10, 20014 Nerviano (MI) (Italy)

** We gratefully acknowledge financial support from Pfizer Central Research for a Postdoctoral Fellowship (to I.R.B.), the BP endowment and the Novartis Research Fellowship (to S.V.L.), and Pharmacia & Upjohn (to C.P.).