

- [16] A. L. Henne, J. W. Shepard, E. J. Young, *J. Am. Chem. Soc.* **1950**, *72*, 3577–3579.
- [17] R. N. Haszeldine, *J. Chem. Soc.* **1953**, 3565–3573.
- [18] R. N. Haszeldine, *J. Chem. Soc.* **1954**, 1273–1279.
- [19] A. T. Morse, P. B. Ayscough, L. C. Leitch, *Can. J. Chem.* **1955**, *33*, 453–457.
- [20] K. Eller, H. Schwarz, *Int. J. Mass Spectrom. Ion Processes* **1989**, *93*, 243–257.
- [21] K. Eller, W. Zummack, H. Schwarz, *J. Am. Chem. Soc.* **1990**, *112*, 621–627.
- [22] D. Schröder, H. Schwarz, D. E. Clemmer, Y. Chen, P. B. Armentrout, V. I. Baranov, D. K. Böhme, *Int. J. Mass Spectrom. Ion Processes* **1997**, *161*, 175–191.
- [23] R. B. Cody, S. C. Burnier, B. S. Freiser, *Anal. Chem.* **1982**, *54*, 96–101.
- [24] M. B. Comisarow, V. Grassi, G. Parisod, *Chem. Phys. Lett.* **1978**, *57*, 413–416.
- [25] U. Mazurek, H. Schwarz, *ICR Kinetics*, v. 3.0.1, TU Berlin, Berlin, **1998**, available upon request.
- [26] T. Su, W. J. Chesnavich, *J. Chem. Phys.* **1982**, *76*, 5183–5185.
- [27] T. Su, *J. Chem. Phys.* **1988**, *89*, 5355.
- [28] Parameters: water: dipole moment 1.854 D,^[30] polarizability $1.45 \times 10^{-24} \text{ cm}^3$,^[30] relative ion gauge sensitivity 0.97;^[31] hexafluoroacetone: dipole moment 0.395 D,^[32] polarizability $7.5 \times 10^{-24} \text{ cm}^3$ (estimation based on the data for acetone and fluorinated methanes $\text{CH}_n\text{F}_{4-n}$),^[30] relative ion gauge sensitivity 2.59.^[31]
- [29] J. N. Harvey, D. Schröder, W. Koch, D. Danovich, S. Shaik, H. Schwarz, *Chem. Phys. Lett.* **1997**, *273*, 164–170.
- [30] *CRC Handbook of Chemistry and Physics*, 79th ed. (Ed.: D. R. Lide), CRC Press, Boca Raton, FL, **1998**.
- [31] J. E. Bartmess, R. M. Georgiadis, *Vacuum* **1983**, *33*, 149–153.
- [32] J. U. Grabow, N. Heineking, W. Stahl, *Z. Naturforsch. A* **1991**, *46*, 229–232.
- [33] **Note added in proof** (June 3, 2002): Since the submission of this paper several publications have appeared on C–F bond activation; see for example: K. Uneyama, H. Amii, *J. Fluorine Chem.* **2002**, *114*, 127–131; K. Guennou de Cadenet, R. Rumin, F. Y. Pétilon, D. S. Yufit, K. W. Muir, *Eur. J. Inorg. Chem.* **2002**, 639–657; D. Zhang, C. Liu, S. Bi, *J. Phys. Chem. A* **2002**, *106*, 4153–4157.

Anti-Markovnikov Hydroamination of Terminal Alkynes**

Annegret Tillack, Ivette Garcia Castro,
Christian G. Hartung, and Matthias Beller*

*Dedicated to Professor Dr. Lutz F. Tietze
on the occasion of his 60th birthday*

Imines are of significant importance as intermediates for the synthesis of various amines and carbonyl compounds. In general, the synthesis of imines includes amination of a suitable aldehyde or ketone. A more atom-efficient route is the direct hydroamination of alkynes.^[1] This method has the additional advantage that no water is produced as a by-

product. Hence, various domino reactions (e.g. direct nucleophilic addition of organometallic reagents) are possible, which do not work in the presence of water.

The homogeneously catalyzed intermolecular hydroamination of alkynes is known to proceed in the presence of Hg and Tl salts,^[2] alkali metals (Cs),^[3] Ti,^[4] Zr,^[5] Nd,^[6] U, and Th^[7] complexes. In addition, complexes of late-transition metals (such as Ru, Pd,^[8] and Rh^[9]) have been used as catalysts for this transformation. Clearly, catalysts based on cheap and easily available titanium and zirconium complexes offer significant advantages compared to those based on toxic metals (Hg, Tl) or more expensive (U, Th, Ru, Pd, and Rh) complexes.

Recently, important progress in the intermolecular hydroamination of alkynes with titanium complexes was made by Johnson and Bergman^[10] and by Doye and co-workers.^[4] While the former group developed the modified titanium complex $[\text{Cp}(\text{ArNH})(\text{py})\text{Ti}=\text{NAr}]$ (Cp = cyclopentadienyl, py = pyridyl) and used it for the reaction of 2,6-dimethylaniline and diphenylacetylene, the latter group described an efficient and general method for the hydroamination of various internal alkynes using dimethyltitanocene as a catalyst. Bytschkov and Doye showed that the turnover frequency of this catalyst can be enhanced by using microwaves.^[4c] Kinetic measurements by Bergman^[10] and Doye^[11] also established a general mechanism of the dimethyltitanocene-catalyzed intermolecular hydroamination of alkynes. Surprisingly little attention was paid to the hydroamination of terminal alkynes using titanium catalysts,^[12] although the regioselective, sequential amination and hydroxylation of compounds that are unsaturated at the terminal position is one of the most challenging goals for industrial catalysis. Herein we report the first example of a titanocene-catalyzed anti-Markovnikov hydroamination of terminal aliphatic alkynes.

Some time ago we started a program on catalytic amination reactions of olefins and alkynes.^[13, 9] Inspired by the work of Doye and Bergman, we also recently looked for easily available and stable titanocene complexes. Here, titanocene alkyne complexes of the type $[\text{Cp}_2\text{Ti}(\eta^2\text{-Me}_3\text{SiC}\equiv\text{CR})]$ (Rosenthal's catalyst)^[14] appeared to be suited as amination catalysts.^[15] These complexes ($[\text{Cp}_2\text{Ti}(\eta^2\text{-Me}_3\text{SiC}\equiv\text{CSiMe}_3)]$ **1**^[14a] and $[\text{Cp}_2\text{Ti}(\eta^2\text{-Me}_3\text{SiC}\equiv\text{CPh})]$ **2**^[14b]) are easily synthesized by reaction of titanocene dichloride with the corresponding silylated alkyne.

Compared to previously used titanocene precatalysts, the titanacyclopentene complexes **1** and **2** are safe and stable under argon at room temperature for many months in German version. Indeed, hydroamination of internal alkynes (diphenylacetylene, 1-phenylpropyne) with aniline or *tert*-butylamine proceeds in excellent yields in the presence of **1** (81–98% yield after hydrolysis with HCl, Table 1). As shown in Table 2, different terminal aliphatic alkynes react with *tert*-butylamine with extremely high regioselectivity (>98%), in high yields (84–98%), and within a short time (2–24 h), by using 0.5–2.5 mol % of catalyst **1**, to give the imines **4a–e** and **5**. Although the reactions proceed smoothly with 0.5 mol % of catalyst, we used 2.5 mol % in general because of the shorter reaction time.

Pleasingly, only the anti-Markovnikov products were obtained, which is explained by the selective formation of the

[*] Prof. Dr. M. Beller, Dr. A. Tillack, Dr. I. Garcia Castro,
Dr. C. G. Hartung
Institut für Organische Katalyseforschung (IfOK)
Universität Rostock e.V.
Buchbinderstrasse 5–6, 18055 Rostock (Germany)
Fax: (+49) 381-466-9324
E-mail: matthias.beller@ifok.uni-rostock.de

[**] We acknowledge financial support of this project from the Deutsche Forschungsgemeinschaft (DFG) and the state of Mecklenburg-West Pomerania. C. Mewes, Dr. C. Fischer, H. Baudisch, and S. Buchholz are thanked for their excellent technical support.

Amines were distilled from CaH_2 . Alkynes were degassed, flushed with argon, and stored over molecular sieves (4 Å). All experiments were carried out under an argon atmosphere.

Catalysts **1** and **2** were synthesized according to a literature procedure.^[14a,b] Imines **4b–h** and **5** were isolated after distillation of the crude hydroamination mixtures and were characterized by NMR spectroscopy, MS, IR spectroscopy, and elemental analyses. Identification of all other products was performed by comparison with authentic products. Compounds **4a** and **4i** were synthesized according to the procedures in references [16] and [9], respectively.

Example of a typical hydroamination experiment (**4b**, Table 2, entry 2): A solution of 1-octyne (3.2 mL, 2.4 g, 21.5 mmol) and *tert*-butylamine (3.5 mL, 2.4 g, 32.2 mmol) was treated with **1** (0.15 g, 0.43 mmol, 2 mol % in toluene (6 mL). This mixture was heated to 85 °C and distilled under vacuum after 24 h. Product **4b** was obtained at 48–49 °C (0.1 mbar); yield: 2.9 g (75 %, >98 % (GC) anti-Markovnikov product); ^1H NMR (400 MHz, CDCl_3): δ = 0.85 (t, 3H, CH_3), 1.14 (s, 9H, C- CH_3), 1.21–1.34 (m, 8H, CH_2), 1.46 (m, 2H, CH_2), 2.20 (m, 2H, CH_2), 7.56 ppm (t, 3J = 5.35 Hz, 1H, CH); ^{13}C NMR (100 MHz, CDCl_3): δ = 14.0 (CH_3), 22.6, 26.4, 29.1, 29.2 (CH_2), 29.6 (C- CH_3), 31.7, 36.4 (CH_2), 56.4 (C_q), 159.3 ppm (CH); IR (neat): $\nu_{\text{C}=\text{N}}$: 1671 cm^{-1} ; MS (EI, 70 eV): m/z = 184 [M^+ +1], 183 [M^+], 168 [M^+ - CH_3], 99 [$\text{C}_7\text{H}_{15}^+$], 84 [$\text{C}_6\text{H}_9\text{NCH}^+$], 57 [C_4H_9^+]; elemental analysis $\text{C}_{12}\text{H}_{25}\text{N}$ calcd (%) C 78.62, H 13.74, N 7.64; found C 78.19, H 13.99, N 7.42.

Received: December 21, 2001 [Z18425]

- [1] Reviews: a) M. Nobis, B. Driessen-Hölscher, *Angew. Chem.* **2001**, *113*, 4105–4108; *Angew. Chem. Int. Ed.* **2001**, *40*, 3983–3986; b) T. E. Müller, M. Beller, *Chem. Rev.* **1998**, *98*, 675–703; c) J.-J. Brunet, D. Neibecker in *Catalytic Heterofunctionalization* (Eds.: A. Togni, H. Grützmaier), Wiley-VCH, Weinheim, **2001**, pp. 91–141; d) T. E. Müller, M. Beller in *Transition Metals for Organic Synthesis*, Vol. 2 (Eds.: M. Beller, C. Bolm), Wiley-VCH, Weinheim, **1998**, pp. 316–330; e) E. Haak, S. Doye, *Chem. Unserer Zeit* **1999**, *33*, 297–303; f) J. J. Brunet, D. Neibecker, F. Niedercorn, *J. Mol. Catal.* **1989**, *49*, 235–259; g) R. Taube in *Applied Homogeneous Catalysis with Organometallic Compounds*, Vol. 1 (Eds.: B. Cornils, W. A. Herrmann), Wiley-VCH, Weinheim, **1996**, S. 507–520.
- [2] a) J. Barluenga, F. Aznar, *Synthesis* **1975**, 704–705; b) J. Barluenga, F. Aznar, *Synthesis* **1977**, 195–196; c) J. Barluenga, F. Aznar, R. Liz, R. Rodes, *J. Chem. Soc. Perkin Trans. 1* **1980**, 2732–2737.
- [3] D. Tzalis, C. Koradin, P. Knochel, *Tetrahedron Lett.* **1999**, *40*, 6193–6195.
- [4] a) E. Haak, I. Bytschkov, S. Doye, *Angew. Chem.* **1999**, *111*, 3584–3586; *Angew. Chem. Int. Ed.* **1999**, *38*, 3389–3391; b) E. Haak, S. Doye, *Org. Lett.* **2000**, *2*, 1935–1937; c) I. Bytschkov, S. Doye, *Eur. J. Org. Chem.* **2001**, 4411–4418; d) E. Haak, I. Bytschkov, S. Doye, *Eur. J. Org. Chem.* **2002**, 457–463.
- [5] a) A. M. Baranger, P. J. Walsh, R. G. Bergman, *J. Am. Chem. Soc.* **1993**, *115*, 2753–2763; b) P. J. Walsh, A. M. Baranger, R. G. Bergman, *J. Am. Chem. Soc.* **1992**, *114*, 1708–1719.
- [6] Y. Li, T. J. Marks, *Organometallics* **1996**, *15*, 3770–3772.
- [7] a) A. Haskel, T. Straub, M. S. Eisen, *Organometallics* **1996**, *15*, 3773–3775; b) M. S. Eisen, T. Straub, A. Haskel, *J. Alloys Compd.* **1998**, *271*–273, 116–122; c) T. Straub, A. Haskel, T. G. Neyroud, M. Kapon, M. Botoshansky, M. S. Eisen, *Organometallics* **2001**, *20*, 5017–5035.
- [8] a) M. Tokunaga, M. Eckert, Y. Wakatsuki, *Angew. Chem.* **1999**, *111*, 3416–3419; *Angew. Chem. Int. Ed.* **1999**, *38*, 3222–3225; b) M. Tokunaga, Y. Wakatsuki, *J. Synth. Org. Chem. Jpn.* **2000**, *58*, 587–596; c) M. Heider, J. Henkelmann, T. Ruehl, *Eur. Pat. Appl.* EP 646571, **1995** [*Chem. Abstr.* **1995**, *123*, 22925s]; d) Y. Uchimaru, *Chem. Commun.* **1999**, 1133–1134; e) I. Kadota, A. Shibuya, L. Mpaka Lutete, Y. Yamamoto, *J. Org. Chem.* **1999**, *64*, 4570–4571.
- [9] C. G. Hartung, A. Tillack, H. Trauthwein, M. Beller, *J. Org. Chem.* **2001**, *66*, 6339–6343.
- [10] J. S. Johnson, R. G. Bergman, *J. Am. Chem. Soc.* **2001**, *123*, 2923–2924.
- [11] F. Pohlki, S. Doye, *Angew. Chem.* **2001**, *113*, 2361–2364; *Angew. Chem. Int. Ed.* **2001**, *40*, 2305–2308.

- [12] a) Y. Shi, J. T. Ciszewski, A. L. Odom, *Organometallics* **2001**, *20*, 3967–3969; b) C. Cao, J. T. Ciszewski, A. L. Odom, *Organometallics* **2001**, *20*, 5011–5013.
- [13] a) M. Beller, M. Eichberger, H. Trauthwein, *Angew. Chem.* **1997**, *109*, 2306–2308; *Angew. Chem. Int. Ed. Engl.* **1997**, *36*, 2225–2227; b) M. Beller, H. Trauthwein, M. Eichberger, C. Breindl, T. E. Müller, A. Zapf, *J. Organomet. Chem.* **1998**, *566*, 277–285; c) M. Beller, H. Trauthwein, M. Eichberger, C. Breindl, T. E. Müller, *Eur. J. Inorg. Chem.* **1999**, 1121–1132; d) M. Beller, O. R. Thiel, H. Trauthwein, C. G. Hartung, *Chem. Eur. J.* **2000**, *6*, 2513–2522; e) A. Tillack, H. Trauthwein, C. G. Hartung, M. Eichberger, S. Pitter, A. Jansen, M. Beller, *Monatsh. Chem.* **2000**, *131*, 1327–1334; f) M. Beller, H. Trauthwein, M. Eichberger, C. Breindl, J. Herwig, T. E. Müller, O. R. Thiel, *Chem. Eur. J.* **1999**, *5*, 1306–1319; g) H. Trauthwein, A. Tillack, M. Beller, *Chem. Commun.* **1999**, 2029–2030.
- [14] a) V. V. Burlakov, U. Rosenthal, P. V. Petrovskii, V. B. Shur, M. E. Vol'pin, *Organomet. Chem. USSR* **1988**, *1*, 526; b) V. V. Burlakov, U. Rosenthal, R. Beckhaus, A. V. Polyakov, Y. T. Struchkov, G. Oehme, V. B. Shur, M. E. Vol'pin, *Organomet. Chem. USSR* **1990**, *3*, 237–238; c) A. Tillack, C. Lefebvre, N. Peulecke, D. Thomas, U. Rosenthal, *Tetrahedron Lett.* **1997**, *38*, 1533–1534; d) N. Peulecke, D. Thomas, W. Baumann, C. Fischer, U. Rosenthal, *Tetrahedron Lett.* **1997**, *38*, 6655–6656; e) N. Peulecke, A. Ohff, P. Kosse, A. Tillack, A. Spannenberg, R. Kempe, W. Baumann, V. V. Burlakov, U. Rosenthal, *Chem. Eur. J.* **1998**, *4*, 1852–1861; f) S. Pulst, F. G. Kirchbauer, B. Heller, W. Baumann, U. Rosenthal, *Angew. Chem.* **1998**, *110*, 2029–2031; *Angew. Chem. Int. Ed.* **1998**, *37*, 1915–1927; g) P. Arndt, A. Spannenberg, W. Baumann, S. Becke, U. Rosenthal, *Eur. J. Inorg. Chem.* **2001**, 2885–2890; Rosenthal's catalysts are now commercially available from Fluka.
- [15] N. Peulecke, Ph.D. thesis, Rostock, **1997** describes an amination of trimethylsilylacetylene and phenylacetylene with aniline in the presence of **1** with a maximum yield of 50 %. However, other amines, such as *tert*-butylamine, do not react.
- [16] N. Y. Grigor'eva, I. M. Avrutov, O. A. Pinsker, O. N. Yudina, A. I. Lutsenko, A. M. Moiseenkov, *Izv. Akad. Nauk SSSR Ser. Khim.* **1985**, *8*, 1824–1835.

Formation of a Butadienenickel-Based Zwitterionic Single-Component Catalyst for Ethylene Polymerization: An Alternative Activation Pathway for Homogeneous Ziegler–Natta Catalysts of Late Transition Metals**

Joachim W. Strauch, Gerhard Erker,* Gerald Kehr, and Roland Fröhlich

Homogeneous Ziegler–Natta catalysis has become very important in recent years.^[1] In addition to the many reported catalysts based on metallocenes with metals from Group 4 and related systems, a variety of Ziegler–Natta systems of late

[*] Prof. Dr. G. Erker, Dipl.-Chem. J. W. Strauch, Dr. G. Kehr, Dr. R. Fröhlich^[+]
Organisch-Chemisches Institut der Universität
Westfälische Wilhelms-Universität Münster
Corrensstrasse 40, 48149 Münster (Germany)
Fax: (+49)251-833-36503
E-mail: erker@uni-muenster.de

[+] X-ray crystal structure analysis.

[**] This work was supported by the Fonds der Chemischen Industrie (and the BMBF) and the Deutsche Forschungsgemeinschaft.