


- 1996, 155, 127–144; c) N. Komatsu in *Organobismuth Chemistry* (Eds.: H. Suzuki, Y. Matano), Elsevier, New York, 2001, chap. 5, pp. 371–440.
- [4] a) S. Maeda in *The Chemistry of Organic Arsenic, Antimony and Bismuth Compounds* (Ed.: S. Patai), Wiley, New York, 1994, chap. 19, pp. 725–759; b) J. Reglinski, in *Chemistry of Arsenic, Antimony and Bismuth* (Ed.: N. C. Norman), Blackie Academic and Professional, London, 1998, . 8, pp. 403–440.
- [5] F. Challenger, O. V. Richards, *J. Chem. Soc.* **1934**, 405–411.
- [6] a) D. H. R. Barton, J. P. Kitchin, W. B. Motherwell, *J. Chem. Soc. Chem. Commun.* **1978**, 1099–1100; b) D. H. R. Barton, D. J. Lester, W. B. Motherwell, M. T. B. Papoula, *J. Chem. Soc. Chem. Commun.* **1979**, 705–707; c) D. H. R. Barton, J. P. Kitchin, D. J. Lester, W. B. Motherwell, M. T. B. Papoula, *Tetrahedron* **1981**, 37, 73–79 (Supplement 9).
- [7] a) V. A. Dodonov, A. V. Gushchin, T. G. Brilkina, *Zh. Obshch. Khim.* **1985**, 55, 73–80; b) V. A. Dodonov, T. G. Brilkina, A. V. Gushchin, *Zh. Obshch. Khim.* **1981**, 51, 2380.
- [8] For instance, the reaction conditions for the oxidations of cinnamyl alcohol and 1-pentanol with a $(\text{Ph}_3\text{BiCl})_2\text{O}$ /base system were reported to be 15 h at 21 °C and 6 h at 60 °C, respectively.^[6a]
- [9] Y. Matano, H. Nomura, *J. Am. Chem. Soc.* **2001**, 123, 6443–6444. Compound **1** was prepared by the reaction of tris(2-methoxyphenyl)-bismuth dichloride with 2 equiv of *t*BuOK in the presence of 5 equiv of water.
- [10] a) V. A. Dodonov, T. I. Zinov'eva, N. N. Osadchaya, *Zh. Obshch. Khim.* **1988**, 58, 712; b) H. Suzuki, T. Ikegami, *J. Chem. Res. (S)* **1996**, 24–25; c) Y. Matano, H. Nomura, H. Suzuki, M. Shiro, H. Nakano, *J. Am. Chem. Soc.* **2001**, 123, 10954–10965.
- [11] H. Suzuki, T. Ikegami, Y. Matano, *Tetrahedron Lett.* **1994**, 35, 8197–8200.
- [12] The initially formed $\text{Ar}_2\text{BiOBiAr}_2$ disproportionated in solution to give Ar_3Bi .
- [13] CCDC-182209 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/conts/retrieving.html (or from the Cambridge Crystallographic Data Centre, 12, Union Road, Cambridge CB21EZ, UK; fax: (+44) 1223-336-033; or deposit@ccdc.cam.ac.uk).
- [14] In the *t*BuOK system, a long column was necessary to completely remove the Bi^{III} by-product (Ar_3Bi) from the carbonyl products in cases where their R_f values are similar.
- [15] A similar mechanism was proposed by Barton et al. for the Ph_3BiX_2 -based oxidation of alcohol.^[6c]
- [16] J. Matsuo, H. Kitagawa, D. Iida, T. Mukaiyama, *Chem. Lett.* **2001**, 150–151.
- [17] Trifluoromethyl carbinols are known to resist oxidation and, at present, only a few methods are available for their conversion into the corresponding trifluoromethyl ketones; see a) R. J. Linderman, D. M. Graves, *J. Org. Chem.* **1989**, 54, 661–668; b) I. E. Markó, P. R. Giles, M. Tsukazaki, I. Chellé-Regnaut, A. Gautier, S. M. Brown, C. J. Urch, *J. Org. Chem.* **1999**, 64, 2433–2439; c) V. Kesavan, D. Bonnet-Delpon, J.-P. Bégué, A. Srikanth, S. Chandrasekaran, *Tetrahedron Lett.* **2000**, 41, 3327–3330.
- [18] a) D. B. Dess, J. C. Martin, *J. Org. Chem.* **1983**, 48, 4155–4156; b) D. B. Dess, J. C. Martin, *J. Am. Chem. Soc.* **1991**, 113, 7277–7287.
- [19] A. Gillmeister, *Chem. Ber.* **1897**, 30, 2843–2850. In this work, **2a** was prepared by oxidative chlorination of tris(2-methylphenyl)bismuthane with sulfuryl chloride.

Synthesis of Cycloalkanones from Dienes and Allylamines through C–H and C–C Bond Activation Catalyzed by a Rhodium(I) Complex**

Dae-Yon Lee, In-Jung Kim, and Chul-Ho Jun*

Among numerous examples of transition-metal-mediated cyclization to obtain medium-sized ring compounds,^[1] intramolecular hydroacylation provides the most promising way to prepare cyclopentanones from 4-pentenal through the C–H-bond activation of an aldehyde.^[2,3] However, its application has been limited to the synthesis of five-membered rings with a few exceptions,^[3] because the competing decarbonylation of the acyl metal hydride intermediate prevails during the formation of the larger ring. In the course of our studies into chelation-assisted C–H- and C–C-bond activation,^[4] allylic amine **1a**, which bears a coordination site, was devised and used as a masked form of formaldehyde in the hydroacylation of 1-alkenes to synthesize dialkyl ketones.^[4d] We envisaged a cyclization of dienes **2**^[5–7] with **1a** to furnish cycloalkanones **3** with various sizes, since decarbonylation cannot occur in the reaction of **1a** (Scheme 1). Herein we present a facile synthesis of various cycloalkanones from the reaction of allylic amines with dienes through chelation-assisted C–H- and C–C-bond activation.



Scheme 1. Formation of cycloalkanone **3** from the reaction of allylamine **1a** and diene **2**.

The reactions of **1a** with various dienes (**2**) are summarized in Table 1. For example, when **1a** was allowed to react with 1,4-pentadiene (**2a**) in the presence of $[(\text{C}_8\text{H}_{14})_2\text{RhCl}]_2$ (**4**, 5 mol %) and PCy_3 (**5**, 15 mol %) at 150 °C for 2 h, cyclohexanone (**3a**) and 2-methylcyclopentanone (**3b**) were obtained in 87 % and 13 % yield, respectively, after hydrolysis (Table 1, entry 1). Furthermore, the reaction with 1,5-hexadiene (**2b**) provided cycloheptanone (**3c**), 2-methylcyclohexanone (**3d**), and 2-ethylcyclopentanone (**3e**) in a ratio of 38:40:22 (Table 1, entry 2).^[8]

The mechanism for this reaction is depicted in Scheme 2. Aldimine **6a** is generated by Rh-catalyzed isomerization of **1a**. Subsequent intermolecular hydroiminoacylation^[9] of **2b**

[*] Prof. Dr. C.-H. Jun, D.-Y. Lee, I.-J. Kim
Department of Chemistry, Yonsei University
Seoul 120-749 (Korea)
Fax: (+82)2-364-7050
E-mail: junch@yonsei.ac.kr

[**] This work was supported by the National Research Laboratory Program (2000-N-NL-01-C-271) administered by Ministry of Science and Technology, and by the Korean Science and Engineering Foundation (20004010).



Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.


Table 1. The cyclization of dienes **2** with **1a**^[a]

Entry	Dienes 2	<i>t</i> [h]	Cycloalkanones 3 ^[b]	Yield
1 ^[c]	2a	2	3a 3b	(100%, 87:13)
2 ^[c,d]	2b	2	3c 3d 3e	(82%, 38:40:22)
3 ^[e]	2c	0.5	3f	(56%)
4 ^[f]	2d	2	3g	86% (100%)
5	2e	4	3h	84% (100%)
6	2f	5	3i	77% (100%)
7 ^[c]	2g	3	3j	67% (100%)
8 ^[g]	2h	18	3b	(77%)

[a] **1a** (0.25 mmol), **2** (0.75 mmol), **4** (0.0063 mmol), **5** (0.039 mmol) in toluene (100 mg) at 170°C. The products were identified after hydrolysis. [b] Yields of isolated cyclohexanones (GC yields are given in parentheses). [c] The reaction temperature was 150°C. [d] A mixture of alkenyl ketones **8a** (9%) and **8b** (9%), which were identified as 1-phenyl-3-nonanone and dihexyl ketone, respectively, were also observed (GC) after hydrogenation. [e] A mixture of alkenyl ketones **8c** (23%) and **8d** (21%) was also obtained. [f] The ratio of meso-**2d** (±-**2d** (ca. 45:55) was retained in **3g** (determined by ¹H NMR spectroscopy). [g] A mixture of alkenyl ketones, identified as a hydrogenated form, 1-phenyl-3-octanone, was obtained (33% yield by GC).

with **6a** gives ketimine **7**, which is hydrolyzed to give acyclic ketone **8a**.^[4d] Further C–C-bond activation of **7**, and subsequent β-hydrogen elimination leads to iminoacylrhodium(III) hydride **9a**, and styrene is liberated.^[4c,d] The intramolecular hydrometalation of **9a** affords ketimines **11a–c**, which furnish corresponding cycloalkanones **3c–e** after hydrolysis. Among the products, ketimine **11a** is derived from anti-Markovnikov hydrometalation of **9a** via metallacyclic intermediate **10a**, whereas **11b** is formed from Markovnikov reaction of **9a** via **10b**. Further skeletal isomerization of **10b** leads to the formation of **11c** via **9b** and **10c**.

Monitoring the change in the ratio of cycloalkanones in the reaction of **1a** with **2b** revealed that the seven-membered ring



Scheme 2. Postulated mechanism for the cyclization of **2b** with **1a**.

3c was the sole product after 5 min (Figure 1). The ratio of **3c** decreased while those of **3d** and **3e** increased as the reaction progressed.^[10] This type of skeletal rearrangement by C–C-





Figure 1. Plot of the ratio of cycloalkanones vs time plot for the reaction of **1a** and **2b** at 150°C. y = Ratio of cycloalkanones determined by GC: ● = **3c**, ▲ = **3d**, ■ = **3e**.

bond activation, which leads to the spontaneous ring contraction of the seven-membered ring to the more stable six- and five-membered rings, has been already studied.^[4f,11] However, the initial formation of a seven-membered ring implies that anti-Markovnikov hydrometalation (Scheme 2, **9a** to **10a**) is favored over Markovnikov hydrometalation (Scheme 2, **9a** to **10b**), maybe as a result of the steric congestion of the (iminoacyl)rhodium system.^[4a]

The exclusive formation of a seven-membered ring was observed with substituted dienes. For instance, the reaction of 2-methyl-1,5-hexadiene (**2c**) with **1a** gave 3-methylcycloheptanone (**3f**) as the sole cyclization product in 56% yield, along with alkenyl ketones **8c** and **8d**, which shows that the initial hydroiminoacylation takes place at the less substituted olefin, that is, C5=C6 in **2c** (Table 1, entry 3). The subsequent cyclization occurs at C1=C2 in **2c** to form **3f**, but further ring contraction did not proceed, because the formation of the resulting metallacycle intermediate **10d** is disfavored as a result of steric congaestion.




Other 1,5- or 1,4-dienes that bear substituents at C2 or C3 also exclusively yielded seven- or six-membered rings, respectively, without forming ring-contraction products (Table 1, entries 4–7). Notably, even the 1,3-diene piperylene (**2h**) also underwent cyclization to give **3b** in good yield (Table 1, entry 8). So far, the hydroacylation of conjugated dienes has been known to afford β,γ -unsaturated ketone via π -allyl intermediates.^[12]


Another interesting substrate is 3-benzyloxy-1,5-hexadiene (**2i**), which reacted with **1a** to give cycloalkanones **3k** and **3l** (67:33; Scheme 3). The formation of **3l** instead of **3m** as a ring-contraction product illustrates that initial hydroiminoacylation occurs exclusively at C1=C2 in **2i** to give intermediate **12a**, which is favored over **12b** as a result of the directing effect of the benzyloxy group, thus forming the stable five-membered metallacycle in **12a**.^[13]

Allylic amines that have no coordination site could also be applied to this reaction. For example, *N*-cinnamylamine (**1b**) reacted with **2i** in the presence of 2-amino-3-picoline (**13**) to give **3k** and **3l** in 72% yield after hydrolysis (Scheme 4). This reaction proceeds through the isomerization of **1b** and subsequent transimination^[4e,14] of the resulting aldimine **6b** to form **6a**.

In summary, the synthesis of various cycloalkanones by the reaction of allylic amines with dienes was achieved through chelation-assisted C–H- and C–C-bond activation. The use of



Scheme 3. Reaction of **2i** with **1a** in the presence of **4** and **5**.



Scheme 4. Reaction of **1b** and **2i** through transimination.

1a as a masked form of formaldehyde allows the formation of six- and seven-membered cycloalkanones, which has rarely been possible by intramolecular hydroacylation as a result of competing decarbonylation. Furthermore, the reaction with substituted 1,4- or 1,5-dienes exclusively gave cyclohexanones or cycloheptanones, respectively. Even allylic amines that have no coordination site can also be applied to this reaction by utilizing a transimination protocol.

Experimental Section

Reaction of 1b and 2i: A screw-capped pressure vial (1 mL) was charged with **1b** (39.2 mg, 0.190 mmol), **2i** (44.9 mg, 0.239 mmol), **13** (43.2 mg, 0.400 mmol), [(C₈H₁₄)₂RhCl]₂ (**4**; 7.3 mg, 0.012 mmol), tricyclohexylphosphane (**5**; 10.8 mg, 0.0599 mmol), and the reaction mixture was dissolved in toluene (100 mg). It was stirred in a preheated oil bath (170 °C) for 12 h. Upon completion, the reaction mixture was hydrolyzed (1 N HCl) and purified by column chromatography (SiO₂, *n*-hexane/ethyl acetate 5:2) to afford a mixture of **3k** and **3l** in 84% yield (24.8 mg). The ratio of **3k**/**3l** was determined by GC analysis as 53:47.

Received: March 30, 2002 [Z19011]

- [1] L. Yet, *Chem. Rev.* **2000**, *100*, 2963.
- [2] a) K. Sakai, J. Ide, O. Oda, N. Nakamura, *Tetrahedron Lett.* **1972**, 1287; b) R. C. Larock, K. Oertle, G. F. Potter, *J. Am. Chem. Soc.* **1980**, *102*, 190; c) D. Milstein, *J. Chem. Soc. Chem. Commun.* **1982**, 1357; d) D. P. Fairlie, B. Bosnich, *Organometallics* **1988**, *7*, 936; e) D. P. Fairlie, B. Bosnich, *Organometallics* **1988**, *7*, 946; f) B. Bosnich, *Acc. Chem. Res.* **1998**, *31*, 667; g) M. Tanaka, M. Imai, M. Fujio, E. Sakamoto, M. Takahashi, Y. Eto-Kato, X. M. Wu, K. Funakoshi, K. Sakai, H. Suemune, *J. Org. Chem.* **2000**, *65*, 5806, and references therein.
- [3] a) K. P. Gable, G. A. Benz, *Tetrahedron Lett.* **1991**, *32*, 3473; b) A. D. Aloise, M. E. Layton, M. D. Shair, *J. Am. Chem. Soc.* **2000**, *122*, 12610; c) Y. Sato, Y. Oonishi, M. Mori, *Angew. Chem.* **2002**, *114*, 1266; *Angew. Chem. Int. Ed.* **2002**, *41*, 1218.
- [4] a) C.-H. Jun, H. Lee, J.-B. Hong, *J. Org. Chem.* **1997**, *62*, 1200; b) C.-H. Jun, J.-B. Hong, D.-Y. Lee, *Synlett* **1999**, *1*; c) C.-H. Jun, H. Lee, *J. Am. Chem. Soc.* **1999**, *121*, 880; d) C.-H. Jun, H. Lee, J.-B. Hong, *Org. Lett.* **1999**, *1*, 2161; e) C.-H. Jun, D.-Y. Lee, H. Lee, J.-B. Hong, *Angew. Chem.* **2000**, *112*, 3214; *Angew. Chem. Int. Ed.* **2000**, *39*, 3070; f) C.-H. Jun, H. Lee, S.-G. Lim, *J. Am. Chem. Soc.* **2001**, *123*, 751; g) C.-H. Jun, H. Lee, C. W. Moon, H.-S. Hong, *J. Am. Chem. Soc.* **2001**, *123*, 8600.
- [5] For the carbonylative cyclization of a 1,4-diene under hydroformylation conditions, see: a) P. Eilbracht, M. Acker, I. Hädrich, *Chem. Ber.* **1988**, *121*, 519; b) P. Eilbracht, E. Balß, M. Acker, *Chem. Ber.* **1985**, *118*, 825.
- [6] For the carbonylative cycloaddition of 1,3-conjugated systems to prepare γ -lactams, see: T. Morimoto, N. Chatani, S. Murai, *J. Am. Chem. Soc.* **1999**, *121*, 1758.
- [7] For the transformation of dienes into cycloalkanones by means of hydroboration, see: a) U. P. Dhokte, P. M. Pathare, V. K. Magubdroo, H. C. Brown, *J. Org. Chem.* **1998**, *63*, 8276; b) H. C. Brown, E.-i. Negishi, *J. Am. Chem. Soc.* **1967**, *89*, 5477.
- [8] A trace amount (ca. 2%) of 2,5-dimethylcyclopentanone was also observed after 6 h, which might be derived from the skeletal rearrangement of **3b**.^[4f]
- [9] J. W. Suggs, *J. Am. Chem. Soc.* **1979**, *101*, 489.
- [10] The total yield of cycloalkanones did not increase much after 2 h, but the amounts of **3d** and **3e** kept increasing as that of **3c** decreased. See Supporting Information.
- [11] For the skeletal rearrangement of cycloalkanones through C–C-bond activation, see also: M. Murakami, K. Takahashi, H. Amii, Y. Ito, *J. Am. Chem. Soc.* **1997**, *119*, 9307.
- [12] a) T. Kondo, N. Hiraishi, Y. Morisaki, K. Wada, Y. Watanabe, T. Mitsudo, *Organometallics* **1998**, *17*, 2131; b) C.-H. Jun, B.-T. Koo, J.-B. Kang, K.-J. Kim, *Bull. Korean Chem. Soc.* **1994**, *15*, 1064.
- [13] C.-H. Jun, J.-S. Han, J.-B. Kang, S.-I. Kim, *J. Organomet. Chem.* **1994**, *474*, 183.
- [14] C.-H. Jun, J.-B. Hong, *Org. Lett.* **1999**, *1*, 887.