



A Pyridylsilyl Group Expands the Scope of Catalytic Intermolecular Pauson–Khand Reactions**

Kenichiro Itami,* Koichi Mitsudo, and Jun-ichi Yoshida*

The [2+2+1] cycloaddition of alkyne, alkene, and carbon monoxide mediated or catalyzed by a transition metal M, which is known as the Pauson–Khand reaction (PKR),^[1] has received particular attention in organic synthesis, since it allows the construction of biologically interesting five-membered carbocycles in a convergent manner. Although tremendous progress has been seen for the catalytic *intramolecular* PKR [Eq. (1)] and many reactions have been reported,^[1] the




development of an efficient and general procedure for a catalytic *intermolecular* PKR [Eq. (2)] lags far behind.^[1,2] In particular, the limitations imposed on the alkene counterpart are serious: 1) No catalytic intermolecular PKR has been reported yet for simple alkenes (the reactions are restricted to ethylene gas and strained alkenes),^[2] and 2) in a stoichiometric PKR, simple alkenes exhibit low reactivity and regioselectivity.^[1,3] This may be due to the low coordinating ability of alkenes in comparison with alkynes and carbon monoxide.

Although such problems might be overcome by incorporating a coordinating heteroatom tethered to the alkenes,^[4,5] the removal of such a directing group after the PKR is strongly desirable from a synthetic point of view. Therefore, the utilization of a removable directing group^[6] is expected to enhance the synthetic potential of the catalytic intermolecular PKR by eliminating the problems associated with reactivity and regioselectivity issues of the alkene.

Our recent demonstration that the dimethyl(2-pyridyl)silyl (2-PyMe₂Si) group serves as an excellent, removable directing group in a number of metal-catalyzed reactions^[7,8] led us to investigate the possibility of alkenyldimethyl(2-pyridyl)silane

as a substrate for a catalytic intermolecular PKR. The oxidative cyclization of alkyne, alkene, and metal can be regarded as a carbometalation of an (alkyne)metal complex across an alkene.^[9] Therefore, we expected the facile and regioselective formation of a metallacyclopentene intermediate owing to the coordination effect of the pyridyl group on silicon (Scheme 1). We have already established the high



Scheme 1. The catalytic intermolecular Pauson–Khand reaction directed by a pyridylsilyl group.

reactivity of alkenyldimethyl(2-pyridyl)silanes in several carbometalation processes.^[7b,f,10] Moreover, the resultant 2-PyMe₂Si-substituted cyclopentenone should be easily converted into the Si-free derivative by using various desilylation methods that we developed.^[11]

We first screened for a catalyst for the intermolecular PKR using dimethyl(2-pyridyl)(vinyl)silane (**1a**). Among various metal complexes (Ti, Fe, Co, Mo, Ru, Rh, Pd, Ir, and Pt), [Ru₃(CO)₁₂] was found to catalyze this process.^[12,13] The reaction of **1a** (1.0 equiv), phenylacetylene (**2a**, 1.5 equiv), and carbon monoxide (1 atm) at 100°C in toluene in the presence of [Ru₃(CO)₁₂] (5 mol%) gave the desilylated cyclopentenone **3aa**, which was isolated in 55% yield with virtually complete regioselectivity (Table 1, entry 1).^[14] The trend in regioselectivity (the Ph group is in the position α to the carbonyl group) is in line with that observed in the stoichiometric PKR.^[1] The realization of a catalytic intermolecular PKR without the need for a high pressure of carbon monoxide is particularly noteworthy. Under the identical conditions, no reaction occurred when dimethylphenyl(vinyl)silane was used in place of **1a**, which clearly implicates the strong directing effect of the pyridyl group. This is not only an example of the successful utilization of vinylsilane as an alkene counterpart in a catalytic intermolecular PKR,^[15] but also a demonstration that **1a** could serve as a nongaseous ethylene equivalent in this reaction.^[16]










When the alkyl-substituted alkyne **2b** was employed, a substantial decline in regioselectivity was observed (Table 1, entry 2). Symmetrical internal alkynes also participate in the catalytic intermolecular PKR with reasonable efficiency. When 1-phenylpropyne (**2e**) was used, cyclopentenone **3ae** was obtained with virtually complete regioselectivity (entry 5). However, when the chloro-substituted alkyne **2f** was allowed to react, there was a substantial decline in the regioselectivity (entry 6). Since **2e** and **2f** are sterically and

[*] Assist. Prof. K. Itami, Prof. J. Yoshida, K. Mitsudo
Department of Synthetic Chemistry and Biological Chemistry
Graduate School of Engineering, Kyoto University
Yoshida, Kyoto 606-8501 (Japan)
Fax: (+81)75-753-5911
E-mail: itami@sbchem.kyoto-u.ac.jp
yoshida@sbchem.kyoto-u.ac.jp

[**] This work was supported by a Grant-in-Aid for Scientific Research from the Japan Society for the Promotion of Science (JSPS). K.M. thanks JSPS for the Promotion of Science for Young Scientists.









Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.

Table 1. Catalytic intermolecular PKR of **1a** and various alkynes **2**.^[a]

			
Entry	2	Products	Yield [%] (3/4)
1	2a (R = H, R' = Ph)	 3aa	55 (100/0) ^[b,c]
2	2b (R = H, R' = C ₆ H ₁₃)	 3ab  4ab	91 (59/41) ^[b]
3	2c (R = R' = Ph)	 3ac	88 ^[d]
4	2d (R = R' = C ₃ H ₇)	 3ad	59 ^[d]
5	2e (R = CH ₃ , R' = Ph)	 3ae	75 (100/0) ^[b,c]
6	2f (R = (CH ₂) ₄ Cl, R' = Ph)	 3af  4af	67 (87/13) ^[d]

[a] Reaction conditions: **1a** (0.50 mmol), **2** (0.75 mmol), [Ru₃(CO)₁₂] (0.025 mmol), CO (1 atm), solvent (1.5 mL). [b] Reaction at 100 °C in toluene. [c] Compound **4** was not detected by NMR spectroscopy and GC analysis. [d] Reaction at 120 °C in xylenes.

Table 2. Catalytic intermolecular PKR of substituted vinylsilanes **1** and alkynes **2**.^[a]


			
1	2	Products	Yield [%] (3/4)
1b (R ¹ = C ₄ H ₉ , R ² = H)	2a (R ³ = H, R ⁴ = Ph)	 3ba	41 (100/0) ^[b,c]
1b (R ¹ = C ₄ H ₉ , R ² = H)	2b (R ³ = H, R ⁴ = C ₆ H ₁₃)	 3bb  4bb	55 (89/11) ^[d]
1b (R ¹ = C ₄ H ₉ , R ² = H)	2c (R ³ = R ⁴ = Ph)	 3bc	44 ^[b]
1c (R ¹ = (CH ₂) ₄ OMe, R ² = H)	2c (R ³ = R ⁴ = Ph)	 3cc	48 ^[b]
1d (R ¹ = H, R ² = CH ₃)	2b (R ³ = H, R ⁴ = C ₆ H ₁₃)	 3db  4db	40 (62/38) ^[e]

[a] Reaction conditions: **1** (0.50 mmol), **2** (0.75 mmol), [Ru₃(CO)₁₂] (0.025 mmol), CO (1 atm), solvent (1.5 mL). [b] Reaction at 140 °C in xylenes. [c] Compound **4** was not detected by NMR spectroscopy and GC analysis. [d] Reaction at 100 °C in toluene. [e] Reaction at 120 °C in xylenes.

electronically similar, the observed differences in regioselectivity may be due to the directing effect of chloride, which has been occasionally observed in Ru and other metal complexes.^[17]


We next turned our attention to the synthesis of more highly substituted cyclopentenones by employing substituted vinylsilanes (Table 2). The advantage of our system is that the use of α - or β -substituted vinylsilanes (**1b–d**)^[18] results in the regioselective production of substituted cyclopentenones with the substituent at the 5- or 4-position, respectively. In all cases, virtually complete regioselectivity with regard to the alkene subunit was observed. Thus, these substituted vinylsilanes not only serve as surrogates for terminal alkenes that cannot be applied in the previously reported catalytic PKR, but also enable the complete regioselective incorporation of the alkene subunit into the cyclopentenone skeleton. Although converting the 2-PyMe₂Si group into functional groups other than a hydrogen atom is not feasible at this stage, the present observations clearly demonstrate the power and usefulness of using the 2-PyMe₂Si group as a removable directing group in the catalytic intermolecular PKR.

Our catalytic intermolecular PKR should be a straightforward approach toward the synthesis of natural and nonnatural cyclopentanoid products. For example, cyclopentenone **6**, which has emerged as an excellent precursor for the synthesis of jasmones,^[19] methyl jasmonate,^[19] and methyl epijasmone,^[19] can be prepared in a single chemical operation from **1a** and enyne **5** [Eq. (3); 60 % yield, 63 % regioselectivity].^[20]



Moreover, the synthesis of rosaprostol,^[21] tetrahydrodicranone B,^[21] equilenins,^[22] or, more importantly, prostaglandins^[23] should be also possible starting from a suitably substituted 2-cyclopentenone that can, in principle, be prepared by our catalytic intermolecular PKR methodology.

In conclusion, we have developed an efficient procedure for the catalytic, intermolecular, and regioselective PKR of unstrained alkenes by utilizing a 2-PyMe₂Si group as a removable directing group. Multisubstituted 2-cyclopentenones can now be analyzed retrosynthetically, as in Equation (4). The advantages of our strategy are that 1) it allows a




formal catalytic intermolecular PKR using simple alkenes, which was previously difficult to achieve; 2) it allows a complete regioselective incorporation of a substituent at the 4- or 5-position of the 2-cyclopentenone skeleton simply by choosing the position of the substituent in the starting alkenylsilane; 3) the position of the C=C bond can be

strategically altered by simply changing the alkyne subunit to be hydrosilylated prior to the intermolecular PKR;^[18] and 4) all the substituents arise from substituents on alkyne subunits which can be easily altered. Thus, this new strategy offers a general approach to a diverse range of cyclopentenoid structures and should find further applications in organic synthesis.

Received: May 22, 2002 [Z19338]

- [1] Reviews: a) K. M. Brummond, J. L. Kent, *Tetrahedron* **2000**, 56, 3263; b) N. E. Schore, *Org. React.* **1991**, 40, 1.
- [2] Examples of catalytic intermolecular PKR: a) I. U. Khand, G. R. Knox, P. L. Pauson, W. E. Watts, M. I. Foreman, *J. Chem. Soc. Perkin Trans. 1* **1973**, 977; b) D. C. Billington, *Tetrahedron Lett.* **1983**, 24, 2905; c) V. Rautenstrauch, P. Mégard, J. Conesa, W. Küster, *Angew. Chem.* **1990**, 102, 1441; *Angew. Chem. Int. Ed. Engl.* **1990**, 29, 1413; d) B. Y. Lee, Y. K. Chung, N. Jeong, Y. Lee, S. E. Hwang, *J. Am. Chem. Soc.* **1994**, 116, 8793; e) N. Y. Lee, Y. K. Chung, *Tetrahedron Lett.* **1996**, 37, 3145; f) N. Jeong, S. H. Hwang, Y. W. Lee, J. S. Lim, *J. Am. Chem. Soc.* **1997**, 119, 10549; g) T. Sugihara, M. Yamaguchi, *J. Am. Chem. Soc.* **1998**, 120, 10782; h) J. W. Kim, Y. K. Chung, *Synthesis* **1998**, 142; i) T. Sugihara, M. Yamaguchi, *Synlett* **1998**, 1384; j) T. Rajesh, M. Periasamy, *Tetrahedron Lett.* **1999**, 40, 817; k) S. W. Kim, S. U. Son, S. I. Lee, T. Hyeon, Y. K. Chung, *J. Am. Chem. Soc.* **2000**, 122, 1550; l) M. Hayashi, Y. Hashimoto, Y. Yamamoto, J. Usuki, K. Saigo, *Angew. Chem.* **2000**, 112, 645; *Angew. Chem. Int. Ed.* **2000**, 39, 631; m) N. Jeong, S. H. Hwang, *Angew. Chem.* **2000**, 112, 650; *Angew. Chem. Int. Ed.* **2000**, 39, 636; n) M. E. Krafft, L. V. R. Boñaga, *Angew. Chem.* **2000**, 112, 3822; *Angew. Chem. Int. Ed.* **2000**, 39, 3676; o) T. Shibata, K. Takagi, *J. Am. Chem. Soc.* **2000**, 122, 9852.
- [3] M. E. Krafft, *Tetrahedron Lett.* **1988**, 29, 999.
- [4] Review on directed chemical reactions: A. H. Hoveyda, D. A. Evans, G. C. Fu, *Chem. Rev.* **1993**, 93, 1307.
- [5] In investigating the stoichiometric PKR, several groups have exploited the directing group strategy for controlling the regio- and stereoselectivity. For example, see a) M. E. Krafft, C. A. Juliano, I. L. Scott, C. Wright, M. D. McEachin, *J. Am. Chem. Soc.* **1991**, 113, 1693; b) S. Fonquerna, A. Moyano, M. A. Pericàs, A. Riera, *J. Am. Chem. Soc.* **1997**, 119, 10225.
- [6] For the use of a removable directing group in metal-catalyzed reactions, see a) D. A. Evans, G. C. Fu, A. H. Hoveyda, *J. Am. Chem. Soc.* **1988**, 110, 6917; b) B. Breit, *Eur. J. Org. Chem.* **1998**, 1123; c) C. H. Jun, H. Lee, *J. Am. Chem. Soc.* **1999**, 121, 880; d) N. D. Buezo, J. C. de la Rosa, J. Priego, I. Alonso, J. C. Carretero, *Chem. Eur. J.* **2001**, 7, 3890; e) S. Ko, Y. Na, S. Chang, *J. Am. Chem. Soc.* **2002**, 124, 750.
- [7] a) J. Yoshida, K. Itami, K. Mitsudo, S. Suga, *Tetrahedron Lett.* **1999**, 40, 3403; b) K. Itami, K. Mitsudo, T. Kamei, T. Koike, T. Nokami, J. Yoshida, *J. Am. Chem. Soc.* **2000**, 122, 12013; c) K. Itami, T. Nokami, J. Yoshida, *J. Am. Chem. Soc.* **2001**, 123, 5600; d) K. Itami, T. Koike, J. Yoshida, *J. Am. Chem. Soc.* **2001**, 123, 6957; e) K. Itami, T. Kamei, J. Yoshida, *J. Am. Chem. Soc.* **2001**, 123, 8773; f) K. Itami, T. Nokami, Y. Ishimura, K. Mitsudo, T. Kamei, J. Yoshida, *J. Am. Chem. Soc.* **2001**, 123, 11577; g) K. Itami, K. Mitsudo, A. Nishino, J. Yoshida, *Chem. Lett.* **2001**, 1088.
- [8] Reviews: a) J. Yoshida, K. Itami, *J. Synth. Org. Chem. Jpn.* **2001**, 59, 1086; b) K. Itami, K. Mitsudo, T. Nokami, T. Kamei, T. Koike, J. Yoshida, *J. Organomet. Chem.* **2002**, 653, 105.
- [9] For recent theoretical studies, see a) M. Yamanaka, E. Nakamura, *J. Am. Chem. Soc.* **2001**, 123, 1703; b) F. Robert, A. Milet, Y. Gimbert, D. Konya, A. E. Greene, *J. Am. Chem. Soc.* **2001**, 123, 5396; c) T. J. M. de Bruin, A. Milet, F. Robert, Y. Gimbert, A. E. Greene, *J. Am. Chem. Soc.* **2001**, 123, 7184.
- [10] K. Itami, K. Mitsudo, J. Yoshida, *Angew. Chem.* **2001**, 113, 2399; *Angew. Chem. Int. Ed.* **2001**, 40, 2337.
- [11] a) K. Itami, K. Mitsudo, J. Yoshida, *J. Org. Chem.* **1999**, 64, 8709 (oxidation); b) ref. [7c] (Pd-catalyzed cross-coupling reaction); c) ref. [7f] (protodesilylation); d) K. Itami, T. Nokami, J. Yoshida, *Tetrahedron* **2001**, 57, 5045 (electrophile-induced substitution reac-

- tion); e) ref. [7e] (fluoride-catalyzed reaction with carbonyl compounds).
- [12] $[\text{Co}_2(\text{CO})_8]$ also catalyzed this process, albeit in lower yield.
- [13] For the use of $[\text{Ru}_3(\text{CO})_{12}]$ as a catalyst in the intramolecular PKR and related reactions, see a) T. Morimoto, N. Chatani, Y. Fukumoto, S. Murai, *J. Org. Chem.* **1997**, 62, 3762; b) T. Kondo, N. Suzuki, T. Okada, T. Mitsudo, *J. Am. Chem. Soc.* **1997**, 119, 6187; c) N. Chatani, T. Morimoto, Y. Fukumoto, S. Murai, *J. Am. Chem. Soc.* **1998**, 120, 5335; d) S. K. Kang, K. J. Kim, Y. T. Hong, *Angew. Chem.* **2002**, 114, 1654; *Angew. Chem. Int. Ed.* **2002**, 41, 1584.
- [14] The silylated cyclopentenone was not observed in the reaction mixture. We assume that desilylation might occur by hydrolysis of the enolsilane, which is produced by the [1,3]-silyl migration of 5-silyl-2-cyclopentenone. Similar desilylation was observed during the course of our investigation of Stille coupling reactions using 2-PyMe₂-SiCH₂SnBu₃.^[7e] The Pd-catalyzed coupling reaction of 2-PyMe₂-SiCH₂SnBu₃ with acid chloride produced the corresponding methyl ketone [Eq. (5), unpublished results]. Moreover, we have also



observed that the oxophilicity of the 2-PyMe₂Si group was substantially higher than that of the PhMe₂Si group.^[11d]

- [15] Very recently, we learned of the results from Pagenkopf and co-workers on the stoichiometric intramolecular PKR of silicon-tethered enynes, which include vinylsilane moieties: J. F. Reichwein, S. T. Iacono, M. C. Patel, B. L. Pagenkopf, *Tetrahedron Lett.* **2002**, 43, 3739.
- [16] The development of nongaseous ethylene equivalents has been an active area of research. For example, see a) W. J. Kerr, M. McLaughlin, P. L. Pauson, S. M. Robertson, *Chem. Commun.* **1999**, 2171; b) W. J. Kerr, M. McLaughlin, P. L. Pauson, S. M. Robertson, *J. Organomet. Chem.* **2001**, 630, 104.
- [17] D. Huang, J. C. Huffman, J. C. Bollinger, O. Eisenstein, K. G. Caulton, *J. Am. Chem. Soc.* **1997**, 119, 7398.
- [18] The β -substituted vinylsilanes can be prepared with virtually complete regio- and stereoselectivity by the Pt/PtBu₃-catalyzed hydrosilylation of alkynes with 2-PyMe₂SiH: K. Itami, K. Mitsudo, A. Nishino, J. Yoshida, *J. Org. Chem.* **2002**, 67, 2645.
- [19] G. Fráter, J. A. Bajgrowicz, P. Kraft, *Tetrahedron* **1998**, 54, 7633, and references therein.
- [20] Cyclopentenone **6** has been prepared in 25% yield by the stoichiometric PKR of the 5- $[\text{Co}_2(\text{CO})_6]$ complex and ethylene gas (35 atm): D. C. Billington, I. M. Helps, P. L. Pauson, W. Thomson, D. Willison, *J. Organomet. Chem.* **1988**, 354, 233.
- [21] B. M. Trost, A. B. Pinkerton, *J. Org. Chem.* **2001**, 66, 7714.
- [22] G. H. Posner, M. J. Chapdelaine, C. M. Lentz, *J. Org. Chem.* **1979**, 44, 3661.
- [23] R. Noyori, M. Suzuki, *Angew. Chem.* **1984**, 96, 854; *Angew. Chem. Int. Ed. Engl.* **1984**, 23, 847.