

In the Communication by **D. M. Hodgson** et al. in Issue 22, **2002**, pp. 4313–4316 errors were inadvertently introduced into Tables 1 and 2 during the printing. The correct sections of the tables are now given below. The editorial office apologizes for these errors.

Table 1. Formation of enantioenriched diols.

Entry <sup>[a]</sup>	Epoxide	RLi/ligand	Product <sup>[b]</sup>	Yield [%] <sup>[c]</sup>	ee [%] <sup>[d]</sup>
1		<i>n</i> BuLi/ <b>1</b>		16	43
2		<i>n</i> BuLi/ <b>2</b>		44	–42
3		<i>n</i> BuLi/ <b>1</b>		46	34
4		<i>n</i> BuLi/ <b>2</b>		34	–40
5		<i>i</i> PrLi/ <b>1</b>		34	63
6		<i>n</i> BuLi/ <b>1</b>		57	27
7		<i>i</i> PrLi/ <b>1</b>		49	59
8		<i>n</i> BuLi/ <b>1</b>		50	51
9		<i>i</i> PrLi/ <b>1</b>		44	74
10		<i>i</i> PrLi/ <b>1</b>		42	56

[a] Entries 1–10 carried out in Et<sub>2</sub>O; entries 11–13 carried out in cumene. [b] Absolute configuration of predominant enantiomer obtained with (–)-sparteine (**1**) is shown (assigned by analogy with the sense of asymmetric induction in deprotonations of other *meso*-epoxides).<sup>[3]</sup> Bisoxazoline **2** preferentially produced the opposite configuration. [c] Yield of isolated product. [d] Determined by HPLC on a chiral stationary phase. TBDMS = *tert*-butyldimethylsilyl.

Table 2. Formation of enantioenriched amino alcohols.

Entry <sup>[a]</sup>	Epoxide	RLi/ligand	Product <sup>[b]</sup>	Yield [%] <sup>[c]</sup>	ee [%] <sup>[d]</sup>
9		<i>n</i> BuLi/ <b>1</b>		83	41
10		<i>n</i> BuLi/ <b>2</b>		60	–67
11		<i>i</i> PrLi/ <b>1</b>		59	65
12		<i>n</i> BuLi/ <b>1</b>		68	79
13		<i>n</i> BuLi/ <b>1</b>		84	66
14		<i>n</i> BuLi/ <b>2</b>		71	–67
15		<i>i</i> PrLi/ <b>1</b>		85	71
16		<i>i</i> PrLi/ <b>2</b>		72	–75
17		<i>i</i> PrLi/ <b>1</b>		69	82
18		TMSCH <sub>2</sub> Li/ <b>1</b>		61	57
19		mixed <sup>[e]</sup> / <b>1</b>		66 <sup>[f]</sup>	64

[a] Quenched at 25°C, apart from entries 9–11 and 18, 19 (–5°C), entry 12 (–78°C), entries 13–16 (–30°C), and entry 17 (–50°C). [b] See Table 1 footnote [b]. [c] Yield of isolated product. [d] Determined by HPLC or GC on a chiral stationary phase. [e] *i*PrLi (1.1 equiv) and TMSCH<sub>2</sub>Li (2.5 equiv). [f] Allylsilane major product, 7% of *i*Pr incorporation additionally observed. TMS = trimethylsilyl.