


Click Intermediate

Isolation of a Copper(I) Triazolide: A “Click” Intermediate**

Christoph Nolte, Peter Mayer, and Bernd F. Straub*

Dedicated to Professor Herbert Mayr on the occasion of his 60th birthday


The reaction of alkynes with organoazides has been known as a valuable triazole synthesis for more than a century.^[1] It is a typical example of a Huisgen-type 1,3-dipolar cycloaddition.^[2] The low regioselectivity and the high reaction temperatures, however, limit the scope of this reaction. Independently, the groups of Meldal and of Sharpless reported an ideal regioselectivity and an extreme rate enhancement in the reaction of terminal alkynes with azides in the presence of copper metal precatalysts or copper salts (Scheme 1).^[3] This method has been categorized as a “click” reaction.^[4]



Scheme 1. Concerted Huisgen-type [3+2] cycloaddition (top)^[5a] and copper-catalyzed click reaction (bottom).^[5b]


A stepwise pathway with copper(I) acetylide catalysts and a copper(I) triazolide intermediate is generally assumed (Scheme 2).^[6] Mechanistic investigations by Fokin and Finn revealed that the alkyne–azide click reaction presumably involves dinuclear copper(I) complexes in the rate-determining step.^[7]

Since a direct proof of the existence of copper(I) intermediates has remained elusive to date,^[8] we considered the isolation and characterization of a molecular copper(I) triazolide complex as a formidable goal in our attempts to unravel the mechanisms of organic reactions catalyzed by transition-metal complexes.^[9] Our strategy for the isolation of



Scheme 2. Some proposed mechanistic steps in the alkyne–azide click reaction; [Cu] represents either mononuclear or dinuclear copper(I) complex fragments.

a triazolide complex comprised the stoichiometric reaction of a copper(I) acetylide complex with a sterically demanding organoazide. In the absence of electrophiles in an aprotic solvent, the resulting copper triazolide species should be stable. However, the preparation of mononuclear copper(I) phenyl acetylides with phenanthroline or phosphane ancillary ligands failed. In our attempts, these ligands dissociated from copper, resulting in precipitation of polymeric copper phenyl acetylide. Since molecular copper(I) acetylide complexes with unsaturated N-heterocyclic carbene (NHC) ligands had already been described,^[10] we developed a straightforward synthesis to copper(I) phenyl acetylide complexes with a saturated NHC ligand (Scheme 3).^[11] NHC copper(I) halide complexes have only recently been reported as highly active catalysts for triazole click reactions.^[12]



Scheme 3. Preparation of a copper(I) triazolide complex 3.

[*] C. Nolte, Dr. P. Mayer, Dr. B. F. Straub^[†]
Department Chemie und Biochemie, LMU München
Butenandtstrasse 5–13 (Haus F), 81377 München (Germany)
E-mail: Bernd.F.Straub@cup.uni-muenchen.de

[†] temporary address:
Chemisch-Geowissenschaftliche Fakultät
Institut für Organische Chemie und Makromolekulare Chemie
Friedrich-Schiller-Universität Jena
Humboldtstrasse 10, 07743 Jena (Germany)

[**] The authors gratefully acknowledge support by the DFG, the LMU München, the Friedrich-Schiller-Universität Jena, the Dr. Otto Röhmer-Gedächtnisstiftung, and Prof. Dr. Herbert Mayr.

Supporting information (X-ray structure of 1, preparation procedures and characterization data) for this article is available on the WWW under <http://www.angewandte.org> or from the author.

Indeed, the reaction of the copper(I) acetate complex **1** with lithium phenyl acetylide yields a suitable precursor complex **2**. Addition of azidodi-4-tolylmethane to a solution of acetylide **2** in toluene results in the precipitation of a mononuclear copper triazolidine complex **3** at room temperature within 12 h. It thus appears that dinuclear copper complexes are not mandatory for Huisgen–Sharpless click reactions, since even mononuclear copper(I) acetylides with extreme steric shielding react with bulky organoazides at room temperature.^[13] The reaction rate is comparable to sterically hindered systems of Nolan and co-workers.^[12]

Both the acetate complex **1** as well as the triazolidine complex **3** are active catalysts in click chemistry. Preliminary results indicate a strong dependence of the reaction rate on the acidity of the medium. The rate-limiting step apparently changes from the protonation of the triazolidine complex (H₂O or alkyne as proton source) to the alkyne deprotonation–cycloaddition sequence (excess of acetic acid). Triazolidine complex **3** in CD₂Cl₂ reacts quantitatively with acetic acid within minutes at room temperature to give the triazole product and acetate complex **1**.

The crystal structure of the product **3** shows a d¹⁰ ML₂ copper(I) species with a linear metal coordination environment (Figure 1).^[14] The uncharged complex is, electronically, closely related to diorganocuprates with 14 valence electrons.^[15] In triazolidine **3**, the Cu–C bond lengths to the formally uncharged NHC ligand and to the formally anionic triazolidine ligand are essentially equal. The five-membered heterocyclic ligands are coplanar, leading to a conformation with minimal steric repulsion. An *ortho* position of the phenyl substituent is




Figure 1. X-ray structure of copper(I) triazolidine **3**.^[16,17] Disordered CH₂Cl₂ is omitted for clarity; only one of the two complexes **3** in the elemental cell is shown. Average bond lengths [pm] and angles [°] (deviations): Cu–C1 190.9(0.5), Cu–C2 190.4(0.2), Cu–H(*o*-Ph) 269.1(7.2), C2–C3 139.8(0.3), C3–N3 136.7(0.1), C2–N5 136.8(0.2), N3–N4 131.2(0.5), N4–N5 136.1(0.1), N2–C1 133.2(0.4), N1–C1 134.1(0.1); C1–Cu–C2 178.0(0.3), Cu–C2–N5 125.3(3.1), Cu–C2–C3 133.7(3.2), C2–C3–C4 130.7(0.3), C2–N5–C5 128.9(0.8). Dihedral angles: N2–C1–C2–C3 1.7(0.2), Cu–C2–N5–N4 178.2(1.9).

in close vicinity to the copper atom, resulting in the Cu–C2–C3 angle being larger than the Cu–C2–N5 angle.

In Figure 2, the ¹H NMR spectrum of copper triazolidine complex **3** in CD₂Cl₂ is shown. The thermal stability of




Figure 2. ¹H NMR spectrum of **3**.

complex **3** in solution is consistent with its high melting point and decomposition temperature of 235 °C. The complex even resists fragmentation under ESI conditions. For the triazolidine substitution pattern present in complex **3**, we have not observed N₂ elimination.

In summary, the isolation of a molecular copper(I) triazolidine complex is direct evidence for the existence of such structures, which have to date only been assumed as intermediates in the Huisgen–Sharpless click reaction. With click intermediates at hand, mechanistic studies can go beyond indirect kinetic measurements of reaction mixtures towards the direct investigation of elementary steps.

Experimental Section

Azidodi(4-tolyl)methane (0.385 g, 1.62 mmol) was added to a Schlenk flask charged with **2** (0.600 g, 1.08 mmol) and dry toluene (10 mL) under an N₂ atmosphere. The solution was stirred at room temperature. After 4 h, a colorless solid began to precipitate. After an additional 8 h, the precipitate was collected by filtration and washed with toluene. The solid was dissolved in CH₂Cl₂ and filtered through Celite 500. Pentane was added to the filtrate, and the solution cooled to –20 °C. The precipitate was collected and dried in vacuo, affording 414 mg (48.4%) of the product as colorless crystals. Single crystals of **3** suitable for an X-ray diffraction study were grown from CH₂Cl₂ at room temperature by slow diffusion of pentane. ¹H NMR (399.92 MHz, CD₂Cl₂): δ = 1.13 (d, ³J_{HH} = 7 Hz, 12 H, CH(CH₃)₂), 1.29 (d, ³J_{HH} = 7 Hz, 12 H, CH(CH₃)₂), 2.29 (s, 6 H, Aryl-CH₃), 3.10 (sept^{pseudo}, ³J_{HH} = 7 Hz, 4 H, CH(CH₃)₂), 3.98 (s, 4 H, NCH₂), 5.72 (s, 1 H, Tol₂CH), 6.64 (d, ³J_{HH} = 8.0 Hz, 4 H, *o*-Tol), 6.86 (t^{pseudo}t^{pseudo}, ³J_{HH} = 7.2 Hz, ⁴J_{HH} = 1.2 Hz, 2 H, *m*-C₆H₅), 6.94 (tt, ³J_{HH} = 7.2 Hz, ⁴J_{HH} = 1.2 Hz, 1 H, *p*-C₆H₅), 6.99 (d, ³J_{HH} = 8.0 Hz, 4 H, *m*-Tol), 7.27 (d, ³J_{HH} = 8.0 Hz, 4 H, *m*-NHC), 7.35 (dd, ³J_{HH} = 7.2 Hz, ⁴J_{HH} = 1.6 Hz, 2 H, *o*-C₆H₅), 7.45 ppm (t, ³J_{HH} = 8.0 Hz, 2 H, *p*-NHC). ¹³C{¹H} NMR (100.56 MHz, CD₂Cl₂): δ = 20.8 (Aryl-CH₃), 23.6 (CHCH₃), 25.1 (CHCH₃), 29.0 (CHCH₃), 54.2 (NCH₂), 70.0 (Tol₂CH), 124.7 (*p*-C₆H₅), 125.0 (*m*-NHC), 125.3 (*o*-C₆H₅), 128.2 (*m*-C₆H₅), 128.2 (*m*-

Tol), 128.9 (s, *o*-Tol) 130.0 (s, *p*-NHC), 135.0 (*o*-NHC), 136.3 (*ipso*-C₆H₅), 136.7 (*p*-Tol), 138.8 (*ipso*-Tol), 146.9 (*ipso*-NHC), 152.2 (triazolide-C), 154.6 (triazolide-C), 204.7 ppm (N₂CCu). MS (+p ESI): *m/z* (%) = 792.406 (100.00) [*M*⁺(⁶³Cu)+H], 793.411 (45.50) [*M*⁺(⁶³Cu)+2H]. IR: $\tilde{\nu}$ = 3057 (w), 2962 (s), 2924 (s), 2868 (m), 2097 (w), 1602 (w), 1513 (m), 1481 (m), 1459 (s), 1428 (s), 1385 (m), 1364 (w), 1327 (m), 1274 (s), 1182 (m), 1058 (w), 988 (m), 805 (s), 788 (m), 770 (s), 758 cm⁻¹ (s). M.p. 235 °C (decomp). Elemental analysis (%) calcd for C₅₀H₅₈CuN₅: C 75.77, H 7.38, N 8.84; found: C 75.24, H 7.33, N 8.73. CCDC-625315 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

Received: October 30, 2006

Revised: January 3, 2007

Published online: February 15, 2007

Keywords: alkynes · azides · click chemistry · copper · cycloadditions · reaction mechanisms

- [1] A. Michael, *J. Prakt. Chem.* **1893**, 48, 94.
- [2] R. Huisgen, *Angew. Chem.* **1963**, 75, 604.
- [3] a) V. V. Rostovtsev, L. G. Green, V. V. Fokin, K. B. Sharpless, *Angew. Chem.* **2002**, 114, 2708; *Angew. Chem. Int. Ed.* **2002**, 41, 2596; b) C. W. Tornøe, C. Christensen, M. Meldal, *J. Org. Chem.* **2002**, 67, 3057.
- [4] K. B. Sharpless, M. G. Finn, *Angew. Chem.* **2001**, 113, 11, 2056; *Angew. Chem. Int. Ed.* **2001**, 40, 2004.
- [5] a) W. Kirmse, L. Horner, *Justus Liebigs Ann. Chem.* **1958**, 614, 1; b) L. D. Pachon, J. H. van Maarseveen, G. Rothenberg, *Adv. Synth. Catal.* **2005**, 347, 811.
- [6] V. D. Bock, H. Hiemstra, J. H. van Maarseveen, *Eur. J. Org. Chem.* **2006**, 51.
- [7] a) F. Himo, T. Lovell, R. Hilgraf, V. V. Rostovtsev, L. Noodleman, K. B. Sharpless, V. V. Fokin, *J. Am. Chem. Soc.* **2005**, 127, 210; b) V. O. Rodionov, V. V. Fokin, M. G. Finn, *Angew. Chem.* **2005**, 117, 2250; *Angew. Chem. Int. Ed.* **2005**, 44, 2210.
- [8] For trapping of assumed copper(I) triazolide intermediates by electrophiles, see a) B. Gerard, J. Ryan, A. B. Beeler, J. A. Porco, Jr., *Tetrahedron* **2006**, 62, 6405; b) B. H. M. Kuipers, G. C. T. Dijkmans, S. Groothuys, P. J. L. M. Quaedflieg, R. H. Blaauw, F. L. van Delft, F. P. J. T. Rutjes, *Synlett* **2005**, 3059; c) J. Deng, Y.-M. Wu, Q.-Y. Chen, *Synthesis* **2005**, 2730.
- [9] a) B. F. Straub, P. Hofmann, *Angew. Chem.* **2001**, 113, 1328; *Angew. Chem. Int. Ed.* **2001**, 40, 1288; b) B. F. Straub, F. Rominger, P. Hofmann, *Organometallics* **2000**, 19, 4305; c) B. F. Straub, I. Gruber, F. Rominger, P. Hofmann, *J. Organomet. Chem.* **2003**, 684, 124; d) B. F. Straub, *Angew. Chem.* **2005**, 117, 6129; *Angew. Chem. Int. Ed.* **2005**, 44, 5974.
- [10] L. A. Goj, E. D. Blue, C. Munro-Leighton, J. L. Petersen, T. B. Gunnoe, *Inorg. Chem.* **2005**, 44, 8647.
- [11] a) N. P. Mankad, T. G. Gray, D. S. Laitar, J. P. Sadighi, *Organometallics* **2004**, 23, 1191; b) A. J. Arduengo III, R. Krafzyk, R. Schmutzler, H. A. Craig, J. R. Goerlich, W. J. Marshall, M. Unverzagt, *Tetrahedron* **1999**, 55, 14523; c) G. A. Grasa, M. S. Viciu, J. Huang, S. P. Nolan, *J. Org. Chem.* **2001**, 66, 7729; d) L. Jafarpour, E. D. Stevens, S. P. Nolan, *J. Organomet. Chem.* **2000**, 606, 49.
- [12] S. D. Díez-González, A. Correa, L. Cavallo, S. P. Nolan, *Chem. Eur. J.* **2006**, 12, 7558.
- [13] Cross experiments are in progress to rule out fast, reversible NHC dissociation, acetylide ligand exchange, and formation of dinuclear complexes.
- [14] With a π -acceptor ligand instead of a strong donor ligand, such as an N-heterocyclic carbene, a coordination number of 3 at a 16-valence-electron copper complex would certainly be a viable alternative for intermediates in click catalysis.
- [15] a) M. John, C. Auel, C. Behrens, M. Marsch, K. Harms, F. Bosold, R. M. Gschwind, P. R. Rajamohanam, G. Boche, *Chem. Eur. J.* **2000**, 6, 3060; b) G. Boche, F. Bosold, M. Marsch, K. Harms, *Angew. Chem.* **1998**, 110, 1779; *Angew. Chem. Int. Ed.* **1998**, 37, 1684.
- [16] a) SIR97: A. Altomare, M. C. Burla, M. Camalli, G. L. Cascarano, C. Giacovazzo, A. Guagliardi, A. G. G. Moliterni, G. Polidori, R. Spagna, *J. Appl. Crystallogr.* **1999**, 32, 115; b) SHELXL-97 (G. M. Sheldrick, University of Göttingen, 97-2 version).
- [17] a) Ortep-3 for Windows: L. J. Farrugia, *J. Appl. Crystallogr.* **1997**, 30, 565; b) <http://www.povray.org>.